Title: Konstantin's Group

Date: Aug 27, 2004 08:45 AM

URL: http://pirsa.org/04080007

Abstract:

Electro-magnetic force

$E_{photon} \cdot \lambda_{photon}$

$$= \frac{e^2}{4\pi\varepsilon_0} \cdot \frac{1}{hc}$$

$$\alpha = \frac{e^2}{} \approx \frac{1}{137}$$

Electro-magnetic force

$$E_{photon} \cdot \lambda_{photon}$$

$$= \frac{e^2}{4\pi\varepsilon_0} \cdot \frac{1}{hc}$$


Electro-magnetic force
$E_{photon} \cdot \lambda_{photon}$
$= \frac{e^2}{4\pi\varepsilon_0} \cdot \frac{1}{hc}$
$\alpha = \frac{e^2}{\hbar c} \approx \frac{1}{137}$

Electro-magnetic force

$E_{photon} \cdot \lambda_{photon}$

$$= \frac{e^2}{4\pi\varepsilon_0} \cdot \frac{1}{hc}$$

$$\alpha = \frac{e^2}{\hbar c} \approx \frac{1}{137}$$

$$E_n = \frac{m_e c^2 \alpha^2}{2n^2}$$

$$E_n = \frac{m_e c^2 \alpha^2}{2n^2}$$


Ionization energy
$E_n = -m_e c \alpha / n^2$
α
OH^- H^+
$Fe^{3+} + 3e^- = Fe$
H
H-H
O=O
C - H
H
E
PE = mgxdxE
E = PE+KE
mv^2
H
H
O
E


$F = \frac{\alpha}{R^2}$
$E_n = -m_e c^2 \alpha^2 \frac{1}{n^2}$
Ionization energy
$\alpha \uparrow$
$F \propto \frac{1}{r^2} \uparrow$ OH^- H^+
$Fe^{3+} + 3e^- =$
H—C—H
H
Pirsa: 04080007
Page 10/25

the de Broglie wavelength of an electron.

$\mathcal{O}$ = orbit diameter

$\mathcal{O} = n\lambda ; n \in \mathbb{N} \rightarrow \lambda = \frac{h}{p}$.

$\mathcal{O} = \frac{nh}{p} \longrightarrow p = mv$.

$$\boxed{\mathcal{O} = \frac{nh}{mv}}$$

$$\boxed{E_n = -m_e c^2 \alpha^2 \frac{1}{n^2}}$$

$$E = \frac{1}{2}mv^2 - k\frac{e^2}{r}$$

in an orbital $F_c = F_e$

$$\frac{mv^2}{r} = k\frac{e^2}{r^2}$$

$$mv^2 = k\frac{e^2}{r}$$

$$-\cancel{m_e} c^2 \alpha^2 \frac{1}{n^2} = -\frac{1}{2}\cancel{m}v^2$$

$$v^2 = \frac{2c^2\alpha^2}{n^2}$$

$$\boxed{v = \pm\sqrt{2}\frac{c\alpha}{n}}$$

we know $\mathcal{O} \propto \frac{1}{v}$

$v \propto \alpha$

$\therefore \mathcal{O} \propto \frac{1}{\alpha}$

$\alpha \propto e^2$

How fundamental charge effects atomic radii

orbit diameters are whole number multiples of the de Broglie wavelength of an electron.

$\mathcal{O}$ = orbit diameter

$\mathcal{O} = n\lambda : n \in \mathbb{N} \rightarrow \lambda = \frac{h}{p}$.

$\mathcal{O} = \frac{nh}{p} \longrightarrow p = mv$.

$$\boxed{\mathcal{O} = \frac{nh}{mv}}$$

$$\boxed{E_n = -m_e c^2 \alpha^2 \frac{1}{n^2}}$$

$E = \frac{1}{2}mv^2 - k\frac{e^2}{r}$

$-\cancel{m}c^2\alpha^2\frac{1}{n^2} = -\frac{1}{2}\cancel{m}v^2$

$v^2 = \frac{2c^2\alpha^2}{n^2}$

$$\boxed{v = \pm\sqrt{2}\frac{c\alpha}{n}}$$

we know $\mathcal{O} \propto \frac{1}{v}$

σ = orbit diameter

$\sigma = n\lambda : n \in \mathbb{N} \rightarrow \lambda = \frac{h}{p}$

$\sigma = \frac{nh}{p} \longrightarrow p = mv$

$$\boxed{\sigma = \frac{nh}{mv}}$$

$$\boxed{E_n = -m_e c^2 \alpha^2 \frac{1}{n^2}}$$

$$E = \frac{1}{2}mv^2 - k\frac{e^2}{r}$$

in an orbital $F_c = F_e$

$$\frac{mv^2}{r} = k\frac{e^2}{r^2}$$

$$mv^2 = k\frac{e^2}{r}$$

$$E = \frac{1}{2}mv^2 - mv^2$$

$$-mc^2\alpha^2\frac{1}{n^2} = -\frac{1}{2}mv^2$$

$$v^2 = \frac{2c^2\alpha^2}{n^2}$$

$$\boxed{v = \pm\sqrt{2}\frac{c\alpha}{n}}$$

we know $\sigma \propto \frac{1}{v}$

$v \propto \alpha$

$\therefore \sigma \propto \frac{1}{\alpha}$

$\alpha \propto e^2$

$\sigma \propto e^2$

orbit diameters are whole number multiples of the de Broglie wavelength of an electron.

σ = orbit diameter

$\sigma = n\lambda : n \in N \rightarrow \lambda = \frac{h}{p}$.

$\sigma = \frac{nh}{p} \longrightarrow p = mv$.

$$\boxed{\sigma = \frac{nh}{mv}}$$

$$\boxed{E_n = -m_e c^2 \alpha^2 \frac{1}{n^2}}$$

$$E = \frac{1}{2}mv^2 - k\frac{e^2}{r}$$

in an orbital $F_c = F_e$

$$\frac{mv^2}{r} = k\frac{e^2}{r^2}$$

$$mv^2 = k\frac{e^2}{r}$$

$$-m_e c^2 \alpha^2 \frac{1}{n^2} = -\frac{1}{2}mv^2$$

$$v^2 = \frac{2c^2\alpha^2}{n^2}$$

$$\boxed{v = \pm\sqrt{2}\frac{c\alpha}{n}}$$

we know $\sigma \propto \frac{1}{v}$

$v \propto \alpha$

$\therefore \sigma \propto \frac{1}{\alpha}$

to electrostatic forces between quarks (r^2)

- Since the charge of protons (quarks) varies with α, this changes mass of proton and neutron

because $\alpha = \frac{e^2}{\hbar c} = e^2$

proton: two up quarks and a down

up $= -\frac{2}{3}$

down $= \frac{1}{3}$



neutron: two down quarks and an up



27 August 200

Alpha and the nucleon

- A great deal of the mass of nucleons is due to electrostatic forces between quarks ($F = K\frac{Qq}{r^2}$)

- Since the charge of protons (quarks) varies with α, this changes mass of proton and neutron

because $\alpha = \frac{e^2}{\hbar c} = e^2$

proton: two up quarks and a down

up $= -\frac{2}{3}$

down $= \frac{1}{3}$



because $\alpha = \frac{e^2}{\hbar c} = e^2$

proton: two up quarks and a down

up $= -\frac{2}{3}$

down $= \frac{1}{3}$



neutron: two down quarks and an up



$E = \frac{e^2}{3d}$ related to alpha $\alpha = e^2$

Bigger alpha = higher E, higher nucleon mass

because $\alpha = \frac{e^2}{\hbar c} = e^2$

proton: two up quarks and a down

up $= -\frac{2}{3}$

down $= \frac{1}{3}$



neutron: two down quarks and an up



$\frac{e^2}{J}$ related to alpha $\alpha = e^2$

ha = higher E, higher nucleon mass

α, this changes mass of proton and electron

because $\alpha \to \frac{e^2}{\hbar c} \to e^2$

proton: two up quarks and a down up $= -\frac{2}{3}$, down $= \frac{1}{3}$

neutron: two down quarks and an up



$F = \frac{e^2}{r^2}$ related to alpha $\alpha \propto e^2$



neutron: two down quarks and an up



$E = \frac{e^2}{3d}$ related to alpha $\alpha = e^2$

Bigger alpha = higher E, higher nucleon mass

Smaller alpha = smaller E, lower nucleon mass


Iron
Binding Energy / nuclei
Element #


Iron
Binding Energy / nucleon
Element #


BIRTHDAY
KONSTANTI


BIRTHD
KONSTA