

Title: Supersymmetric Black Rings and Supertubes

Date: Sep 30, 2004 01:00 PM

URL: <http://pirsa.org/04090004>

Abstract:

Supersymmetric Black Rings and Supertubes

David Mateos

Perimeter Institute

Based on:

A Supersymmetric Black Ring, hep-th/0407065;
Supersymmetric Black Rings and Three-charge Supertubes, hep-th/0408120,

with H. Elvang, R. Emparan, and H. Reall

Supertubes, hep-th/0103030,

with P. Townsend

Supergravity Supertubes, hep-th/0106012,

with R. Emparan and P. Townsend

Tachyons, Supertubes and Brane/Anti-Brane Systems, hep-th/0112054,

with S. Ng and P. Townsend

Supersymmetric Black Ring = Supersymmetric,




Supersymmetric Black Ring = Supersymmetric,
Asymptotically Flat,
Stationary Black Hole Solution of 5D Supergravity



Supersymmetric Black Ring = Supersymmetric,
Asymptotically Flat,
Stationary Black Hole Solution of 5D Supergravity
with Horizon Topology $S^1 \times S^2$



Supersymmetric Black Ring = Supersymmetric,
Asymptotically Flat,
Stationary Black Hole Solution of 5D Supergravity
with Horizon Topology $S^1 \times S^2$



Times flat directions = Black **Supertube**

2-charge Supertubes: DBI (Worldvolume) Description

Townsend & D.M.

Supersymmetric Brane Expansion in Flat Space by Angular Momentum




$$\Gamma_{F1} \varepsilon = \varepsilon$$

$$\Gamma_{D0} \varepsilon = \varepsilon$$

2-charge Supertubes: DBI (Worldvolume) Description

Townsend & D.M.

Supersymmetric Brane Expansion in Flat Space by Angular Momentum



$$\Gamma_{F1} \varepsilon = \varepsilon$$


$$\Gamma_{D0} \varepsilon = \varepsilon$$

2-charge Supertubes: DBI (Worldvolume) Description

Townsend & D.M.

Supersymmetric Brane Expansion in Flat Space by Angular Momentum

1/4-SUSY preserved




2-charge Supertubes: DBI (Worldvolume) Description

Townsend & D.M.

Supersymmetric Brane Expansion in Flat Space by Angular Momentum

1/4-SUSY preserved

Q_{F1} and Q_{D0} dissolved as fluxes




2-charge Supertubes: DBI (Worldvolume) Description

Townsend & D.M.

Supersymmetric Brane Expansion in Flat Space by Angular Momentum

1/4-SUSY preserved

Q_{F1} and Q_{D0} dissolved as fluxes
 J generated as integrated **Poynting**



**Tubular D2/ $F1$ /D0
Bound State**

$$\Gamma_{F1} \varepsilon = \varepsilon$$

$$\Gamma_{D0} \varepsilon = \varepsilon$$


2-charge Supertubes: DBI (Worldvolume) Description

Townsend & D.M.

Supersymmetric Brane Expansion in Flat Space by Angular Momentum

1/4-SUSY preserved

Q_{F1} and Q_{D0} dissolved as fluxes
 J generated as integrated **Poynting**
 $E = Q_{F1} + Q_{D0}$



**Tubular D2/ $F1$ /D0
Bound State**

$$\Gamma_{F1} \varepsilon = \varepsilon$$

$$\Gamma_{D0} \varepsilon = \varepsilon$$

2-charge Supertubes: DBI (Worldvolume) Description


Townsend & D.M.

Supersymmetric Brane Expansion in Flat Space by Angular Momentum

1/4-SUSY preserved

Q_{F1} and Q_{D0} dissolved as fluxes
 J generated as integrated **Poynting**
 $E = Q_{F1} + Q_{D0}$


Arbitrary Cross-section C in $\mathbb{E}^8$:
 (and charge densities)



2-charge Supertubes: DBI (Worldvolume) Description

Townsend & D.M.

Supersymmetric Brane Expansion in Flat Space by Angular Momentum




1/4-SUSY preserved

Q_{F1} and Q_{D0} dissolved as fluxes
 J generated as integrated **Poynting**
 $E = Q_{F1} + Q_{D0}$

Arbitrary Cross-section C in $\mathbb{E}^8$:
 (and charge densities)


TS-Dualizing = 'Helical' String with
 Left-moving wave on it



2-charge Supertubes: DBI (Worldvolume) Description

Townsend & D.M.

Supersymmetric Brane Expansion in Flat Space by Angular Momentum




1/4-SUSY preserved

Q_{F1} and Q_{D0} dissolved as fluxes
 J generated as integrated **Poynting**
 $E = Q_{F1} + Q_{D0}$

Arbitrary Cross-section C in $\mathbb{E}^8$:
 (and charge densities)


TS-Dualizing = 'Helical' String with
 Left-moving wave on it



No net D2-brane charge
 but dipole $q_{D2} \sim n_{D2}$


2-charge Supertubes: DBI (Worldvolume) Description

For circular Supertube of radius R :



$$R = \frac{\sqrt{Q_{F1} Q_{D0}}}{q_{D2}} \quad J = \frac{Q_{F1} Q_{D0}}{q_{D2}} = q_{D2} R^2$$

2-charge Supertubes: DBI (Worldvolume) Description




For circular Supertube of radius R :

$$R = \frac{\sqrt{Q_{F1}Q_{D0}}}{q_{D2}} \quad J = \frac{Q_{F1}Q_{D0}}{q_{D2}} = q_{D2}R^2$$

For general system with F1/D0 charges:

$$R \leq \frac{\sqrt{Q_{F1}Q_{D0}}}{q_{D2}} \quad J \leq \frac{Q_{F1}Q_{D0}}{q_{D2}}$$

2-charge Supertubes: DBI (Worldvolume) Description




For circular Supertube of radius R :

$$R = \frac{\sqrt{Q_{F1}Q_{D0}}}{q_{D2}} \quad J = \frac{Q_{F1}Q_{D0}}{q_{D2}} = q_{D2}R^2$$

For general system with F1/D0 charges:

$$R \leq \frac{\sqrt{Q_{F1}Q_{D0}}}{q_{D2}} \quad J \leq \frac{Q_{F1}Q_{D0}}{q_{D2}}$$

2-charge Supertubes: DBI (Worldvolume) Description




For circular Supertube of radius R :

$$R = \frac{\sqrt{Q_{F1}Q_{D0}}}{q_{D2}} \quad J = \frac{Q_{F1}Q_{D0}}{q_{D2}} = q_{D2}R^2$$

For general system with F1/D0 charges:

$$R \leq \frac{\sqrt{Q_{F1}Q_{D0}}}{q_{D2}} \quad J \leq \frac{Q_{F1}Q_{D0}}{q_{D2}}$$

2-charge Supertubes: DBI (Worldvolume) Description



For circular Supertube of radius R :

$$R = \frac{\sqrt{Q_{F1} Q_{D0}}}{q_{D2}} \quad J = \frac{Q_{F1} Q_{D0}}{q_{D2}} = q_{D2} R^2$$

For general system with F1/D0 charges:

$$R \leq \frac{\sqrt{Q_{F1} Q_{D0}}}{q_{D2}} \quad J \leq \frac{Q_{F1} Q_{D0}}{q_{D2}}$$

Or eliminating q_{D2} :

$$J^2 \leq R^2 Q_{F1} Q_{D0}$$

2-charge Supertubes: Supergravity Description

Emparan, Townsend & D.M.


$$ds^2 = -U^{-1}V^{-1/2}(dt + \omega)^2 + U^{-1}V^{1/2}dz^2 + V^{1/2}d\vec{x} \cdot d\vec{x}$$

$$B_2 = -U^{-1}(dt + \omega) \wedge dz$$

$$C_1 = -V^{-1}(dt + \omega)$$

$$C_3 = -U^{-1}dt \wedge dz \wedge \omega$$

$$e^\phi = U^{-1/2}V^{3/4}$$



2-charge Supertubes: Supergravity Description

Emparan, Townsend & D.M.

$$ds^2 = -U^{-1}V^{-1/2}(dt + \omega)^2 + U^{-1}V^{1/2}dz^2 + V^{1/2}d\vec{x} \cdot d\vec{x}$$


$$B_2 = -U^{-1}(dt + \omega) \wedge dz$$

$$C_1 = -V^{-1}(dt + \omega)$$

$$C_3 = -U^{-1}dt \wedge dz \wedge \omega$$

$$e^\phi = U^{-1/2}V^{3/4}$$

$U(x)$ and $V(x)$ harmonic functions on $\mathbb{E}^8$



2-charge Supertubes: Supergravity Description

Emparan, Townsend & D.M.

$$ds^2 = -U^{-1}V^{-1/2}(dt + \omega)^2 + U^{-1}V^{1/2}dz^2 + V^{1/2}d\vec{x} \cdot d\vec{x}$$

$$B_2 = -U^{-1}(dt + \omega) \wedge dz$$


$$C_1 = -V^{-1}(dt + \omega)$$

$$C_3 = -U^{-1}dt \wedge dz \wedge \omega$$

$$e^\phi = U^{-1/2}V^{3/4}$$

$U(x)$ and $V(x)$ harmonic functions on $\mathbb{E}^8$

1-form $\omega(x)$ satisfies Maxwell on $\mathbb{E}^8$: $d * d \omega = 0$



2-charge Supertubes: Supergravity Description

Emparan, Townsend & D.M.

$$ds^2 = -U^{-1}V^{-1/2}(dt + \omega)^2 + U^{-1}V^{1/2}dz^2 + V^{1/2}d\vec{x} \cdot d\vec{x}$$

$$B_2 = -U^{-1}(dt + \omega) \wedge dz$$

$$C_1 = -V^{-1}(dt + \omega)$$


$$C_3 = -U^{-1}dt \wedge dz \wedge \omega$$

$$e^\phi = U^{-1/2}V^{3/4}$$

$U(x)$ and $V(x)$ harmonic functions on $\mathbb{E}^8$

1-form $\omega(x)$ satisfies Maxwell on $\mathbb{E}^8$: $d * d \omega = 0$

$$U(\vec{x}) = 1 + Q_{F1} \int_C \frac{d\sigma}{|\vec{x} - \vec{x}(\sigma)|^6} \quad \omega(\vec{x}) = q_{D2} \int_C d\sigma \frac{x'_i(\sigma)}{|\vec{x} - \vec{x}(\sigma)|^6} dx^i$$



2-charge Supertubes: Supergravity Description

Emparan, Townsend & D.M.


$$ds^2 = -U^{-1}V^{-1/2}(dt + \omega)^2 + U^{-1}V^{1/2}dz^2 + V^{1/2}d\vec{x} \cdot d\vec{x}$$

$$B_2 = -U^{-1}(dt + \omega) \wedge dz$$

$$C_1 = -V^{-1}(dt + \omega)$$

$$C_3 = -U^{-1}dt \wedge dz \wedge \omega$$

$$e^\phi = U^{-1/2}V^{3/4}$$



$U(x)$ and $V(x)$ harmonic functions on $\mathbb{E}^8$

1-form $\omega(x)$ satisfies Maxwell on $\mathbb{E}^8$: $d * d \omega = 0$

$$U(\vec{x}) = 1 + Q_{F1} \int_C \frac{d\sigma}{|\vec{x} - \vec{x}(\sigma)|^6} \quad \omega(\vec{x}) = q_{D2} \int_C d\sigma \frac{x'_i(\sigma)}{|\vec{x} - \vec{x}(\sigma)|^6} dx^i$$

Net charges: $\int_{S^7_\infty} *H_3 = Q_{F1} \quad \int_{S^8_\infty} *G_2 = Q_{D0} \quad \mathbf{J} = q_{D2} \mathbf{R}^2$

2-charge Supertubes: Supergravity Description

Emparan, Townsend & D.M.


$$ds^2 = -U^{-1}V^{-1/2}(dt + \omega)^2 + U^{-1}V^{1/2}dz^2 + V^{1/2}d\vec{x} \cdot d\vec{x}$$

$$B_2 = -U^{-1}(dt + \omega) \wedge dz$$

$$C_1 = -V^{-1}(dt + \omega)$$

$$C_3 = -U^{-1}dt \wedge dz \wedge \omega$$

$$e^\phi = U^{-1/2}V^{3/4}$$



$U(x)$ and $V(x)$ harmonic functions on $\mathbb{E}^8$

1-form $\omega(x)$ satisfies Maxwell on $\mathbb{E}^8$: $d * d \omega = 0$

$$U(\vec{x}) = 1 + Q_{F1} \int_C \frac{d\sigma}{|\vec{x} - \vec{x}(\sigma)|^6} \quad \omega(\vec{x}) = q_{D2} \int_C d\sigma \frac{x'_i(\sigma)}{|\vec{x} - \vec{x}(\sigma)|^6} dx^i$$

Net charges: $\int_{S^7_\infty} *H_3 = Q_{F1} \quad \int_{S^8_\infty} *G_2 = Q_{D0} \quad \mathbf{J} = q_{D2} \mathbf{R}^2$


$$\mathbf{M} = Q_{F1} + Q_{D0} \longleftrightarrow \frac{1}{4}\text{-SUSY}$$

2-charge Supertubes: Supergravity Description

No net D2 charge, but D2 dipole (and higher) moments:

2-charge Supertubes: Supergravity Description


No net D2 charge, but D2 dipole (and higher) moments:



$$G_4 = d \left[\frac{J r^2}{(r^2 + \rho^2)^4} dt \wedge dz \wedge d\psi \right] + \dots$$

2-charge Supertubes: Supergravity Description

No net D2 charge, but D2 dipole (and higher) moments:




$$G_4 = d \left[\frac{J r^2}{(r^2 + \rho^2)^4} dt \wedge dz \wedge d\psi \right] + \dots$$

$$\longrightarrow \int_{S_\infty^6} \star G_4 = 0$$

2-charge Supertubes: Supergravity Description

No net D2 charge, but D2 dipole (and higher) moments:




$$G_4 = d \left[\frac{J r^2}{(r^2 + \rho^2)^4} dt \wedge dz \wedge d\psi \right] + \dots$$

$$\longrightarrow \int_{S_\infty^6} \star G_4 = 0$$


Easily understood ~ D2/anti-D2 pair:

$$\longrightarrow \int_{S_{small}^6} \star G_4 = q_{D2}$$



2-charge Supertubes: Supergravity Description

No net D2 charge, but D2 dipole (and higher) moments:




$$G_4 = d \left[\frac{J r^2}{(r^2 + \rho^2)^4} dt \wedge dz \wedge d\psi \right] + \dots$$

$$\longrightarrow \int_{S_\infty^6} \star G_4 = 0$$

Easily understood ~ D2/anti-D2 pair:

$$\longrightarrow \int_{S_{small}^6} \star G_4 = q_{D2}$$




Bound recovered: Absence of CCCs

$$\longrightarrow J^2 \leq R^2 Q_{F1} Q_{D0}$$

2-charge Supertubes: Supergravity Description

No net D2 charge, but D2 dipole (and higher) moments:




$$G_4 = d \left[\frac{J r^2}{(r^2 + \rho^2)^4} dt \wedge dz \wedge d\psi \right] + \dots$$

$$\longrightarrow \int_{S_\infty^6} \star G_4 = 0$$

Easily understood ~ D2/anti-D2 pair:

$$\longrightarrow \int_{S_{small}^6} \star G_4 = q_{D2}$$




Bound recovered: Absence of CCCs

$$\longrightarrow J^2 \leq R^2 Q_{F1} Q_{D0}$$

Linearity of eqs $\longrightarrow$ Multiple-tube solutions

2-charge Supertubes $\rightarrow$ D1/D5 System


Lunin&Mathur
Maoz&Maldacena
Lunin,Maldacena&Maoz
.....



F1:	-	1
D0:	-	-
d2:	ψ	1


2-charge Supertubes $\rightarrow$ D1/D5 System

Lunin&Mathur
Maoz&Maldacena
Lunin,Maldacena&Maoz
.....




2-charge Supertubes $\rightarrow$ D1/D5 System

Lunin&Mathur
Maoz&Maldacena
Lunin, Maldacena&Maoz
.....



F1: - 1
D0: - -
d2: ψ 1

T1
 $\rightarrow$




P: - 1
D1: - 1
d1: ψ -

T2345
 $\rightarrow$

P: - 1 - - - -
D5: - 1 2 3 4 5
d5: ψ - 2 3 4 5


2-charge Supertubes $\rightarrow$ D1/D5 System

Lunin&Mathur
Maoz&Maldacena
Lunin,Maldacena&Maoz
.....



F1: - 1
D0: - -
d2: ψ 1

T1
→



P: - 1
D1: - 1
d1: ψ -

T2345
→


P: - 1 - - -
D5: - 1 2 3 4 5
d5: ψ - 2 3 4 5

S
→

P: - 1 - - -
NS5: - 1 2 3 4 5
ns5: ψ - 2 3 4 5


2-charge Supertubes $\rightarrow$ D1/D5 System

Lunin&Mathur
Maoz&Maldacena
Lunin,Maldacena&Maoz
.....




2-charge Supertubes $\rightarrow$ D1/D5 System

Lunin&Mathur
Maoz&Maldacena
Lunin,Maldacena&Maoz
.....



F1: - 1
D0: - -
d2: ψ 1

T1
 $\rightarrow$



P: - 1
D1: - 1
d1: ψ -

T2345
 $\rightarrow$

P: - 1 - - -
D5: - 1 2 3 4 5
d5: ψ - 2 3 4 5

S
 $\rightarrow$

P: - 1 - - -
NS5: - 1 2 3 4 5
ns5: ψ - 2 3 4 5

T1
 $\rightarrow$

F1: - 1 - - -
NS5: - 1 2 3 4 5
kk5: ψ • 2 3 4 5

(Trivial T) **S**
 $\rightarrow$

D1: - 1 - - -
D5: - 1 2 3 4 5
kk5: ψ • 2 3 4 5

3-charge Supertubes and Supersymmetric Black Rings: Supergravity Description

Elvang, Emparan, Reall & D.M.
Bena & Warner
Gauntlett & Gutowski

3-charge Supertubes and Supersymmetric Black Rings: Supergravity Description

Elvang, Emparan, Reall & D.M.
Bena & Warner
Gauntlett & Gutowski

First, lift 2-charge supertube to M-theory:

F1:	–	1
D0:	–	–
d2:	ψ	1

3-charge Supertubes and Supersymmetric Black Rings: Supergravity Description

Elvang, Emparan, Reall & D.M.
Bena & Warner
Gauntlett & Gutowski

First, lift 2-charge supertube to M-theory:

$$\begin{array}{lcl}
 \text{F1:} & - & 1 \\
 \text{D0:} & - & - \\
 \text{d2:} & \psi & 1
 \end{array}
 \xrightarrow{\text{T34}}
 \begin{array}{lcl}
 \text{F1:} & - & 1 & - & - & - \\
 \text{D2:} & - & - & - & 3 & 4 \\
 \text{d4:} & \psi & 1 & - & 3 & 4
 \end{array}$$

3-charge Supertubes and Supersymmetric Black Rings: Supergravity Description

Elvang, Emparan, Reall & D.M.
Bena & Warner
Gauntlett & Gutowski

First, lift 2-charge supertube to M-theory:

$$\begin{array}{lcl}
 \begin{array}{l} \text{F1:} \quad - \quad 1 \\ \text{D0:} \quad - \quad - \\ \text{d2:} \quad \psi \quad 1 \end{array} & \xrightarrow{\text{T34}} & \begin{array}{l} \text{F1:} \quad - \quad 1 \quad - \quad - \quad - \\ \text{D2:} \quad - \quad - \quad - \quad 3 \quad 4 \\ \text{d4:} \quad \psi \quad 1 \quad - \quad 3 \quad 4 \end{array} \\
 & & \xrightarrow{\text{Lift}} \\
 & & \begin{array}{l} \text{M2:} \quad - \quad 1 \quad 2 \quad - \quad - \\ \text{M2:} \quad - \quad - \quad - \quad 3 \quad 4 \\ \text{m5:} \quad \psi \quad 1 \quad 2 \quad 3 \quad 4 \end{array}
 \end{array}$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-	-

With 3 charges, each pair expands:


Q_1 M2:	1	2	-	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\}$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\underbrace{\quad\quad\quad}_{\mathbb{T}^6 = \{z^1, \dots, z^6\}} \quad \underbrace{\quad\quad\quad}_{\mathbb{E}^4 = \{r, \Theta, \psi, \phi\}}$$




With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$

$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$




With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$

$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$


$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$



With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\underbrace{\mathbb{T}^6 = \{z^1, \dots, z^6\}}_{\text{charges}} \quad \underbrace{\mathbb{E}^4 = \{r, \Theta, \psi, \phi\}}_{\text{coordinates}}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$


$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}$$


$$H_2 = 1 + \frac{Q_2 - q_1 q_3}{\Sigma} + q_1 q_3 \frac{r^2}{\Sigma^2}$$

$$H_3 = 1 + \frac{Q_3 - q_1 q_2}{\Sigma} + q_1 q_2 \frac{r^2}{\Sigma^2}$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}$$

$$\Sigma(r, \Theta) = \sqrt{(r^2 - R^2)^2 + 4R^2 r^2 \cos^2 \Theta}$$


$$H_2 = 1 + \frac{Q_2 - q_1 q_3}{\Sigma} + q_1 q_3 \frac{r^2}{\Sigma^2}$$

$$H_3 = 1 + \frac{Q_3 - q_1 q_2}{\Sigma} + q_1 q_2 \frac{r^2}{\Sigma^2}$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}$$

$$H_2 = 1 + \frac{Q_2 - q_1 q_3}{\Sigma} + q_1 q_3 \frac{r^2}{\Sigma^2}$$

$$H_3 = 1 + \frac{Q_3 - q_1 q_2}{\Sigma} + q_1 q_2 \frac{r^2}{\Sigma^2}$$


$$\Sigma(r, \Theta) = \sqrt{(r^2 - R^2)^2 + 4R^2 r^2 \cos^2 \Theta}$$

Harmonic with ring source

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}$$

$$\Sigma(r, \Theta) = \sqrt{(r^2 - R^2)^2 + 4R^2 r^2 \cos^2 \Theta}$$

$$H_2 = 1 + \frac{Q_2 - q_1 q_3}{\Sigma} + q_1 q_3 \frac{r^2}{\Sigma^2}$$


$$\omega(r, \Theta; Q_i, q_i, R) = \omega_\psi d\psi + \underbrace{\omega_\phi d\phi}_{\text{Extra angular momentum}}$$

$$H_3 = 1 + \frac{Q_3 - q_1 q_2}{\Sigma} + q_1 q_2 \frac{r^2}{\Sigma^2}$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\underbrace{\quad\quad\quad}_{\mathbb{T}^6 = \{z^1, \dots, z^6\}} \quad \underbrace{\quad\quad\quad}_{\mathbb{E}^4 = \{r, \Theta, \psi, \phi\}}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}$$

$$\Sigma(r, \Theta) = \sqrt{(r^2 - R^2)^2 + 4R^2 r^2 \cos^2 \Theta}$$

$$H_2 = 1 + \frac{Q_2 - q_1 q_3}{\Sigma} + q_1 q_3 \frac{r^2}{\Sigma^2}$$

$$\omega(r, \Theta; Q_i, q_i, R) = \omega_\psi d\psi + \underbrace{\omega_\phi d\phi}_{\text{Extra angular momentum}}$$

$$H_3 = 1 + \frac{Q_3 - q_1 q_2}{\Sigma} + q_1 q_2 \frac{r^2}{\Sigma^2}$$


Globally well-defined (no Dirac-Misner strings) since

$$\omega_\psi(\Theta = 0) = \omega_\phi(\Theta = \pi/2) = 0$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$


$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}, \quad \mathcal{A} = A^1 dz_1 \wedge dz_2 + A^2 dz_3 \wedge dz_4 + A^3 dz_5 \wedge dz_6$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\underbrace{\mathbb{T}^6 = \{z^1, \dots, z^6\}}_{\text{charges}} \quad \underbrace{\mathbb{E}^4 = \{r, \Theta, \psi, \phi\}}_{\text{coordinates}}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$


$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}, \quad \mathcal{A} = A^1 dz_1 \wedge dz_2 + A^2 dz_3 \wedge dz_4 + A^3 dz_5 \wedge dz_6$$

$$A^i = H_i^{-1} (dt + \omega) + \frac{q_i}{2\Sigma} [(r^2 + \Sigma^2 - R^2) d\psi + (r^2 + \Sigma^2 + R^2) d\phi]$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}, \quad \mathcal{A} = A^1 dz_1 \wedge dz_2 + A^2 dz_3 \wedge dz_4 + A^3 dz_5 \wedge dz_6$$

$$A^i = H_i^{-1} (dt + \omega) + \frac{q_i}{2\Sigma} [(r^2 + \Sigma^2 - R^2) d\psi + (r^2 + \Sigma^2 + R^2) d\phi]$$


→ Q_i = M2 charges

q_i = M5 dipoles

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}, \quad \mathcal{A} = A^1 dz_1 \wedge dz_2 + A^2 dz_3 \wedge dz_4 + A^3 dz_5 \wedge dz_6$$

$$A^i = H_i^{-1} (dt + \omega) + \frac{q_i}{2\Sigma} [(r^2 + \Sigma^2 - R^2) d\psi + (r^2 + \Sigma^2 + R^2) d\phi]$$


$$\longrightarrow Q_i = \text{M2 charges} \quad Q_1 \equiv Q_1 - q_2 q_3, \quad Q_2 \equiv Q_2 - q_1 q_3, \quad Q_3 \equiv Q_3 - q_1 q_2$$

$$q_i = \text{M5 dipoles}$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}, \quad \mathcal{A} = A^1 dz_1 \wedge dz_2 + A^2 dz_3 \wedge dz_4 + A^3 dz_5 \wedge dz_6$$

$$A^i = H_i^{-1} (dt + \omega) + \frac{q_i}{2\Sigma} [(r^2 + \Sigma^2 - R^2) d\psi + (r^2 + \Sigma^2 + R^2) d\phi]$$

→ Q_i = M2 charges

$$\mathcal{Q}_1 \equiv Q_1 - q_2 q_3, \quad \mathcal{Q}_2 \equiv Q_2 - q_1 q_3, \quad \mathcal{Q}_3 \equiv Q_3 - q_1 q_2$$


q_i = M5 dipoles

$$H_i \geq 0 \text{ and no CCCs} \rightarrow \mathcal{Q}_i \geq 0$$

With 3 charges, each pair expands:

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-

$$\mathbb{T}^6 = \{z^1, \dots, z^6\} \quad \mathbb{E}^4 = \{r, \Theta, \psi, \phi\}$$



$$ds^2 = ds_5^2 + X^1 (dz_1^2 + dz_2^2) + X^2 (dz_3^2 + dz_4^2) + X^3 (dz_5^2 + dz_6^2)$$

$$X^i = H_i^{-1} (H_1 H_2 H_3)^{1/3}, \quad X^1 X^2 X^3 = 1$$

$$ds_5^2 = -(H_1 H_2 H_3)^{-2/3} (dt + \omega)^2 + (H_1 H_2 H_3)^{1/3} dx_4^2$$

$$H_1 = 1 + \frac{Q_1 - q_2 q_3}{\Sigma} + q_2 q_3 \frac{r^2}{\Sigma^2}, \quad \mathcal{A} = A^1 dz_1 \wedge dz_2 + A^2 dz_3 \wedge dz_4 + A^3 dz_5 \wedge dz_6$$

$$A^i = H_i^{-1} (dt + \omega) + \frac{q_i}{2\Sigma} [(r^2 + \Sigma^2 - R^2) d\psi + (r^2 + \Sigma^2 + R^2) d\phi]$$


→ Q_i = M2 charges

$$Q_1 \equiv Q_1 - q_2 q_3, \quad Q_2 \equiv Q_2 - q_1 q_3, \quad Q_3 \equiv Q_3 - q_1 q_2$$

q_i = M5 dipoles


$$H_i \geq 0 \text{ and no CCCs} \rightarrow Q_i \geq 0 \quad q \equiv (q_1 q_2 q_3)^{1/3}$$

Physical Properties



Physical Properties


Solution preserves 4 supercharges (1/8-Susy in 11D)



Physical Properties

Solution preserves 4 supercharges (1/8-Susy in 11D)

Stationary and asymptotically flat (in 5D)




Physical Properties

Solution preserves 4 supercharges (1/8-Susy in 11D)

Stationary and asymptotically flat (in 5D)

Independent Conserved Charges: Three Gauge Charges Q_i and two ang mom:



Physical Properties


Solution preserves 4 supercharges (1/8-Susy in 11D)

Stationary and asymptotically flat (in 5D)

Independent Conserved Charges: Three Gauge Charges Q_i and two ang mom:

$$J_\phi = \frac{\pi}{8G_5} (q_1 Q_1 + q_2 Q_2 + q_3 Q_3 - q_1 q_2 q_3)$$

$$J_\psi = \frac{\pi}{8G_5} 2R^2 (q_1 + q_2 + q_3) + J_\phi$$



Physical Properties

Solution preserves 4 supercharges (1/8-Susy in 11D)


Stationary and asymptotically flat (in 5D)

Independent Conserved Charges: Three Gauge Charges Q_i and two ang mom:

$$J_\phi = \frac{\pi}{8G_5} (q_1 Q_1 + q_2 Q_2 + q_3 Q_3 - q_1 q_2 q_3)$$

$$J_\psi = \frac{\pi}{8G_5} 2R^2 (q_1 + q_2 + q_3) + J_\phi$$

$$\longrightarrow J_\psi > J_\phi$$



Physical Properties

Solution preserves 4 supercharges (1/8-Susy in 11D)

Stationary and asymptotically flat (in 5D)


Independent Conserved Charges: Three Gauge Charges Q_i and two ang mom:

$$J_\phi = \frac{\pi}{8G_5} (q_1 Q_1 + q_2 Q_2 + q_3 Q_3 - q_1 q_2 q_3)$$

$$J_\psi = \frac{\pi}{8G_5} 2R^2 (q_1 + q_2 + q_3) + J_\phi$$

$$\longrightarrow J_\psi > J_\phi$$

(Mass is fixed by BPS bound: $M = Q_1 + Q_2 + Q_3$)



Physical Properties

Solution preserves 4 supercharges (1/8-Susy in 11D)

Stationary and asymptotically flat (in 5D)

Independent Conserved Charges: Three Gauge Charges Q_i and two ang mom:


$$J_\phi = \frac{\pi}{8G_5} (q_1 Q_1 + q_2 Q_2 + q_3 Q_3 - q_1 q_2 q_3)$$

$$J_\psi = \frac{\pi}{8G_5} 2R^2 (q_1 + q_2 + q_3) + J_\phi$$

$$\longrightarrow J_\psi > J_\phi$$

(Mass is fixed by BPS bound: $M = Q_1 + Q_2 + Q_3$)


Absence of CCCs $\longleftrightarrow L^2 = \frac{1}{4q^4} \left(2 \sum_{i < j} Q_i q_i Q_j q_j - \sum_i Q_i^2 q_i^2 - 4R^2 q^3 \sum_i q_i \right) \geq 0$



Physical Properties

Solution preserves 4 supercharges (1/8-Susy in 11D)

Stationary and asymptotically flat (in 5D)



Independent Conserved Charges: Three Gauge Charges Q_i and two ang mom:

$$J_\phi = \frac{\pi}{8G_5} (q_1 Q_1 + q_2 Q_2 + q_3 Q_3 - q_1 q_2 q_3)$$

$$J_\psi = \frac{\pi}{8G_5} 2R^2 (q_1 + q_2 + q_3) + J_\phi$$

$$\longrightarrow J_\psi > J_\phi$$

(Mass is fixed by BPS bound: $M = Q_1 + Q_2 + Q_3$)


Absence of CCCs $\longleftrightarrow L^2 = \frac{1}{4q^4} \left(2 \sum_{i < j} Q_i q_i Q_j q_j - \sum_i Q_i^2 q_i^2 - 4R^2 q^3 \sum_i q_i \right) \geq 0$

$\longrightarrow$ Upper bound on R, or equivalently on $J_\psi - J_\phi$

Physical Properties

Solution preserves 4 supercharges (1/8-Susy in 11D)

Stationary and asymptotically flat (in 5D)



Independent Conserved Charges: Three Gauge Charges Q_i and two ang mom:

$$J_\phi = \frac{\pi}{8G_5} (q_1 Q_1 + q_2 Q_2 + q_3 Q_3 - q_1 q_2 q_3)$$

$$J_\psi = \frac{\pi}{8G_5} 2R^2 (q_1 + q_2 + q_3) + J_\phi$$

$$\longrightarrow J_\psi > J_\phi$$

(Mass is fixed by BPS bound: $M = Q_1 + Q_2 + Q_3$)

Absence of CCCs $\longleftrightarrow L^2 = \frac{1}{4q^4} \left(2 \sum_{i < j} Q_i q_i Q_j q_j - \sum_i Q_i^2 q_i^2 - 4R^2 q^3 \sum_i q_i \right) \geq 0$


$\longrightarrow$ Upper bound on R , or equivalently on $J_\psi - J_\phi$

$L^2 > 0 \longrightarrow$ Regular event horizon at $r = R, \Theta = \pi/2$ of topology $S^1 \times S^2$:

Physical Properties

Solution preserves 4 supercharges (1/8-Susy in 11D)

Stationary and asymptotically flat (in 5D)



Independent Conserved Charges: Three Gauge Charges Q_i and two ang mom:

$$J_\phi = \frac{\pi}{8G_5} (q_1 Q_1 + q_2 Q_2 + q_3 Q_3 - q_1 q_2 q_3)$$

$$J_\psi = \frac{\pi}{8G_5} 2R^2 (q_1 + q_2 + q_3) + J_\phi$$

$$\longrightarrow J_\psi > J_\phi$$

(Mass is fixed by BPS bound: $M = Q_1 + Q_2 + Q_3$)

Absence of CCCs $\longleftrightarrow L^2 = \frac{1}{4q^4} \left(2 \sum_{i < j} Q_i q_i Q_j q_j - \sum_i Q_i^2 q_i^2 - 4R^2 q^3 \sum_i q_i \right) \geq 0$

$\longrightarrow$ Upper bound on R, or equivalently on $J_\psi - J_\phi$

$L^2 > 0 \longrightarrow$ Regular event horizon at $r = R, \Theta = \pi/2$ of topology $S^1 \times S^2$:

$$ds_H^2 = L^2 d\psi'^2 + \frac{q^2}{2} d\Omega_2^2 \longrightarrow L \text{ sets physical length, } A_H = 2\pi^2 L q^2$$

Example:

$$Q_i = Q, q_i = q$$

Example:



$$Q_i = Q, q_i = q$$

$$J_\phi \propto q(3Q - q^2)$$

$$J_\psi \propto J_\phi + 6qR^2$$

Example: $\longrightarrow$
 $Q_i = Q, q_i = q$

$$J_\phi \propto q(3Q - q^2)$$

$$J_\psi \propto J_\phi + 6qR^2$$

$$A_H = \pi^2 q \sqrt{3 \left[(Q - q^2)^2 - 4q^2 R^2 \right]}$$

$$L^2 = 3 \left(\frac{(Q - q^2)^2}{4q^2} - R^2 \right)$$

$$R < \frac{Q - q^2}{2q}$$

Example:
 $Q_i = Q, q_i = q$


$$J_\phi \propto q(3Q - q^2)$$

$$J_\psi \propto J_\phi + 6qR^2$$

$$A_H = \pi^2 q \sqrt{3 \left[(Q - q^2)^2 - 4q^2 R^2 \right]}$$

$$L^2 = 3 \left(\frac{(Q - q^2)^2}{4q^2} - R^2 \right)$$

$$R < \frac{Q - q^2}{2q}$$



Example: $\longrightarrow$
 $Q_i = Q, q_i = q$


$$J_\phi \propto q(3Q - q^2)$$

$$J_\psi \propto J_\phi + 6qR^2$$

$$A_H = \pi^2 q \sqrt{3 \left[(Q - q^2)^2 - 4q^2 R^2 \right]}$$

$$L^2 = 3 \left(\frac{(Q - q^2)^2}{4q^2} - R^2 \right)$$

$$R < \frac{Q - q^2}{2q}$$



Example:
 $Q_i = Q, q_i = q$


$$J_\phi \propto q(3Q - q^2)$$

$$J_\psi \propto J_\phi + 6qR^2$$

$$A_H = \pi^2 q \sqrt{3 \left[(Q - q^2)^2 - 4q^2 R^2 \right]}$$

$$L^2 = 3 \left(\frac{(Q - q^2)^2}{4q^2} - R^2 \right)$$

$$R < \frac{Q - q^2}{2q}$$



Example: $\longrightarrow$
 $Q_i = Q, q_i = q$


$$J_\phi \propto q(3Q - q^2)$$

$$J_\psi \propto J_\phi + 6qR^2$$

$$A_H = \pi^2 q \sqrt{3 \left[(Q - q^2)^2 - 4q^2 R^2 \right]}$$

$$L^2 = 3 \left(\frac{(Q - q^2)^2}{4q^2} - R^2 \right)$$

$$R < \frac{Q - q^2}{2q}$$



Example: $\longrightarrow$
 $Q_i = Q, q_i = q$


$$J_\phi \propto q(3Q - q^2)$$

$$J_\psi \propto J_\phi + 6qR^2$$

$$A_H = \pi^2 q \sqrt{3 \left[(Q - q^2)^2 - 4q^2 R^2 \right]}$$

$$L^2 = 3 \left(\frac{(Q - q^2)^2}{4q^2} - R^2 \right)$$

$$R < \frac{Q - q^2}{2q}$$



Non-uniqueness

7 parameters: Q_i , q_i , R

Non-uniqueness


7 parameters: Q_i, q_i, R

5 independent charges: Q_i, J_ψ, J_ϕ

Non-uniqueness

7 parameters: Q_i, q_i, R

5 independent charges: Q_i, J_ψ, J_ϕ




Classical infinite non-uniqueness
parametrised by two continuous variables

Non-uniqueness

7 parameters: Q_i, q_i, R

5 independent charges: Q_i, J_ψ, J_ϕ




Classical infinite non-uniqueness
parametrised by two continuous variables

Contour-plot area as function of q_1 and q_2 keeping $Q_i = Q$, J_ψ and J_ϕ fixed :

Non-uniqueness


7 parameters: Q_i, q_i, R

5 independent charges: Q_i, J_ψ, J_ϕ



Classical infinite non-uniqueness
parametrised by two continuous variables


Contour-plot area as function of q_1 and q_2 keeping $Q_i = Q$, J_ψ and J_ϕ fixed :



Non-uniqueness


7 parameters: Q_i, q_i, R

5 independent charges: Q_i, J_ψ, J_ϕ



Classical infinite non-uniqueness
parametrised by two continuous variables

Contour-plot area as function of q_1 and q_2 keeping $Q_i = Q$, J_ψ and J_ϕ fixed :



Even with extra requirement
of constant area




Infinitely-many
black rings

Non-uniqueness


7 parameters: Q_i, q_i, R

5 independent charges: Q_i, J_ψ, J_ϕ




Classical infinite non-uniqueness
parametrised by two continuous variables

Contour-plot area as function of q_1 and q_2 keeping $Q_i = Q$, J_ψ and J_ϕ fixed :



Even with extra requirement
of constant area




Infinitely-many
black rings

(except for maximum area)

Non-uniqueness


7 parameters: Q_i, q_i, R

5 independent charges: Q_i, J_ψ, J_ϕ



Classical infinite non-uniqueness
parametrised by two continuous variables

Contour-plot area as function of q_1 and q_2 keeping $Q_i = Q$, J_ψ and J_ϕ fixed :



Even with extra requirement
of constant area




Infinitely-many
black rings

(except for maximum area)

Non-uniqueness


7 parameters: Q_i, q_i, R

5 independent charges: Q_i, J_ψ, J_ϕ




Classical infinite non-uniqueness
parametrised by two continuous variables

Contour-plot area as function of q_1 and q_2 keeping $Q_i = Q$, J_ψ and J_ϕ fixed :



Even with extra requirement
of constant area



Infinitely-many
black rings

(except for maximum area)

Particular cases and limits: $R=0$

Example: $\longrightarrow$
 $Q_i = Q, q_i = q$


$$J_\phi \propto q(3Q - q^2)$$

$$J_\psi \propto J_\phi + 6qR^2$$

$$A_H = \pi^2 q \sqrt{3 \left[(Q - q^2)^2 - 4q^2 R^2 \right]}$$


$$L^2 = 3 \left(\frac{(Q - q^2)^2}{4q^2} - R^2 \right)$$

$$R < \frac{Q - q^2}{2q}$$




2-charge Supertubes $\rightarrow$ D1/D5 System

Lunin&Mathur
Maoz&Maldacena
Lunin,Maldacena&Maoz
.....



F1: - 1
D0: - -
d2: ψ 1

T1
 $\rightarrow$



P: - 1
D1: - 1
d1: ψ -

T2345
 $\rightarrow$

P: - 1 - - -
D5: - 1 2 3 4 5
d5: ψ - 2 3 4 5

S
 $\rightarrow$

P: - 1 - - -
NS5: - 1 2 3 4 5
ns5: ψ - 2 3 4 5

T1
 $\rightarrow$

F1: - 1 - - -
NS5: - 1 2 3 4 5
kk5: ψ • 2 3 4 5

(Trivial T) **S**
 $\rightarrow$

D1: - 1 - - -
D5: - 1 2 3 4 5
kk5: ψ • 2 3 4 5

Particular cases and limits: $R=0$

Particular cases and limits: $R=0$

$R=0 \rightarrow$ BMPV (Breckenridge, Myers, Peet & Vafa) $J_\psi = J_\phi = J$ Horizon = S^3

Particular cases and limits: $R=0$

$R=0 \rightarrow$ BMPV (Breckenridge, Myers, Peet & Vafa) $J_\psi = J_\phi = J$ Horizon = S^3

Change in topology $\rightarrow$ Discontinuous change in entropy:

Particular cases and limits: $R=0$

$R=0 \rightarrow$ BMPV (Breckenridge, Myers, Peet & Vafa) $J_\psi = J_\phi = J$ Horizon = S^3

Change in topology $\rightarrow$ Discontinuous change in entropy:

$$A_{BMPV} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$

Particular cases and limits: $R=0$

$R=0 \rightarrow$ BMPV (Breckenridge, Myers, Peet & Vafa) $J_\psi = J_\phi = J$ Horizon = S^3

Change in topology $\rightarrow$ Discontinuous change in entropy:

$$A_{BMPV} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$

$$\lim_{R \rightarrow 0} A_{ring} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$

Particular cases and limits: $R=0$

$R=0 \rightarrow$ BMPV (Breckenridge, Myers, Peet & Vafa) $J_\psi = J_\phi = J$ Horizon = S^3

Change in topology $\rightarrow$ Discontinuous change in entropy:

$$A_{BMPV} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2} \quad \longrightarrow \quad \lim_{R \rightarrow 0} A_{ring} \leq A_{BMPV}$$
$$\lim_{R \rightarrow 0} A_{ring} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$

Particular cases and limits: $R=0$

$R=0 \rightarrow$ BMPV (Breckenridge, Myers, Peet & Vafa) $J_\psi = J_\phi = J$ Horizon = S^3


Change in topology $\rightarrow$ Discontinuous change in entropy:

$$A_{BMPV} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$

$$\lim_{R \rightarrow 0} A_{ring} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$

$$\longrightarrow \lim_{R \rightarrow 0} A_{ring} \leq A_{BMPV}$$

Eg. $Q_i = Q$, $q_i = q$:



Particular cases and limits: $R=0$

$R=0 \rightarrow$ BMPV (Breckenridge, Myers, Peet & Vafa) $J_\psi = J_\phi = J$ Horizon = S^3

Change in topology $\rightarrow$ Discontinuous change in entropy:


$$A_{BMPV} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$


$$\lim_{R \rightarrow 0} A_{ring} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$

$$\longrightarrow \lim_{R \rightarrow 0} A_{ring} \leq A_{BMPV}$$

Eg. $Q_i = Q$, $q_i = q$:

Analogous to the merging
of two 4D RN black holes:


 $A = 2 \times 4\pi Q^2$



Particular cases and limits: $R=0$

$R=0 \rightarrow$ BMPV (Breckenridge, Myers, Peet & Vafa) $J_\psi = J_\phi = J$ Horizon = S^3

Change in topology $\rightarrow$ Discontinuous change in entropy:

$$A_{BMPV} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$

$$\lim_{R \rightarrow 0} A_{ring} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$


$$\rightarrow \lim_{R \rightarrow 0} A_{ring} \leq A_{BMPV}$$

Eg. $Q_i = Q$, $q_i = q$:

Analogous to the merging
of two 4D RN black holes:


 $A = 2 \times 4\pi Q^2$
 $R > 0$


 $A = 4\pi (2Q)^2$
 $R = 0$



Particular cases and limits: $R \rightarrow \infty$

Particular cases and limits: $R=0$

$R=0 \rightarrow$ BMPV (Breckenridge, Myers, Peet & Vafa) $J_\psi = J_\phi = J$ Horizon = S^3

Change in topology $\rightarrow$ Discontinuous change in entropy:

$$A_{BMPV} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$

$$\lim_{R \rightarrow 0} A_{ring} = 2\pi^2 \sqrt{Q_1 Q_2 Q_3 - Q_1 Q_2 Q_3 - \left(\frac{4G_5 J}{\pi}\right)^2}$$


$$\longrightarrow \lim_{R \rightarrow 0} A_{ring} \leq A_{BMPV}$$

Eg. $Q_i = Q$, $q_i = q$:

Analogous to the merging
of two 4D RN black holes:


 $A = 2 \times 4\pi Q^2$
 $R > 0$


 $A = 4\pi (2Q)^2$
 $R = 0$




Particular cases and limits: $R \rightarrow \infty$

Particular cases and limits: $R \rightarrow \infty$

Send $R \rightarrow \infty$ keeping Q_i / R and q_i fixed

Particular cases and limits: $R \rightarrow \infty$


Send $R \rightarrow \infty$ keeping Q_i / R and q_i fixed $\longrightarrow$ 'Straight line' solution of Bena



Particular cases and limits: $R \rightarrow \infty$

Send $R \rightarrow \infty$ keeping Q_i / R and q_i fixed $\longrightarrow$ 'Straight line' solution of Bena

Important: $J_\psi = P_\psi \neq 0$ but $J_\phi \rightarrow 0$!




Particular cases and limits: $R \rightarrow \infty$

Send $R \rightarrow \infty$ keeping Q_i / R and q_i fixed $\longrightarrow$ 'Straight line' solution of Bena

Important: $J_\psi = P_\psi \neq 0$ but $J_\phi \rightarrow 0$!

In fact, required by $SO(3)$ invariance:




Particular cases and limits: $R \rightarrow \infty$

Send $R \rightarrow \infty$ keeping Q_i / R and q_i fixed $\longrightarrow$ 'Straight line' solution of Bena

Important: $J_\psi = P_\psi \neq 0$ but $J_\phi \rightarrow 0$!

In fact, required by $SO(3)$ invariance:



Speculation: J_ϕ Poynting-generated by SUGRA fields

Worldvolume 3-charge Supertubes = Calibrated Supertubes

Worldvolume 3-charge Supertubes = Calibrated Supertubes

First step: F1/D4/D0 bound state with D2/D6/NS5 dipoles

Bena & Kraus

Worldvolume 3-charge Supertubes = Calibrated Supertubes

First step: F1/D4/D0 bound state with D2/D6/NS5 dipoles

Bena & Kraus

Q_1 M2:	1	2	-	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-	-
q_1 m5:	-	-	3	4	5	6	ψ	-	-	-
q_2 m5:	1	2	-	-	5	6	ψ	-	-	-
q_3 m5:	1	2	3	4	-	-	ψ	-	-	-

Worldvolume 3-charge Supertubes = Calibrated Supertubes

First step: F1/D4/D0 bound state with D2/D6/NS5 dipoles

Bena & Kraus


q_1 m5:	-	-	3	4	5	6	7	-	-	-
q_2 m5:	1	2	-	-	5	6	7	-	-	-
q_3 m5:	1	2	3	4	-	-	7	-	-	-

Worldvolume 3-charge Supertubes = Calibrated Supertubes

First step: F1/D4/D0 bound state with D2/D6/NS5 dipoles

Bena & Kraus

q_1 m5:	-	-	3	4	5	6	7	-	-	-
q_2 m5:	1	2	-	-	5	6	7	-	-	-
q_3 m5:	1	2	3	4	-	-	7	-	-	-



Single M5-brane = 2-dim complex surface S in $\mathbb{C}^3$

Gibbons & Papadopoulos
Gauntlett, Lambert & West

Calibrated by $h \wedge h$, with $h = dz^1 \wedge dz^2 + dz^3 \wedge dz^4 + dz^5 \wedge dz^6$ Kahler form

Worldvolume 3-charge Supertubes = Calibrated Supertubes

First step: F1/D4/D0 bound state with D2/D6/NS5 dipoles

Bena & Kraus

Q_1 M2:	1	2	-	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-	-
q_1 m5:	-	-	3	4	5	6	C	-	-	-
q_2 m5:	1	2	-	-	5	6	C	-	-	-
q_3 m5:	1	2	3	4	-	-	C	-	-	-



Single M5-brane = 2-dim complex surface S in $\mathbb{C}^3$

Gibbons & Papadopoulos
Gauntlett, Lambert & West

Calibrated by $h \wedge h$, with $h = dz^1 \wedge dz^2 + dz^3 \wedge dz^4 + dz^5 \wedge dz^6$ Kahler form

Turning on $H = h \wedge d\sigma + H'$ induces M2 charges and allows arbitrary C

Elvang, Emparan, Reall & D.M.

Worldvolume 3-charge Supertubes = Calibrated Supertubes

First step: F1/D4/D0 bound state with D2/D6/NS5 dipoles

Bena & Kraus

Q_1 M2:	1	2	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-
q_1 m5:	-	-	3	4	5	6	C	-	-
q_2 m5:	1	2	-	-	5	6	C	-	-
q_3 m5:	1	2	3	4	-	-	C	-	-



Single M5-brane = 2-dim complex surface S in $\mathbb{C}^3$

Gibbons & Papadopoulos
Gauntlett, Lambert & West

Calibrated by $h \wedge h$, with $h = dz^1 \wedge dz^2 + dz^3 \wedge dz^4 + dz^5 \wedge dz^6$ Kahler form

Turning on $H = h \wedge d\sigma + H'$ induces M2 charges and allows arbitrary C

Elvang, Emparan, Reall & D.M.


Captures 3 dipoles but not $J_\phi \rightarrow$ Defect or Poynting-generated?

Worldvolume 3-charge Supertubes = Calibrated Supertubes

First step: F1/D4/D0 bound state with D2/D6/NS5 dipoles

Bena & Kraus

Q_1 M2:	1	2	-	-	-	-	-	-	-	-
Q_2 M2:	-	-	3	4	-	-	-	-	-	-
Q_3 M2:	-	-	-	-	5	6	-	-	-	-
q_1 m5:	-	-	3	4	5	6	C	-	-	-
q_2 m5:	1	2	-	-	5	6	C	-	-	-
q_3 m5:	1	2	3	4	-	-	C	-	-	-



Single M5-brane = 2-dim complex surface S in $\mathbb{C}^3$

Gibbons & Papadopoulos
Gauntlett, Lambert & West

Calibrated by $h \wedge h$, with $h = dz^1 \wedge dz^2 + dz^3 \wedge dz^4 + dz^5 \wedge dz^6$ Kahler form

Turning on $H = h \wedge d\sigma + H'$ induces M2 charges and allows arbitrary C

Elvang, Emparan, Reall & D.M.

Captures 3 dipoles but not $J_\phi \rightarrow$ Defect or Poynting-generated?

(locally 2-charge $\rightarrow$ susy junction)

3-charge Supertubes → D1/D5/P System

3-charge Supertubes $\rightarrow$ D1/D5/P System

Reducing to 10D and dualising:

Q_1 D5:	1	2	3	4	5	-	-	-	-
Q_2 D1:	-	-	-	-	5	-	-	-	-
Q_3 P:	-	-	-	-	5	-	-	-	-
q_1 d1:	-	-	-	-	-	ψ	-	-	-
q_2 d5:	1	2	3	4	-	ψ	-	-	-
q_3 kk5:	1	2	3	4	•	ψ	-	-	-

3-charge Supertubes $\rightarrow$ D1/D5/P System

Elvang, Emparan, Reall & D.M.

Reducing to 10D and dualising:

Decoupling limit: $\alpha' \rightarrow 0$, r/α' fixed, etc.

Q_1 D5:	1	2	3	4	5	-	-	-	-
Q_2 D1:	-	-	-	-	5	-	-	-	-
Q_3 P:	-	-	-	-	5	-	-	-	-
q_1 d1:	-	-	-	-	-	ψ	-	-	-
q_2 d5:	1	2	3	4	-	ψ	-	-	-
q_3 kk5:	1	2	3	4	•	ψ	-	-	-

$$AdS_3 \times S^3 \times T^4 \xrightarrow{\text{RG}} AdS_3 \times S^3/Z_{q_3} \times T^4$$

$$\ell_{UV} = (Q_1 Q_2)^{1/4} > \ell_{IR} = \sqrt{q_3} (q_1 q_2)^{1/4}$$

3-charge Supertubes $\rightarrow$ D1/D5/P System

Elvang, Emparan, Reall & D.M.

Reducing to 10D and dualising:

Decoupling limit: $\alpha' \rightarrow 0$, r/α' fixed, etc.

Q_1 D5:	1	2	3	4	5	-	-	-	-
Q_2 D1:	-	-	-	-	5	-	-	-	-
Q_3 P:	-	-	-	-	5	-	-	-	-
q_1 d1:	-	-	-	-	-	ψ	-	-	-
q_2 d5:	1	2	3	4	-	ψ	-	-	-
q_3 kk5:	1	2	3	4	•	ψ	-	-	-

$$AdS_3 \times S^3 \times T^4 \xrightarrow{\text{RG}} AdS_3 \times S^3/Z_{q_3} \times T^4$$


$$\ell_{UV} = (Q_1 Q_2)^{1/4} > \ell_{IR} = \sqrt{q_3} (q_1 q_2)^{1/4}$$

Reproduction of the entropy by 'splitting string':

Bena & Kraus

$$J_\phi = \frac{\pi}{8G_5} (q_1 Q_1 + q_2 Q_2 + q_3 Q_3 - q_1 q_2 q_3)$$

$$J_\psi = \frac{\pi}{8G_5} 2R^2 (q_1 + q_2 + q_3) + J_\phi$$



Supertube-like, carried by fermion zero modes

BMPV-like, carried by fermion momentum modes

End of slide show, click to exit.