

Title: M-Theory inflation from multi M5-Brane dynamics

Date: Apr 26, 2005 11:00 AM

URL: <http://pirsa.org/05040066>

Abstract:

# M-THEORY INFLATION FROM MULTI M5-BRANE DYNAMICS

Axel Krause

University of Maryland

Perimeter Institute

String Seminar

April 26, 2005

based on

K. & M. Becker, AK: hep-th/0501130, NPB

+ older work w/ M. Becker, G. Curio

+ work in progress

## Overview

### 1. Some Background:

- Heterotic M-Theory Flux Compactification Backgrounds & Newton's Constant
- Moduli Stabilization & de Sitter Vacua

### 2. Inflation:


- Single Field Inflation in String-Theory?
- Power-Law and Assisted Inflation
- Realizing Assisted Inflation in M-Theory
- The Whole Evolution: Cascade Inflation
- Number of E-Foldings

### 3. Conclusion



## The Framework: Heterotic M-Theory


M-Theory on  $S^1/Z_2$



$\Rightarrow$  10d fixed planes become **boundaries** of 11d interior bulk spacetime such that 11d M-theory universe looks like:

Visible Boundary = Us:  
10d  $E_8$  SYM Gauge Theory

Hidden Boundary:  
10d  $E_8$  SYM Gauge Theory



To render the theory **anomaly-free** a supersymmetric Yang Mills theory with gauge group  $E_8$  has to reside on each of the fixed planes.

$L$  is a measure for the **string coupling**  $g_s$  since  $L \sim g_s^{2/3}$   
 $\Rightarrow$  recover weakly coupled heterotic string at  $L \rightarrow 0$



Important: **boundaries** act as **magnetic sources** for the 4-form field-strength  $G = dC$ . Therefore, Bianchi identity acquires a non-trivial right-hand-side

$$dG \sim \left(\frac{\kappa}{4\pi}\right)^{2/3} \sum_{i=1,2} \left( \text{tr} F^{(i)} \wedge F^{(i)} - \frac{1}{2} \text{tr} R \wedge R \right) \delta(x^{11} - L^{(i)}) \wedge dx^{11}$$

### What is the Geometry for M-theory on the Spacetime $\mathcal{M}^4 \times CY_3 \times S^1/Z_2$ ?

$\Rightarrow$  non-vanishing "flux"  $G \neq 0$  implies a **warped product** structure [Witten '96]

$$ds^2 = (1 - f_{lin}(x^n, x^{11})) \eta_{\mu\nu} dx^\mu dx^\nu + (1 + f_{lin}(x^n, x^{11})) (g_{lm}^{CY}(x^n) dx^l dx^m + dx^{11} dx^{11})$$

with (approximate solution)

$$f_{lin} = -\frac{2}{3} x^{11} Q_v, \quad |f_{lin}| \ll 1$$

where  $Q_v$  is the visible boundary charge determining the coupling to the 3-form potential  $C$



# Exact warped geometry for compactification in the presence of $G$ flux

[Curio, AK '00, '03]

$$ds^2 = e^{-f(x^n, x^{11})} \eta_{\mu\nu} dx^\mu dx^\nu + e^{f(x^n, x^{11})} (g_{lm}^{CY}(x^n) dx^l dx^m + dx^{11} dx^{11})$$

with warp-factor


$$e^{f(x^{11})} = |1 - x^{11} Q_v|^{2/3}$$

Let's illustrate this geometry with the volume dependence of the deformed Calabi-Yau:

E. Witten '96 -----

G. Curio, AK '00, '03 ———

Warped CY Volume



⇒ solves problem with negative CY-volume (and negative metric!). Now localized naked singularity. Quantum corrections give  $R^4$  terms which lead to small positive shift in volume if Euler-number of CY is positive  $\chi > 0$  (Strominger '98)



what can we use it for?

⇒ Derivation of 4d Newton's Constant, 4d de Sitter Vacua with Dark Matter (Cosmology), Inflation (Early Universe Cosmology), to which we now turn ...

### Newton's Constant

Exact integration over the internal 7 dimensions gives (with  $R_v = 1/M_{GUT}$ ,  $M_{GUT} = 3 \times 10^{16}$  GeV,  $\alpha_v = 1/25$ )

$$G_N = \frac{G_{N,c}}{(1 - \text{sign}(1 - Q_v L)|1 - Q_v L|^{8/3})} \\ \geq G_{N,c} = (1.2 \times 10^{19} \text{ GeV})^{-2}$$

since  $L \leq L_c = 1/Q_v$

Compare with the measured value

$$G_N^{exp} = (1.22 \times 10^{19} \text{ GeV})^{-2}$$

⇒ Can  $L$  be Stabilized at its critical value  $L_c$ ?





- Now that we know the background geometry, what can we use it for?

⇒ Derivation of 4d Newton's Constant, 4d de Sitter Vacua with Dark Matter (Cosmology), **Inflation** (Early Universe Cosmology), to which we now turn ...

### Newton's Constant

Exact integration over the internal 7 dimensions gives  
(with  $R_v = 1/M_{GUT}$ ,  $M_{GUT} = 3 \times 10^{16} \text{ GeV}$ ,  $\alpha_v = 1/25$ )

$$G_N = \frac{G_{N,c}}{(1 - \text{sign}(1 - Q_v L)|1 - Q_v L|^{8/3})}$$

$$\geq G_{N,c} = (1.2 \times 10^{19} \text{ GeV})^{-2}$$

since  $L \leq L_c = 1/Q_v$

Compare with the measured value

$$G_N^{exp} = (1.22 \times 10^{19} \text{ GeV})^{-2}$$

⇒ Can  $L$  be Stabilized at its critical value  $L_c$ ?




## Moduli Stabilization and De Sitter Vacua

- Cosmological Data compatible with (approximate) de Sitter Universe today. So can we break supersymmetry and stabilize the moduli to obtain the correct  $G_N$  and a realistic (gauge, matter fields,  $M_{SUSY}$ ) de Sitter Vacuum? [Becker, Curio, AK '04]

Let me here focus on the stabilization of  $L$  for which there exist two competing non-perturbative effects

- 1) Open Membrane Instantons = M2 Branes wrapped around the spacelike 3-cycle  $\Sigma_2 \times S^1/\mathbb{Z}_2$

[Berglund et al. '95; Witten '99; Moore, Peradze, Saulina '00; etc.]




repelling non-perturbative force exerted by these open membrane instantons. Can be seen from the resulting superpotential

$$|W_{OM}| \sim e^{-T}, \quad \text{Re}(T) \sim L$$




## 2) Gaugino Condensation on the Hidden Boundary:



gives an attractive force between the two boundaries

$$W_{GC} \sim e^{-\frac{1}{c_H}(S - \gamma_v T)}$$

Superpotentials for both effects:




$|W_{Om}|$  (left curve) and  $|W_{GC}|$  (right curve)

By working out the full effective potential

$$U = M_{Pl}^4 e^K \left( \sum K^{I\bar{J}} D_{\bar{I}} \bar{W} D_J W - 3|W|^2 \right)$$

one confirms a robust minimum through the balancing of both effects (OM and GC)



the resulting minimum potential as a function of  $\mathcal{L} = L/l_{norm}$

Moreover:

Balancing  $H$ -Flux with GC ("Perfect Square")  $\Rightarrow$  Stabilization of Complex Structure Moduli

Balancing M5-instantons with GC  $\Rightarrow$  Stabilization of Volume Modulus

Results:

- > local de Sitter minima
- > vacua lie in a regime where  $\mathcal{V}_{OM}, \mathcal{V} \gg 1$ , i.e. where supergravity is under good control
- >  $D_S W = -\frac{W_{GC}}{C_H} \neq 0 \rightarrow$  supersymmetry is broken spontaneously through F-terms
- > whereas the dilaton  $L \sim g_s^{2/3}$  shows a runaway behaviour at weak coupling it gets naturally stabilized at strong coupling

[Curio, AK'01]



## What Do we Learn about the Hidden Gauge Group?

The values for

$$\tilde{M}_{SUSY} = M_{SUSY}/e^{K(z)}, \quad \tilde{m}_{3/2} = m_{3/2}/e^{K(z)}$$

are given in the following table at the critical point (notice the factor  $e^{K(z)} \simeq 1/V_v \simeq 1/300$  as compared to the actual physical parameters  $M_{SUSY}, m_{3/2}$ )

$H, C_H$	$E_8, 30$	$E_6, 12$	$SO(10), 8$	$SU(5), 5$	$SU(3), 3$
$L_0$	1.1	3.0	4.1	5.5	7.1
$L_c$	7.2	7.2	7.2	7.2	7.2
$M_{SUSY}/TeV$	$3.1 \times 10^{10}$	$1.1 \times 10^8$	$8.5 \times 10^6$	527697	39158
$\tilde{m}_{3/2}/TeV$	$3.4 \times 10^8$	$1.3 \times 10^6$	103752	6991	567

Result: Hidden Gauge Group of small rank favored

- it brings  $M_{SUSY}$  close to the **TeV regime**
- it leads to stabilization of hidden boundary close to  $L_c$ . This leads to the **correct 4d Newton's Constant**

Important: once the hidden  $E_8$  is broken to something smaller like, say a hidden SM sector, one has **hidden matter** which might account for **dark matter** (interacts with visible matter or gauge fields only (super-) gravitationally and can be expected to exhibit similar clustering as visible matter!



## Single-Field Inflation in String-Theory?

To embed single-field slow-roll inflation into string- or M-theory, take your favorite model, use the available fluxes and non-perturbative effects and stabilize all moduli except for one,  $\varphi$ .

Since the aim is to derive a single extremely flat  $\varphi$  direction, the masses of the stabilized moduli should be heavier than the one for  $\varphi$ , such that we can integrate these moduli out and remain with an effective potential  $U(\varphi)$ .

To see whether the resulting  $U(\varphi)$  is flat enough to give rise to a period of inflation which is long enough (50-60 e-foldings usually), you have to check that both

$$\epsilon = \frac{M_{Pl}^2}{2} \left( \frac{U'(\varphi)}{U(\varphi)} \right)^2 \ll 1$$
$$|\eta| = \left| M_{Pl}^2 \frac{U''(\varphi)}{U(\varphi)} \right| \ll 1$$

While the  $\epsilon$  condition is typically easy to satisfy, the contrary is true for the  $\eta$  constraint and hence developed into the

**Eta-Problem = Inflaton-Mass Problem**



Various proposals for its solution were made (warp factors, shift symmetries, etc.) but when analyzed carefully the problem persists [see e.g. McAllister hep-th/0502001].

⇒ Let's try to embed inflation into string- and M-theory through multi-field inflation (more natural in view of the multitude of scalars)?

### Power-Law and Assisted Inflation

To this end, let us emphasize that there are various different types of inflation through which one might try to embed inflation into string-theory:

**New Inflation:** is based on a vacuum energy dominated de Sitter expansion for which

$$a(t) = a_0 e^{Ht}$$

Many efforts concentrated on embedding inflation into string-theory via new inflation (or hybrid inflation)

**Power-Law Inflation:**  
relies on an exponential potential

[Lucchin, Matarrese 1985]

$$U(\varphi) = U_0 e^{-\sqrt{\frac{2}{p}} \frac{\varphi}{M_{Pl}}}$$



with parameter  $p > 1$  leading to a scale-factor

$$a(t) = a_0 t^p$$

and an evolution of the inflaton

$$\varphi(t) = \sqrt{2p} M_{Pl} \ln \left( \sqrt{\frac{U_0}{p(3p-1) M_{Pl}^2}} t \right)$$

(solution is valid for  $p > 1/3$  but inflation arises only if  $p > 1 \Leftrightarrow \ddot{a} > 0$ )

power-law inflation implies very simple constant slow-roll parameters

$$\epsilon = \frac{1}{p} \quad \eta = \frac{2}{p}$$

**constant slow-roll parameters** mean there is no exit from power-law inflation. When embedded into M/string theory this presents, however, no problem as additional contributions will eventually modify the simple exponential potential causing inflation to end.

exponential potentials arise naturally from various **non-perturbative effects** which need to be included anyway for moduli stabilization (and spontaneous supersymmetry breaking)



**Single** D-brane or Membrane instantons lead however to  $p = \mathcal{O}(1)$  which is not sufficient for a sustained period of inflation!

So what can be done to obtain inflation nevertheless?

**Assisted Inflation:**

[Liddle, Mazumdar, Schunck 1998]

consider instead a multi-inflaton extension of the single scalar power-law inflation scenario, termed assisted inflation

each of the  $N$  scalar fields  $\varphi_i$ ,  $i = 1, \dots, N$ , has a potential

$$U = U_0 e^{-\sqrt{\frac{2}{p}} \frac{\varphi_i}{M_{Pl}}}, \quad \forall i = 1, \dots, N.$$

(for  $p = \mathcal{O}(1)$  as in M/string theory, individual potentials too steep to give power-law inflation) Since all scalars obey identical dynamics, one can map this multi-field problem by rescaling to the single field power-law problem and show that it gives again a power-law solution

$$a(t) = a_0 t^{p(N)}$$

but this time with

$$p(N) = Np$$

$\Rightarrow$  even though single exponential contributions are too steep to support inflation individually, it is easy to



obtain large  $p(N) \gg 1$  by increasing  $N$

$\Rightarrow$  this mechanism gives inflation with small  $\epsilon, \eta$  by using many exponential potentials which would usually be discarded as too steep when considered individually (increased Hubble-friction experienced by every individual scalar)!

Is it possible to embed inflation via the method of assisted inflation into M/string-theory, thereby using the naturally available "too steep" potentials?

## Realizing Assisted Inflation in M-Theory

[K. & M. Becker, AK '05]


### 1) The Multi M5-Brane Potential

For definiteness and because of its direct contact with a realistic GUT or MSSM sector [see e.g. B. Ovrut et al. '04] – which is important for the issue of reheating – we will focus subsequently on heterotic M-theory.

When compactified down to 4d on a 6-dim. manifold preserving  $N = 1$  supersymmetry, background is given either by a warped Calabi-Yau threefold or a warped non-Kähler manifold depending on type of flux. But warp-factor along  $S^1/Z_2$  stays the same [Curio, AK 2000, 2003]



Let's focus on the CY case and consider the following setup of  $N$  parallel M5-branes distributed along the  $S^1/Z_2$  interval (all M5-branes fill 4d spacetime and wrap same genus zero holomorphic 2-cycle on the CY; for simplicity  $h^{1,1} = 1$ , i.e. one Kähler modulus  $T$  only).



### Moduli and Kähler-Potential

effective 4d  $N=1$  supergravity is described in terms of

- $h^{2,1}$  complex structure moduli  $Z^\alpha$
- CY volume modulus  $S$
- $T$  modulus measuring  $S^1/Z_2$  length
- M5-brane position fields  $Y_i$

defined as

$$S = \mathcal{V} + \mathcal{V}_{OM} \sum_{i=1}^N \left( \frac{x_i^{11}}{L} \right)^2 + i\sigma_S$$

$$T = \mathcal{V}_{OM} + i\sigma_T$$

$$Y_i = \mathcal{V}_{OM} \left( \frac{x_i^{11}}{L} \right) + i\sigma_i, \quad i = 1, \dots, N$$

where

- $\mathcal{V}$  = average CY volume over  $S^1/\mathbb{Z}_2$
- $\mathcal{V}_{OM}$  = average volume of 3-cycle  $\Sigma_2 \times S^1/\mathbb{Z}_2$
- $L$  = length of  $S^1/\mathbb{Z}_2$  interval
- $0 \leq x_i^{11} \leq L$ : position of the  $i$ th M5-brane

in addition it's useful to define

$$s = S + \bar{S}, \quad t = T + \bar{T}, \quad y_i = Y_i + \bar{Y}_i,$$

$$y = \left( \sum_{i=1}^N y_i^2 \right)^{1/2}$$

plus


$$Q = s - \frac{y^2}{t}$$

$$R = 3Q^2 - 2\frac{y^4}{t^2}$$

such that the Kähler-potential becomes



Let's focus on the CY case and consider the following setup of  $N$  parallel M5-branes distributed along the  $S^1/\mathbb{Z}_2$  interval (all M5-branes fill 4d spacetime and wrap same genus zero holomorphic 2-cycle on the CY; for simplicity  $h^{1,1} = 1$ , i.e. one Kähler modulus  $T$  only).



### Moduli and Kähler-Potential

effective 4d  $N=1$  supergravity is described in terms of

- $h^{2,1}$  complex structure moduli  $Z^\alpha$
- CY volume modulus  $S$
- $T$  modulus measuring  $S^1/\mathbb{Z}_2$  length
- M5-brane position fields  $Y_i$

defined as



$$K = K_{(S)} + K_{(T)} + K_{(Y)} + K_{(Z)}$$

where

$$K_{(S)} + K_{(Y)} = -\ln Q$$

$$K_{(T)} = -\ln \left( \frac{d}{6} t^3 \right)$$

$$K_{(Z)} = -\ln \left( i \int_{CY} \Omega \wedge \overline{\Omega} \right),$$

( $d$  = CY intersection number)

Obviously  $Q = 2V > 0$ ; likewise  $R > 0$  to ensure a positive definite Kähler metric  $K_{I\bar{J}}$  for which one finds ( $I, J, \dots$  run over all complex moduli,  $G_{\alpha\bar{\beta}}$  = metric on the complex structure moduli space)


$$\det K_{I\bar{J}} = \frac{16R}{Q^{2N} t^6} \det G_{\alpha\bar{\beta}}$$

### Which Superpotentials Need to be Considered?

a priori contributions to superpotential come from open membrane instantons (wrapping same genus zero curve as M5's) stretching between:

- both boundaries (99),
- between two of the M5-branes (55),
- between the visible boundary and an M5-brane (95)
- or between an M5-brane and the hidden boundary (59)





$$W_{OM} = W_{99} + W_{55} + W_{95} + W_{59}$$

where

$$W_{99} = h e^{-T}, \quad W_{95} = h \sum_{i=1}^N e^{-Y_i}, \quad W_{59} = h \sum_{i=1}^N e^{-(T-Y_i)}$$

$$W_{55} = h \sum_{i < j} e^{-Y_{ji}}$$

with

$$Y_{ji} = Y_j - Y_i$$

describing the distance between the  $j$ th and the  $i$ th M5-brane

When  $T$  would have been stabilized at the critical length [Curio, AK 2001; M. Becker, Curio, AK 2004] where volume (gauge coupling) of hidden boundary becomes small (large), then one would also consider gaugino condensation on

$$S = \mathcal{V} + \mathcal{V}_{OM} \sum_{i=1}^N \left( \frac{x_i^{11}}{L} \right)^2 + i\sigma_S$$

$$T = \mathcal{V}_{OM} + i\sigma_T$$

$$Y_i = \mathcal{V}_{OM} \left( \frac{x_i^{11}}{L} \right) + i\sigma_i, \quad i = 1, \dots, N$$

where

- $\mathcal{V}$  = average CY volume over  $S^1/Z_2$
- $\mathcal{V}_{OM}$  = average volume of 3-cycle  $\Sigma_2 \times S^1/Z_2$
- $L$  = length of  $S^1/Z_2$  interval
- $0 \leq x_i^{11} \leq L$ : position of the  $i$ th M5-brane

in addition it's useful to define

$$s = S + \bar{S}, \quad t = T + \bar{T}, \quad y_i = Y_i + \bar{Y}_i,$$

$$y = \left( \sum_{i=1}^N y_i^2 \right)^{1/2}$$

plus

$$Q = s - \frac{y^2}{t}$$

$$R = 3Q^2 - 2\frac{y^4}{t^2}$$

such that the Kähler-potential becomes



Among the open membrane contributions, we can focus on the **dominant nearest neighbor 55 contributions**. This is correct as long as nearest M5 brane distances are smaller than the orbifold size implying that the neglected OM instantons have stretch over longer distances. Hence

$$W = W_{55}$$

### M5-brane Interaction Potential

Potential follows from standard F-term expression

$$U = M_{Pl}^4 e^K \left( \sum K^{\bar{I}J} D_{\bar{I}} \overline{W}_{55} D_J W_{55} - 3 |W_{55}|^2 \right),$$

which leads to

$$\begin{aligned} \frac{U}{M_{Pl}^4 e^K} &= G^{\bar{\alpha}\beta} D_{\bar{\alpha}} W_{55} D_{\beta} W_{55} \\ &+ Qt \sum_{i,j=1}^N \left( \frac{1}{2} \delta_{ij} + \frac{Q}{Rt} y_i y_j \right) \overline{D_i W_{55}} D_j W_{55} \\ &+ \left( \frac{3Q^2}{R} - \frac{2y^2}{Qt} \right) |W_{55}|^2 \end{aligned}$$

with Kähler factor  $e^K = 6/(i \int \Omega \wedge \overline{\Omega}) Qt^3 d$



the only term which in principle could become negative is the **last term** which arises from

$$\sum K^{\bar{I}J} K_{\bar{I}} K_J |W_{55}|^2 - 3|W_{55}|^2$$

It is however easy to check that  $3Q^2/R > 2y^2/Qt$  when  $Q > 0$  and  $R > 0$ , as required for a positive definite Kähler metric. Hence this term and therefore the whole potential will be **positive** – an important requisite for the derivation of assisted inflation!

## 2) Mapping M5-Brane Dynamics to Assisted Inflation Dynamics

Since M5-brane interaction potential is positive we can **partially minimize** it by demanding

$$D_\alpha W_{55} = 0$$

$$D_{\bar{i}} W_{55} = 0$$

Let us see what they imply. The first equation is equivalent to

$$\frac{\partial \ln h}{\partial Z^\alpha} = -\frac{\partial K_{(Z)}}{\partial Z^\alpha}$$

and implies

$$h = i \int_{CY} \Omega \wedge \bar{\Omega}$$

It will fix the  $h^{2,1}$  complex structure moduli



The second equation has a simple geometric meaning: in the large volume regime, where we can trust the supergravity analysis, we have  $Qt \simeq st \gg t > y_i$  and therefore

$$0 = D_i W_{55} = W_{55,i} + \frac{2y_i}{Qt} W_{55} \rightarrow W_{55,i}$$

With

$$W_{55} = h \sum_{i=1}^{N-1} e^{-Y_{i+1,i}}$$

this implies

$$Y_{i+1,i} \equiv \Delta Y \quad \forall i$$

Hence partially minimizing the energy through setting  $D_i W_{55} = 0$  forces the inter M5-brane distances to be **equidistant**

### Specifying the Regime where Inflation Occurs

So far, by partially minimizing the energy, we have arrived at

$$U \propto \left( \frac{3Q}{Rt^3} - \frac{2y^2}{Q^2 t^4} \right) |W_{55}|^2$$

If this potential is to be mapped to an assisted inflation dynamics with the inflatons arising from the M5-brane position differences  $y_{i+1} - y_i$ , we have to make sure that



there is a  $y_i$  dependence only in the exponentials of  $W_{55}$   
 can be achieved by working in the regime where

$$Qt \gg y^2$$

which is consistent with large volume  $Q \simeq s \gg 1$  and  
 implies  $3Q/Rt^3 \gg 2y^2/Q^2t^4$ . In this regime we simply  
 have

$$U \propto \frac{1}{st^3} |W_{55}|^2$$

and therefore finally

$$\frac{Ud}{6M_{Pl}^4(i \int \Omega \wedge \bar{\Omega})} = \frac{(N-1)^2}{st^3} e^{-\Delta y}, \quad \Delta y = \Delta Y + \overline{\Delta Y}$$

## The Mapping

1) Transformation to canonically normalized scalars:  
 $Y_i$  kinetic term

$$S_{kin} = -M_{Pl}^2 \int d^4x \sqrt{-g} K_{i\bar{j}} \partial_\mu Y_i \partial^\mu \bar{Y}_{\bar{j}},$$

where

$$K_{i\bar{j}} = \frac{4y_i y_j + 2Qt \delta_{ij}}{Q^2 t^2}.$$

In the regime which we had just specified, we have  
 $Qt \gg y^2 = \sum y_i^2 > y_i y_j$ . Thus (under the sum) we can



The second equation has a simple geometric meaning: in the large volume regime, where we can trust the supergravity analysis, we have  $Qt \simeq st \gg t > y_i$  and therefore

$$0 = D_i W_{55} = W_{55,i} + \frac{2y_i}{Qt} W_{55} \rightarrow W_{55,i}$$

With

$$W_{55} = h \sum_{i=1}^{N-1} e^{-Y_{i+1,i}}$$

this implies

$$Y_{i+1,i} \equiv \Delta Y \quad \forall i$$

Hence partially minimizing the energy through setting  $D_i W_{55} = 0$  forces the inter M5-brane distances to be **equidistant**

### Specifying the Regime where Inflation Occurs

So far, by partially minimizing the energy, we have arrived at

$$U \propto \left( \frac{3Q}{Rt^3} - \frac{2y^2}{Q^2 t^4} \right) |W_{55}|^2$$

If this potential is to be mapped to an assisted inflation dynamics with the inflatons arising from the M5-brane position differences  $y_{i+1} - y_i$ , we have to make sure that



there is a  $y_i$  dependence only in the exponentials of  $W_{55}$

can be achieved by working in the regime where

$$Qt \gg y^2$$

which is consistent with large volume  $Q \simeq s \gg 1$  and implies  $3Q/Rt^3 \gg 2y^2/Q^2t^4$ . In this regime we simply have

$$U \propto \frac{1}{st^3} |W_{55}|^2$$

and therefore finally

$$\frac{Ud}{6M_{Pl}^4(i \int \Omega \wedge \bar{\Omega})} = \frac{(N-1)^2}{st^3} e^{-\Delta y}, \quad \Delta y = \Delta Y + \overline{\Delta Y}$$

### The Mapping

1) Transformation to canonically normalized scalars:

$Y_i$  kinetic term

$$S_{kin} = -M_{Pl}^2 \int d^4x \sqrt{-g} K_{ij} \partial_\mu Y_i \partial^\mu \bar{Y}_j,$$

where

$$K_{ij} = \frac{4y_i y_j + 2Qt \delta_{ij}}{Q^2 t^2}.$$

In the regime which we had just specified, we have  $Qt \gg y^2 = \sum y_i^2 > y_i y_j$ . Thus (under the sum) we can



neglect the first piece and obtain  $K_{ij} = 2\delta_{ij}/Qt$  which leads to the following canonically normalized real M5-brane position and difference fields

$$\phi_i = \frac{2M_{Pl}}{\sqrt{Qt}} y_i, \quad \Delta\phi = \frac{2M_{Pl}}{\sqrt{Qt}} \Delta y$$

## 2) Switching to COM and Relative Coordinates:

A potential is only generated for distances between adjacent M5-branes but not for their combined com position. Let us therefore switch from the  $N$  position fields  $\phi_i$  to the more adequate description in terms of the M5-brane com field

$$\phi_{com} = \frac{1}{N}(\phi_1 + \dots + \phi_N),$$

and the difference field  $\Delta\phi$ . The relation between the two sets of fields is provided by the relation

$$\phi_i = \phi_{com} + \left(i - \frac{N+1}{2}\right) \Delta\phi$$

Since there is no potential for  $\phi_{com}$ , its value will stay constant and its kinetic term vanishes. The sum of the



$\phi_i$  kinetic terms then becomes

$$\begin{aligned} \frac{1}{2} \sum_{i=1}^N \partial_\mu \phi_i \partial^\mu \phi_i &= \partial_\mu \Delta \phi \partial^\mu \Delta \phi \sum_{i=1}^N \left( i - \frac{N+1}{2} \right)^2 \\ &= \frac{N(N^2-1)}{12} \partial_\mu \Delta \phi \partial^\mu \Delta \phi \end{aligned}$$

requiring us to perform a **second renormalization**

Eventually, we arrive at the **final canonically normalized difference field**  $\varphi$

$$\varphi = \sqrt{\frac{N(N^2-1)}{6}} \Delta \phi = M_{Pl} \sqrt{\frac{2N(N^2-1)}{3Qt}} \Delta y$$

in terms of which the **potential** reads

$$U(\varphi) = \tilde{U}_0 (N-1)^2 e^{-\sqrt{\frac{3Qt}{2N(N^2-1)}} \frac{\varphi}{M_{Pl}}}$$

with  $\tilde{U}_0 = 6M_{Pl}^4 (i \int \Omega \wedge \bar{\Omega}) / st^3 d$  (the approximate constancy of  $s, t$  will be discussed shortly)

For a spatially flat 4d FRW universe we then have a **Hubble parameter**

$$H^2 = \frac{1}{3M_{Pl}^2} \left( U(\varphi) + \frac{1}{2} \dot{\varphi}^2 \right),$$

and the dynamics of  $\varphi$  is determined by

$$\ddot{\varphi} + 3H\dot{\varphi} + \frac{dU}{d\varphi} = 0.$$



This is precisely the dynamics which gives **power-law inflation** once we identify the parameters

$$p = \frac{4N(N^2 - 1)}{3Qt}$$

$$U_0 = \tilde{U}_0(N - 1)^2$$

This completes the mapping and therefore embedding of assisted inflation into M-theory.

### Bounds on $N$

Contrary to what one might think, the value for  $N$  is rather constrained

1) We had the condition

$$Qt \gg y^2$$

which implies an **upper bound** on  $N$  as  $y$  grows with  $N$ . For typical values  $\mathcal{V} = 341$ ,  $\mathcal{V}_{OM} = 7$  and  $x_i^{11}/L = \mathcal{O}(1/2)$  we have  $s = 682 + 3.5N$ ,  $t = 14$ ,  $y^2 \simeq 49N$  which leads to

$$N \ll 195$$

2) to obtain inflation we need  $p > 1$  which implies a **lower bound** on  $N$

$$p > 1 \quad \Leftrightarrow \quad 4N(N^2 - 1) > 3Qt$$

For the same  $s, t$  values as before we get

$$19 < N$$




There is thus a non-empty set of  $N$ 's satisfying both constraints. Moreover also observational constraints coming from the number of e-foldings or the scalar spectral index require values for  $N$  within same range!

### Moduli Stabilization

There is a crucial difference between new inflaton models and this assisted inflation model for the embedding into M/string-theory

New inflaton models have to select a single very flat direction. Necessarily one has to stabilize all other moduli before inflation.

Here, we have made use of the very steepest directions available. The universe is rolling down exponentially steep directions during inflation





(up to one) moduli before inflation. Mild, i.e. power-law runaways might be tolerable  $(s, t)$

### The Whole Evolution: Cascade Inflation

So far we have kept  $N = \#$  of M5-branes fixed and analyzed therefore only part of the full process. Since the M5-brane distances are growing  $N$  will jump to  $N - 2$  as soon as the two outermost M5-branes hit the boundaries


This can also be seen from the anomaly cancellation equation for the  $G$  flux

$$\beta_v + \beta_h + N = 0$$

( $\beta_{v,h}$  are integers characterizing the boundary's 2nd Chern classes). When the M5-branes coalesce with the boundaries they change the boundaries topological data via small instanton transitions which has to be compensated for by a change in  $N$

Hence, we have the following "cascade-like" evolution:





For each interval we obtain power-law inflation via the same derivation as before, as long as

$$p(N_q) > 1 \quad (\text{Exit Condition : } p(N_q) = 1)$$

(2nd constraint  $Qt \gg y^2$  always remains satisfied as  $y$  decreases with decreasing  $N$ )

The full "cascade inflation" process has the following structure

$$\begin{aligned} a_1(t) &= a_1 t^{p_1}, & t_0 \leq t \leq t_1 \\ a_2(t) &= a_2 t^{p_2}, & t_1 \leq t \leq t_2 \\ &\vdots \\ a_q(t) &= a_q t^{p_q}, & t_{q-1} \leq t \leq t_q \\ &\vdots \end{aligned}$$

Matching them at the transition times  $t_q$  determines the constant prefactors

$$a_q = a_1 t_1^{p_1} \left( \frac{t_2}{t_1} \right)^{p_2} \left( \frac{t_3}{t_2} \right)^{p_3} \dots \left( \frac{t_{q-1}}{t_{q-2}} \right)^{p_{q-1}} \frac{1}{t_{q-1}^{p_q}}$$



## Number of E-Foldings

$$N_e \equiv \ln \left( \frac{a(t_f)}{a(t_0)} \right) = \sum_{q=1}^{q_f} p_q \ln \left( \frac{t_q}{t_{q-1}} \right)$$

with

$$p_q = \frac{4N_q(N_q^2 - 1)}{3st}$$

the constant ratios  $t_q/t_{q-1}$  can be determined by using the exact solution for the inflaton (distances between adjacent M5-branes) evolution

$$\begin{aligned} \frac{t_q}{M_{Pl}} \simeq \frac{t_q - t_0}{M_{Pl}} &= \frac{1}{\sqrt{U_0}} \sum_{a=1}^q \frac{p_a(3p_a - 1)}{N_a - 1} e^{t(\frac{\Delta x_a}{L} - \frac{\Delta x_{q-1}}{L})} \\ &= \frac{1}{\sqrt{U_0}} \sum_{a=1}^q \frac{p_a(3p_a - 1)}{N_a - 1} e^{t(\frac{1}{N_a - 1} - \frac{1}{N_{a-1} - 1})} \end{aligned}$$

Taking typical values for  $s, t$  as before, this series can be summed numerically.

$$\text{Exit from Inflation: } p(N_{q_f}) \stackrel{!}{=} 1 \Rightarrow N_{q_f} = 19$$

Knowing  $N_{q_f}$ , we can carry out the summation and obtain

$$N = 40 \Rightarrow N_e = 13.3$$

$$N = 50 \Rightarrow N_e = 28.6$$

$$N = 60 \Rightarrow N_e = 53.2$$

$$N = 70 \Rightarrow N_e = 89.7$$

$\Rightarrow$  Since  $19 \leq N \ll 195$ , we see that a realistic  $N_e = 50 - 60$  can be obtained within required regime!