

Title: Perturbative Theory of the Phase Structure of Very Dense QCD Matter

Date: May 25, 2007 10:20 AM

URL: <http://pirsa.org/07050074>


Abstract:

Phase Structure of (Very) Dense Quark Matter

Thomas Schaefer

North Carolina State

Schematic Phase Diagram



Dense Baryonic Matter

Low Density



Equation of state of nuclear/neutron matter

Neutron/proton superfluidity, pairing gaps

Moderate Density

Pion/kaon condensation, hyperon matter

Pairing, equation of state at high density

High Density

Quark matter


Color superconductivity, Color-flavor-locking

Dense Baryonic Matter

Low Density	(constrained by NN interaction, phenomenology)
	Equation of state of nuclear/neutron matter
	Neutron/proton superfluidity, pairing gaps
Moderate Density	(very poorly known)
	Pion/kaon condensation, hyperon matter
	Pairing, equation of state at high density
High Density	(weak coupling methods apply)
	Quark matter
	Color superconductivity, Color-flavor-locking

Pairing in Quark Matter

QQ scattering in perturbative QCD




$$(\vec{T})_{ac}(\vec{T})_{bd} = -\frac{1}{3}(\delta_{ab}\delta_{cd} - \delta_{ad}\delta_{bc}) + \frac{1}{6}(\delta_{ab}\delta_{cd} + \delta_{ad}\delta_{bc})$$

[3] \times [3] = [3] + [6]

Fermi surface: pairing instability in weak coupling

$$\Phi_{ij}^{ab,\alpha\beta} = \langle \psi_i^{a,\alpha} C \psi_j^{b,\beta} \rangle$$



Phase structure in perturbation theory

$$\text{Minimize } \Omega(\Phi_{ij}^{ab,\alpha\beta})$$

In practice: consider $\Phi_{ij}^{ab,\alpha\beta}$ with residual symmetries

Superconductivity

Thermodynamic potential

$$\Omega = \text{Diagram A} + \text{Diagram B} + \text{Diagram C} + \dots$$

Variational principle $\delta\Omega/\delta\Phi$ gives gap equation

$$= \frac{g^2}{18\pi^2} \int dq_0 \log \left(\frac{\Lambda_{BCS}}{|p_0 - q_0|} \right) \frac{\Delta(q_0)}{\sqrt{q_0^2 + \Delta(q_0)^2}}$$

$\Lambda_{BCS} = c_i 256\pi^4 \mu g^{-5}$ determined by symmetries of order parameter

$$\Omega = \frac{\mu^2}{4\pi^2} \sum_i \Delta_i^2 \quad \Delta_i = c_i \mu g^{-5} \exp \left(-\frac{3\pi^2}{\sqrt{2}g} \right)$$

Remarks

Behavior of perturbative expansion

$$\Delta = \mu g^{-5} \exp\left(-\frac{c}{g}\right) \left(c_0 + c_1 g \log(g) + c_2 g + O(g^{n/3} \log^m(g))\right)$$

Also: non-perturbative effects (become large for $\mu \sim 1$ GeV)

$$\Delta \sim \mu \exp\left(-\frac{8\pi^2}{g^2}\right)$$

Magnitude of gap quite uncertain

But: Phase Structure not sensitive to interaction

$N_f = 1$: Color-Spin-Locking



$N_f = 1$, color-anti-symmetric: spin-1 condensate

$$\langle \psi^b C \gamma_i \psi^c \rangle = \Phi_i^a \epsilon^{abc}$$

Ground state: Color-Spin-Locking (CSL): $\Phi_i^a \sim \delta_i^a$

$SU(3)_c \times SO(3) \rightarrow SO(3)$: rotational symmetry

$U(1)_B$ broken: massless Goldstone boson

($s=3/2$) + ($s=1/2$) gapped fermions

$N_f = 2$: 2SC Phase

$N_f = 2$, color-anti-symmetric: spin-0 BCS condensate

$$\langle \psi_i^b C \gamma_5 \psi_j^c \rangle = \Phi^a \epsilon^{abc} \epsilon_{ij}$$

Order parameter $\phi^a \sim \delta^{a3}$ breaks $SU(3)_c \rightarrow SU(2)$

$SU(2)_L \times SU(2)_R$ unbroken

4 gapped, 2 (almost) gapless fermions

light $U(1)_A$ Goldstone boson

$SU(2)$ confined ($\Lambda_{conf} \ll \Delta$)

$N_f = 3$: CFL Phase

Consider $N_f = 3$ ($m_i = 0$)

$$\langle q_i^a q_j^b \rangle = \phi \epsilon^{abI} \epsilon_{ijI}$$



$$\langle ud \rangle = \langle us \rangle = \langle ds \rangle$$

$$\langle rb \rangle = \langle rg \rangle = \langle bg \rangle$$


Symmetry breaking pattern:

$$SU(3)_L \times SU(3)_R \times [SU(3)]_C \\ \times U(1) \rightarrow SU(3)_{C+F}$$

All quarks and gluons acquire a gap
 $[8] + [1]$ fermions, Q integer



Rotate left flavor Compensate by rotating color



... have to rotate right flavor also !

$$\langle \psi_L \psi_L \rangle = -\langle \psi_R \psi_R \rangle$$

QCD with many flavors/colors


QCD with $N_f > 3$ flavors: CFL-like, fully gapped, phases

$$N_f = 4 \quad SU(4)_L \times SU(4)_R \times U(1)_V \rightarrow [SU(2)_V]^2 \times SU(2)_A$$

$$N_f = 5 \quad SU(5)_L \times SU(5)_R \times U(1)_V \rightarrow SU(2)_V$$

$$N_f = 6 \quad SU(6)_L \times SU(6)_R \times U(1)_V \rightarrow SU(3)_V \times U(1)_V \times U(1)_A$$

$N_c \rightarrow \infty$: qq pairing suppressed, qq^{-1} chiral density wave



$$\langle \bar{q}(x)q(x) \rangle \sim \Sigma e^{2i\vec{q}_F \cdot \vec{x}}$$

Requires very large $N_c > 1000$

$N_f = 3$: CFL Phase

Consider $N_f = 3$ ($m_i = 0$)

$$\langle q_i^a q_j^b \rangle = \phi \epsilon^{abI} \epsilon_{ijI}$$


$$\langle ud \rangle = \langle us \rangle = \langle ds \rangle$$

$$\langle rb \rangle = \langle rg \rangle = \langle bg \rangle$$

Symmetry breaking pattern:

$$SU(3)_L \times SU(3)_R \times [SU(3)]_C \\ \times U(1) \rightarrow SU(3)_{C+F}$$

All quarks and gluons acquire a gap
 $[8] + [1]$ fermions, Q integer



$$\langle \psi_L \psi_L \rangle = -\langle \psi_R \psi_R \rangle$$

QCD with many flavors/colors


QCD with $N_f > 3$ flavors: CFL-like, fully gapped, phases

$$N_f = 4 \quad SU(4)_L \times SU(4)_R \times U(1)_V \rightarrow [SU(2)_V]^2 \times SU(2)_A$$

$$N_f = 5 \quad SU(5)_L \times SU(5)_R \times U(1)_V \rightarrow SU(2)_V$$

$$N_f = 6 \quad SU(6)_L \times SU(6)_R \times U(1)_V \rightarrow SU(3)_V \times U(1)_V \times U(1)_A$$

$N_c \rightarrow \infty$: qq pairing suppressed, qq^{-1} chiral density wave




$$\langle \bar{q}(x)q(x) \rangle \sim \Sigma e^{2i\vec{q}_F \cdot \vec{x}}$$

Requires very large $N_c > 1000$

Towards the real world: Non-zero strange quark mass

Have $m_s > m_u, m_d$: Unequal Fermi surfaces



$$\delta p_F \simeq \frac{m_s^2}{2p_F}$$

Also: If $p_F^s < p_F^{u,d}$ have unequal densities

Charge neutrality not automatic


Strategy

Consider $N_f = 3$ at $\mu \gg \Lambda_{QCD}$ (CFL phase)

Study response to $m_s \neq 0$

Constrained by chiral symmetry

Very Dense Matter: Effective Field Theories




High Density Effective Theory

Effective field theory on v -patches



$$\psi_{v\pm} = e^{-i\mu v \cdot x} \left(\frac{1 \pm \vec{\alpha} \cdot \vec{v}}{2} \right) \psi$$



Effective lagrangian for $p_0 < m$


$$\mathcal{L} = \psi_v^\dagger \left(i v \cdot D - \frac{D_\perp^2}{2\mu} \right) \psi_v + \mathcal{L}_{4f} - \frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a + \mathcal{L}_{HDL}$$

$$\mathcal{L}_{HDL} = -\frac{m^2}{2} \sum_v G_{\mu\alpha}^a \frac{v^\alpha v^\beta}{(v \cdot D)^2} G_{\mu\beta}^b$$

Mass Terms: Match HDET to QCD

$$\mathcal{L} = \psi_R^\dagger \frac{MM^\dagger}{2\mu} \psi_R + \psi_L^\dagger \frac{M^\dagger M}{2\mu} \psi_L$$

$$+ \frac{V_M^0}{\mu^2} (\psi_R^\dagger M \lambda^a \psi_L) (\psi_R^\dagger M \lambda^a \psi_L)$$



mass corrections to FL parameters $\hat{\mu}_{L,R}$ and $V^0(RR \rightarrow LL)$

EFT in the CFL Phase

Consider HDET with a CFL gap term

$$\mathcal{L} = \text{Tr} \left(\psi_L^\dagger (iv \cdot D) \psi_L \right) + \frac{\Delta}{2} \left\{ \text{Tr} (X^\dagger \psi_L X^\dagger \psi_L) - \kappa [\text{Tr} (X^\dagger \psi_L)]^2 \right\} + (L \leftrightarrow R, X \leftrightarrow Y)$$


$$\psi_L \rightarrow L \psi_L C^T, \quad X \rightarrow L X C^T, \quad \langle X \rangle = \langle Y \rangle = \mathbb{1}$$

Quark loops generate a kinetic term for X, Y

Integrate out gluons, identify low energy fields ($\xi = \Sigma^{1/2}$)


$$\Sigma = XY^\dagger$$

[8]+[1] GBs



$$N_L = \xi (\psi_L X^\dagger) \xi^\dagger$$

[8]+[1] Baryons



Effective theory: (CFL) baryon chiral perturbation theory

$$\begin{aligned}
\mathcal{L} = & \frac{f_\pi^2}{4} \left\{ \text{Tr} \left(\nabla_0 \Sigma \nabla_0 \Sigma^\dagger \right) - v_\pi^2 \text{Tr} \left(\nabla_i \Sigma \nabla_i \Sigma^\dagger \right) \right\} \\
& + A \left\{ [\text{Tr} (M \Sigma)]^2 - \text{Tr} (M \Sigma M \Sigma) + h.c. \right\} \\
& + \text{Tr} (N^\dagger i v^\mu D_\mu N) - D \text{Tr} (N^\dagger v^\mu \gamma_5 \{\mathcal{A}_\mu, N\}) \\
& - F \text{Tr} (N^\dagger v^\mu \gamma_5 [\mathcal{A}_\mu, N]) + \frac{\Delta}{2} \left\{ \text{Tr} (NN) - [\text{Tr} (N)]^2 \right\}
\end{aligned}$$

$$\nabla_0 \Sigma = \partial_0 \Sigma + i \hat{\mu}_L \Sigma - i \Sigma \hat{\mu}_R$$

$$D_\mu N = \partial_\mu N + i[\mathcal{V}_\mu, N]$$

$$f_\pi^2 = \frac{21 - 8 \log 2}{18} \frac{\mu^2}{2\pi^2} \quad v_\pi^2 = \frac{1}{3} \quad A = \frac{3\Delta^2}{4\pi^2} \quad D = F = \frac{1}{2}$$

Phase Structure and Spectrum

Phase structure determined by effective potential

$$V(\Sigma) = \frac{f_\pi^2}{2} \text{Tr} (\hat{\mu}_L^\dagger \Sigma \hat{\mu}_R \Sigma^\dagger) - A \text{Tr}(M \Sigma^\dagger) - B_1 [\text{Tr}(M \Sigma^\dagger)]^2 + \dots$$


$$V(\Sigma_0) \equiv \min$$

Fermion spectrum determined by

$$\mathcal{L} = \text{Tr} (N^\dagger i v^\mu D_\mu N) + \text{Tr} (N^\dagger \gamma_5 \rho_A N) + \frac{\Delta}{2} \left\{ \text{Tr} (NN) - [\text{Tr} (N)]^2 \right\},$$

$$\rho_{V,A} = \frac{1}{2} \{ \xi \hat{\mu}_L \xi^\dagger \pm \xi^\dagger \hat{\mu}_R \xi \} \quad \xi = \sqrt{\Sigma_0}$$

Phase Structure of CFL Phase




$$m_s^{crit} \sim 3.03 m_d^{1/3} \Delta^{2/3}$$

$$m^* \sim 0.017 \alpha_s^{4/3} \Delta$$

QCD realization of s-wave meson condensation

Driven by strangeness oversaturation of CFL state

Fermion Spectrum



$$m_s^{crit} \sim (8\mu\Delta/3)^{1/2}$$

gapless fermion modes (gCFLK)
(chromomagnetic) instabilities ?

Instabilities

Consider meson current

$$\Sigma(x) = U_Y(x) \Sigma_K U_Y(x)^\dagger \quad U_Y(x) = \exp(i\phi_K(x)\lambda_8)$$

$$\vec{V}(x) = \frac{\vec{\nabla}\phi_K}{4}(-2\hat{I}_3 + 3\hat{Y}) \quad \vec{A}(x) = \vec{\nabla}\phi_K(e^{i\phi_K}\hat{u}^+ + e^{-i\phi_K}\hat{u}^-)$$


Gradient energy

$$\mathcal{E} = \frac{f_\pi^2}{2} v_\pi^2 j_K^2 \quad \vec{j}_K = \vec{\nabla}\phi_K$$



Fermion spectrum

$$\omega_l = \Delta + \frac{l^2}{2\Delta} - \frac{4\mu_s}{3} - \frac{1}{4}\vec{v} \cdot \vec{j}_K$$

$$\mathcal{E} = \frac{\mu^2}{2\pi^2} \int dl \int d\hat{\Omega} \omega_l \Theta(-\omega_l)$$



Energy Functional



$$\left. \frac{3\mu_s - 4\Delta}{\Delta} \right|_{crit} = a_{crit} \quad \left. \frac{JK}{\Delta} \right|_{crit} = c_{crit}$$

current strongly suppressed by electric charge neutrality

$m_s^2 \sim 2\mu\Delta$: multiple currents? crystalline state?

Instabilities

Consider meson current

$$\Sigma(x) = U_Y(x) \Sigma_K U_Y(x)^\dagger \quad U_Y(x) = \exp(i\phi_K(x)\lambda_8)$$

$$\vec{V}(x) = \frac{\vec{\nabla}\phi_K}{4}(-2\hat{I}_3 + 3\hat{Y}) \quad \vec{A}(x) = \vec{\nabla}\phi_K(e^{i\phi_K}\hat{u}^+ + e^{-i\phi_K}\hat{u}^-)$$


Gradient energy

$$\mathcal{E} = \frac{f_\pi^2}{2} v_\pi^2 j_K^2 \quad \vec{j}_K = \vec{\nabla}\phi_K$$


Fermion spectrum

$$\omega_l = \Delta + \frac{l^2}{2\Delta} - \frac{4\mu_s}{3} - \frac{1}{4}\vec{v} \cdot \vec{j}_K$$

$$\mathcal{E} = \frac{\mu^2}{2\pi^2} \int dl \int d\hat{\Omega} \omega_l \Theta(-\omega_l)$$



Fermion Spectrum



$$m_s^{crit} \sim (8\mu\Delta/3)^{1/2}$$

gapless fermion modes (gCFLK)
(chromomagnetic) instabilities ?

Summary

Rich, complicated phase diagram even for $\mu \gg \Lambda_{QCD}$

Depends crucially on N_f, N_c and m_q

Color-flavor-locked (CFL) phase provides weak coupling realization of non-perturbative phenomena

Chiral symmetry breaking, s and p-wave meson condensation

Issues not covered in this talk: Transition to nuclear matter, nuclear exotics, etc.

Constraints from compact star phenomenology

tr
File Edit View Termin
tmschaef@tmschaef ~
tmschaef@tmschaef t
L] 2909
tmschaef@tmschaef t

File Edit View Document Tools Window Help

70% | + - | ? | Search Web | Y!

Pages

Summary

Rich, complicated phase diagram even for $\mu \gg \Lambda_{QCD}$

Depends crucially on N_f, N_c and m_q

Color-flavor-locked (CFL) phase provides weak coupling realization of non-perturbative phenomena

Chiral symmetry breaking, s and p-wave meson condensation

Issues not covered in this talk: Transition to nuclear matter, nuclear exotics, etc.

Constraints from compact star phenomenology

23

| < < | 23 of 23 | > > | + - |

Pirsa: 07050074

Page 29/32

$$\sum_{l=1}^L N_l \approx \sum_{l=1}^L N_p$$

tr
File Edit View Termin
tmschaef@tmschaef ~
tmschaef@tmschaef t
L] 2909
tmschaef@tmschaef t

File Edit View Document Tools Window Help

Close Wind

Pages

Summary

Rich, complicated phase diagram even for $\mu \gg \Lambda_{QCD}$

Depends crucially on N_f , N_c and m_q

Color-flavor-locked (CFL) phase provides weak coupling realization of non-perturbative phenomena

Chiral symmetry breaking, s and p-wave meson condensation

Issues not covered in this talk: Transition to nuclear matter, nuclear exotics, etc.

Constraints from compact star phenomenology

23

23 of 23

Pirsa: 07050074

Page 31/32

Adobe Reader - talk_perimeter.pdf

File Edit View Document Tools Window Help

Close Wind

Pages

Attachments

Comments

Search Web Y!

70%

70% 19 of 23

Phase Structure of CFL Phase

$m_s^{crit} \sim 3.03 m_d^{1/3} \Delta^{2/3}$

$m^* \sim 0.017 \alpha_s^{4/3} \Delta$

QCD realization of s-wave meson condensation

Driven by strangeness oversaturation of CFL state

19