


Title: Searching for stellar counterparts to dark matter halos on the smallest possible scales


Date: Jun 07, 2008 09:00 AM

URL: <http://pirsa.org/08060004>

Abstract:



*Searching for stellar counterparts to  
the lowest mass dark matter halos  
(that are able to host stars)*



*Searching for stellar counterparts to  
the lowest mass dark matter halos  
(that are able to host stars)*


Erik Tollerud, Louie Strigari,  
James Bullock (UC Irvine)

Shane Walsh, Helmut  
Jerjen (Stromlo)

=0.0




=0.0



- Methodology to find the least luminous galaxies
- The results and detection limits -  
Quantifying what we do and don't know;  
Predicting the luminosity function; Their  
spatial distribution
- Prospects for all sky and more sensitive  
searches


=0.0




- Methodology to find the least luminous galaxies
- The results and detection limits -  
Quantifying what we do and don't know;  
Predicting the luminosity function; Their  
spatial distribution
- Prospects for all sky and more sensitive  
searches











No Signal

VGA-1

No Signal

VGA-1

No Signal

VGA-1


# Sloan Digital Sky Survey

9,500 square  
degrees of  
imaging

21,439,777  
stars  $r < 22$ ,  
 $g-r < 1$

$u, g, r, i, z$


1 minute  
exposures  
on 2.5m  
telescope



Leo I



*Resolved stars  
reveal the least  
luminous systems*




Leo I




*Resolved stars  
reveal the least  
luminous systems*

V-band apparent magnitude



# How To Find Resolved Dwarf Galaxies


bright



faint

blue

red




From ARAA, V26, 1968

# How To Find Resolved Dwarf Galaxies


bright

faint



blue

red




blue

red

# How To Find Resolved Dwarf Galaxies


bright

faint



blue


red



# How To Find Resolved Dwarf Galaxies


bright

faint




blue


red



blue


red






**SDSS DR6**  
**Footprint -**  
**9500 deg<sup>2</sup>**


# All stars




# Red Stars, $18 < r < 19.5$




# Red Stars, $18 < r < 19.5$



## Binned array




# Array smoothed with 3' filter









# Bootes







# Bootes




# *Detection and Confirmation of new Milky Way Dwarfs: example, Bootes II*









# *Detection and Confirmation of new Milky Way Dwarfs: example, Bootes II*





# Detection and Confirmation of new Milky Way Dwarfs: example, Bootes II





# Detection and Confirmation of new Milky Way Dwarfs: example, Bootes II














# *Detection and Confirmation of new Milky Way Dwarfs: ???*




# Detection and Confirmation of new Milky Way Dwarfs: ???



MMT/Megacam data



Detection - SDSS




## Dwarfs or clusters?


*Dwarf - dark matter halo*


*Cluster - no dark matter halo*


- traditionally classified by their size and luminosity
- avg distance, age, [Fe/H] measurements don't get us much farther
- Need dark matter to explain kinematics?





# *Detection and Confirmation of new Milky Way Dwarfs: ???*










# Detection and Confirmation of new Milky Way Dwarfs: example, Bootes II












# Detection and Confirmation of new Milky Way Dwarfs: ???



MMT/Megacam data




Detection - SDSS



## Dwarfs or clusters?

*Dwarf - dark matter halo*  
*Cluster - no dark matter halo*

- traditionally classified by their size and luminosity
- avg distance, age, [Fe/H] measurements don't get us much farther
- Need dark matter to explain kinematics?



No Signal

VGA-1


No Signal

VGA-1

## Dwarfs or clusters?

*Dwarf - dark matter halo*  
*Cluster - no dark matter halo*

- traditionally classified by their size and luminosity
- avg distance, age, [Fe/H] measurements don't get us much farther
- Need dark matter to explain kinematics?



No Signal


VGA-1

## Dwarfs or clusters?

*Dwarf - dark matter halo*


*Cluster - no dark matter halo*

- traditionally classified by their size and luminosity
- avg distance, age, [Fe/H] measurements don't get us much farther
- Need dark matter to explain kinematics?



Figures from Zucker et al 2006; Belokurov et al 2007


Ursa Major II



May still have very high M/L if tidally disturbed, but would affect dynamical modeling

Many of the least luminous objects appear distorted

Tides?




Coma Berenices

## Dwarfs or clusters?

*Dwarf - dark matter halo*


*Cluster - no dark matter halo*

- traditionally classified by their size and luminosity
- avg distance, age, [Fe/H] measurements don't get us much farther
- Need dark matter to explain kinematics?




Figures from Zucker  
et al 2006;  
Belokurov et al 2007

Ursa Major II




May still have very high  
M/L if tidally disturbed,  
but would affect  
dynamical modeling




Coma Berenices

Many of the least  
luminous objects  
appear distorted



Tides?



There may be lots more of these least luminous objects



Many of the least luminous objects appear distorted  
Tides?



	$R_{\text{tidal}}$ if mass = $10^7 M_{\text{Sun}}$	Extent (lower limit)	Mass if $r_{\text{tid}} = \text{extent}$
Segue 1	570 pc	30 pc	$1 \times 10^3 M_{\text{sun}}$
Bootes II	845 pc	82 pc	$9 \times 10^3 M_{\text{sun}}$
Wil 1	740 pc	43 pc	$2 \times 10^3 M_{\text{sun}}$
Com Ber	830 pc	80 pc	$8 \times 10^3 M_{\text{sun}}$
Ursa Major II	670 pc	170 pc	$2 \times 10^5 M_{\text{sun}}$


Distances of the 5 least luminous dwarfs are 25 - 45 kpc


Luminosities are ~ few thousand  $L_{\text{Sun}}$

Since 2004: ~20  
 dwarfs around  
 Milky Way and  
 M31 discovered  
 with only 1,000  
 to 100,000 stars  
 (-2.5 >  $M_V$  > -8)  
 (25 < d < 400 kpc)

Previous searches:  
 Willman et al 02, 05a&b  
 Zucker et al 06a&b  
 Belokurov et al 06, 07  
 Martin et al 07, Ibata et al 07

Ongoing searches:  
 Walsh, Willman & Jerjen 08, in prep  
 Koposov et al 08








Since 2004: ~20  
 dwarfs around  
 Milky Way and  
 M31 discovered  
 with only 1,000  
 to 100,000 stars  
 ( $-2.5 > M_V > -8$ )  
 ( $25 < d < 400$  kpc)


Previous searches:  
 Willman et al 02, 05a&b  
 Zucker et al 06a&b  
 Belokurov et al 06, 07  
 Martin et al 07, Ibata et al 07

Ongoing searches:  
 Walsh, Willman & Jerjen 08, in prep  
 Koposov et al 08











No Signal

VGA-1

No Signal

VGA-1







No Signal


VGA-1


No Signal

VGA-1









abc[

WordArt WordA

Graphics

Insert...

Font

Name: Arial


Size: 16 Color: [Color Picker]

B I U S | A<sup>2</sup> A<sub>2</sub>

Toggle Case: aA

Change Slides

[Slide Navigation Icons]



abc|

WordArt WordA

Graphics

Insert...

Font

Name: Arial

Size: 16 Color: [Color Picker]

B I U S | A<sup>2</sup> A<sub>2</sub>

Toggle Case: aA

Change Slides

[Slide thumbnails]



No Transition

perimeter.ppt

16 17 18 19 20

21 22 23 24 25

26 27 28 29 30

31 32 33 34 35

Change Slides

Color scheme: Custom Schei



26 27 28 29 30


31 32 33 34 35

36 37 38 39 40

41 42 43 44 45



perimeter.ppt




26



27




28



29




30




31

- Things to consider:
- 1. Is there a number of threshold?
  - 2. Is there a number of objects?
  - 3. Is there a number of...

32



33



34




35


What does the future hold?  
Deeper, larger sky surveys

- Dark Energy Survey (DES) (2011-2017)
- Dark Energy Survey (DES) (2011-2017)
- LSST (2014) - full of stars, galaxies, supernovae in Galactic plane and nearby, Cloud and globular clusters (2012)

36




37



38




39




40

The end

41



42




43

What does the future hold?  
Deeper, larger sky surveys

- Dark Energy Survey (DES) (2011-2017)
- Dark Energy Survey (DES) (2011-2017)
- LSST (2014) - full of stars, galaxies, supernovae in Galactic plane and nearby, Cloud and globular clusters (2012)

44



45

# *Take home messages*

- The least luminous galaxies are presently the most direct way to learn about the least massive dark matter halos; these galaxies can only be found very nearby.
- With uniform datasets and detection limits, we move toward a robust comparison of the abundance, spatial and mass distribution of luminous dark matter subs with predictions
- There may be 300-1000 luminous subs around the Milky Way; Will, Bootes II, Segue 1 are the big wildcards
- PanStarrs, SkyMapper, Dark Energy Survey and LSST will provide the necessary dataset to test present empirical predictions of the abundance and spatial distribution of luminous subs. LSST will expand known ultra-faint census throughout the Local Group.
- It could become very tricky to disentangle dwarfs from tidal garbage.

No Signal

VGA-1

No Signal

VGA-1

# *Take home messages*

- The least luminous galaxies are presently the most direct way to learn about the least massive dark matter halos; these galaxies can only be found very nearby.
- With uniform datasets and detection limits, we move toward a robust comparison of the abundance, spatial and mass distribution of luminous dark matter subs with predictions
- There may be 300-1000 luminous subs around the Milky Way; Will1, Bootes II, Segue 1 are the big wildcards
- PanStarrs, SkyMapper, Dark Energy Survey and LSST will provide the necessary dataset to test present empirical predictions of the abundance and spatial distribution of luminous subs. LSST will expand known ultra-faint census throughout the Local Group.
- It could become very tricky to disentangle dwarfs from tidal garbage.


# *Take home messages*

- The least luminous galaxies are presently the most direct way to learn about the least massive dark matter halos; these galaxies can only be found very nearby.
- With uniform datasets and detection limits, we move toward a robust comparison of the abundance, spatial and mass distribution of luminous dark matter subs with predictions
- There may be 300-1000 luminous subs around the Milky Way; Will1, Bootes II, Segue 1 are the big wildcards
- PanStarrs, SkyMapper, Dark Energy Survey and LSST will provide the necessary dataset to test present empirical predictions of the abundance and spatial distribution of luminous subs. LSST will expand known ultra-faint census throughout the Local Group.
- It could become very tricky to disentangle dwarfs from tidal garbage.




No Transition

perimeter.ppt




26



27




28



29




30



31

- Regression
1. Is a function of the number of observations
  2. Is a function of the number of variables
  3. Is a function of the number of variables

32



33



34

Graphics

Insert...


Change Slides

35


What does the future hold?  
Deeper, larger sky surveys

- SDSS (2005) - red, 2.5 degree (2000 sq deg)
- SDSS (2005) - red, 2.5 degree (2000 sq deg)
- SDSS (2005) - red, 2.5 degree (2000 sq deg)
- SDSS (2005) - red, 2.5 degree (2000 sq deg)

36



37



38





39

Color scheme

Custom Scheme


40

The end



What does the future hold?  
Deeper, larger sky surveys

- SDSS (2005) - red, 2.5 degree (2000 sq deg)
- SDSS (2005) - red, 2.5 degree (2000 sq deg)
- SDSS (2005) - red, 2.5 degree (2000 sq deg)
- SDSS (2005) - red, 2.5 degree (2000 sq deg)



## Local Group with micro-to-milli arcsecond astrometry

- Internal stellar proper motions
- Dark matter density of dwarfs
- Halo mass, lumpiness from streams and hyper velocity stars
- Angular momentum distribution of outer halo stars
- Systematic proper motions
- Orbits of Galactic satellites and star clusters
- Local Group galaxies [M31, LMC, SMC]

## Local Group with micro-to-milli arcsecond astrometry

- Internal stellar proper motions
  - Dark matter density of dwarfs
  - Halo mass, lumpiness from streams and hyper velocity stars
  - Angular momentum distribution of outer halo stars
- Systematic proper motions
  - Orbits of Galactic satellites and star clusters
  - Local Group galaxies [M31, LMC, SMC]

# Dynamics of Milky Way Dwarf Satellites

Louie Strigari  
University of California, Irvine

Small-Scale Structure of Dark Matter  
Perimeter Institute  
June 7, 2008

Collaborators: James Bullock, Manoj Kaplinghat, Marla Geha, Josh Simon, Beth Willman