

Title: Results from the MiniBooNE Experiment

Date: Jun 03, 2008 09:35 AM

URL: <http://pirsa.org/08060036>

Abstract: We present the results from the MiniBooNE neutrino oscillations search in which no significant excess of events is observed above background in the energy range from 475 MeV to 3000 MeV. For lower energies an excess of events that is not consistent with a two neutrino oscillation model is observed. We present recent advances in the understanding of this excess, including a study of muon and electron neutrinos from the nearby NuMI neutrino source. The techniques used in the first oscillation analysis are discussed as well as those of a recent analysis that combines two different electron neutrino candidate samples with a high statistics muon neutrino sample in the oscillations fit to reduce systematic uncertainties.

## Outline

The experiment and the oscillations result

NC  $\pi^0$  rate measurement

Combining analyses

Compatibility of high  $\Delta m^2$  experiments

Event excess below oscillations analysis threshold

Data from the NuMI beam at MiniBooNE

Summary

## The MiniBooNE Collaboration

A. A. Aguilar-Arevalo, A. O. Bazarin, S. J. Brice, B. C. Brown,  
 L. Bugel, J. Cao, L. Coney, J. M. Conrad, D. C. Cox, A. Curioni,  
 Z. Djuricic, D. A. Finley, B. T. Fleming, R. Ford, F. G. Garcia,  
 G. T. Garvey, J. A. Green, C. Green, T. L. Hart, E. Hawker,  
 R. Imhof, R. A. Johnson, P. Kasper, T. Katon, T. Kobilarcik,  
 I. Kourbanis, S. Koutsoliotas, J. M. Link, Y. Liu, Y. Liu,  
 W. C. Louis, K. B. M. Mahn, W. Marsh, P. S. Martin, G. McGregor,  
 W. Metcalf, P. D. Meyers, F. Mills, G. B. Mills, J. Monroe,  
 C. D. Moore, R. H. Nelson, P. Nienaber, S. Ouedraogo,  
 R. B. Patterson, D. Perevalov, C. C. Polly, E. Prabbs, J. L. Raaf,  
 H. Ray, B. P. Roe, A. D. Russell, V. Sandberg, R. Schirato,  
 D. Schmitz, M. H. Shaevitz, F. C. Shoemaker, D. Smith, M. Sorel,  
 P. Spentzouris, I. Stancu, R. J. Stefanski, M. Sung, H. A. Tanaka,  
 R. Taylor, M. Tzanov, M. O. Wascko, R. Van de Wazer, D. H. White,  
 M. J. Wilking, H. J. Yan, G. P. Zeller, E. D. Zimmerman



University of Alabama  
 Bucknell University  
 University of Cincinnati  
 University of Colorado  
 Columbia University  
 Embry Riddle University  
 Fermi National Accelerator Laboratory  
 Indiana University

Los Alamos National Laboratory  
 Louisiana State University  
 University of Michigan  
 Princeton University  
 Saint Mary's University of Minnesota  
 Virginia Polytechnic Institute  
 Western Illinois University  
 Yale University

## The MiniBooNE Strategy

Test the LSND indication of anti-electron neutrino oscillations  
 keep same  $L/E$ , change beam energy and systematic errors

$$P(\nu_\mu \rightarrow \nu_e) = \sin^2 2\theta \sin^2(1.27 \Delta m^2 L/E)$$

Neutrino Energy (E):


MiniBooNE: ~600 MeV

LSND: ~30 MeV

Baseline (L):

MiniBooNE: ~540 m


LSND: ~30 m



Integrated Fluxes:  $\nu_\mu = 93.8\%$ ,  $\nu_e = 0.5\%$ ,  $\bar{\nu}_\mu = 5.7\%$ ,  $\bar{\nu}_e = 0.08\%$



## The MiniBooNE Detector



541 meters downstream of target

3 meter overburden of dirt

12 meter diameter sphere

Filled with 800 t of pure mineral oil  
( $\text{CH}_2$  -- density  $0.845 \text{ gr/cm}^3$ ,  $n=1.47$ )


Fiducial volume 450 t

1280 inner 8" phototubes – 10% coverage


240 veto phototubes

Less than 2% tubes failed during run

## Neutrino Interactions




MiniBooNE typical  $\nu$  energy





- $5.6 \times 10^{20}$  POT in neutrino mode (10/02-12/05).
  - 193,709  $\nu_\mu$  CCQE interaction candidates
- Phys. Rev. Lett. 98, 231801 (2007).

## $\nu$ events in the detector

- Cosmic  $\mu$  rejected with low veto activity cut.
- Exponential decay:  $e$  from  $\mu$  decay: Rejected with minimum tank hits cut.



Sub-events:



- $\mu$  from  $\nu_{\mu}$  CCQE interactions have typically two sub-events.
- $\nu_e$  CCQE interactions, typically one sub-event.





## Oscillation analysis structure

Two algorithms used:


- (1) Track Based Likelihood (TBL\*)  
 Uses direct reconstruction of particle types and likelihood ratios for PID.
- (2) Boosted Decision Tree (BDT)  
 Uses less detailed reconstruction, and a set of "low level" variables combined in BDT algorithm into a PID score.

The TBL analysis had higher sensitivity to oscillations, hence was chosen for primary results.






## MiniBooNE First Results (April, 2007)



Data consistent with expected background  
 $\Rightarrow$  Inconsistent with a  $\nu_\mu \rightarrow \nu_e$  oscillations

Exclude region in parameter space:



Oscillation Search Region  
 $475 < E_\nu < 1250$  MeV

data:  $380 \pm 19$  (stat) events  
 expectation:  $358 \pm 35$  (sys) events  
 significance:  $0.55 \sigma$

Best Fit (dashed):  
 $(\sin^2 2\theta, \Delta m^2) = (0.001, 4 \text{ eV}^2)$

Probability of Null Fit: 93%  
 Probability of Best Fit: 99%

Published: Phys. Rev. Lett. 98, 231801 (2007)

## Oscillation Signal

⇒ An Excess of " $\nu_e$ " Events over Expectation


All the major backgrounds for the oscillation search can be constrained directly from measurements using MiniBooNE data

- **NC  $\pi^0$  production:** (arXiv:0803.3423, accepted for publication by Phys. Rev. Lett.)  
Largest mis-ID background, where one of the decay photons is missed.  
MiniBooNE cannot distinguish electrons from single gammas.  
Rate constrained from dedicated NC  $\pi^0$  sample. Also constrains radiative  $\Delta$  decays:  
 $\Delta \rightarrow N\gamma$ .
- **External events (Dirt):**  
Backgrounds from interactions with material outside of the detector. Rate constrained from dedicated sample.
- **Intrinsic kaon decay  $\nu_e$ 's:**  
Partially constrained by observed  $\nu_e$  events at high energy where there are no oscillation events.
- **Intrinsic muon decay  $\nu_e$ 's:**  
Largest intrinsic  $\nu_e$  background. Highly constrained by the observed  $\nu_\mu$  events. The constraint can be applied by using the combined  $\nu_e/\nu_\mu$  oscillation fit.



## Measurement of $\nu_e$ NC $\pi^0$ Rate and constraint of $\nu_e$ of Mis-IDs

Largest NC  $\pi^0$  sample ever collected (28,600  $\pi^0$  events)




- $\pi^0$  rate measured to a few percent.
- Critical to oscillation analysis  $\rightarrow$  without  $\pi^0$  rate errors would be  $\sim 25\%$
- First measurement of coherent NC  $\pi^0$  production off  $^{12}\text{C}$  below 2 GeV ( $19.5 \pm 2.7\%$ ).

arXiv: 0803.3423, accepted by Phys. Lett. B



## TBL Analysis: expected events

$\nu_e$  candidate sample composition shown below:



Counting experiment:

475 MeV – 1250 MeV	
$\nu_e^K$	$94 \pm 27$
$\nu_e^\mu$	$132 \pm 10$
NC $\pi^0$	$62 \pm 10$
Dirt	$17 \pm 3$
$\Delta \rightarrow N\gamma$	$20 \pm 4$
Other	$33 \pm 6$
<b>Total</b>	<b><math>358 \pm 35</math></b>
LSND best fit $\nu_\mu \rightarrow \nu_e$	$126 \pm 21$

Counting experiment range

Sig/ $\sqrt{Bkgd}$  = 6.8

$$E_V^{QE} = \frac{1}{2} \frac{2M_p E_\ell - m_\ell^2}{M_p - E_\ell + \sqrt{(E_\ell^2 - m_\ell^2)} \cos\theta_\ell}$$

## A Combined $\nu_e$ BDT, $\nu_e$ TBL, $\nu_\mu$ CCQE Oscillations Fit

Do oscillation fit to the observed and  $\nu_e$  BDT,  $\nu_e$  TBL, and  $\nu_\mu$  CCQE energy distributions by minimizing the following  $\chi^2$  statistic:

$$\chi^2 = \begin{pmatrix} \Delta_i^{\nu_e \text{BDT}} & \Delta_i^{\nu_e \text{TBL}} & \Delta_i^{\nu_\mu \text{CCQE}} \end{pmatrix} \begin{pmatrix} M_{ij}^{\nu_e \text{BDT}, \nu_e \text{BDT}} & M_{ij}^{\nu_e \text{BDT}, \nu_e \text{TBL}} & M_{ij}^{\nu_e \text{BDT}, \nu_\mu \text{CCQE}} \\ M_{ij}^{\nu_e \text{TBL}, \nu_e \text{BDT}} & M_{ij}^{\nu_e \text{TBL}, \nu_e \text{TBL}} & M_{ij}^{\nu_e \text{TBL}, \nu_\mu \text{CCQE}} \\ M_{ij}^{\nu_\mu \text{CCQE}, \nu_e \text{BDT}} & M_{ij}^{\nu_\mu \text{CCQE}, \nu_e \text{TBL}} & M_{ij}^{\nu_\mu \text{CCQE}, \nu_\mu \text{CCQE}} \end{pmatrix}^{-1} \begin{pmatrix} \Delta_j^{\nu_e \text{BDT}} \\ \Delta_j^{\nu_e \text{TBL}} \\ \Delta_j^{\nu_\mu \text{CCQE}} \end{pmatrix}$$

where  $\Delta_i^{\nu_e \text{BDT/TBL}} = \text{Data}_i^{\nu_e \text{BDT/TBL}} - \text{Pred}_i^{\nu_e \text{BDT/TBL}}(\Delta m^2, \sin^2 2\theta)$ , and  $\Delta_j^{\nu_\mu \text{CCQE}} = \text{Data}_j^{\nu_\mu \text{CCQE}} - \text{Pred}_j^{\nu_\mu \text{CCQE}}$

Systematic (and statistical) uncertainties are included in  $(M_{ij})^{-1}$  matrix

- Covariance matrix includes correlations between  $\nu_e$  and  $\nu_\mu$  events.
- Statistical error component takes care of event overlap in  $\nu_e$  samples.
- 68% of TBL  $\nu_e$ 's are also BDT  $\nu_e$ 's  $\Rightarrow$  improvement is expected.

Need to define which  $\nu_\mu$  CCQE sample to use. In this calculation we use the  $\nu_\mu$  CCQE sample of the BDT analysis in the combination. This causes a loss of sensitivity in the TBL component (not identical to first result).



## The $\nu_e$ BDT + $\nu_e$ TBL + $\nu_\mu$ CCQE results:

Paper at draft stage


The combination of the three samples gives a significant increase in coverage in the region  $\Delta m^2 < 1 \text{ eV}^2$ .

Differences in the details are due to the specific fluctuations in the three data samples and the interplay with correlations among them.

$3\sigma$  and  $5\sigma$  limits improve significantly:  $5\sigma$  is comparable to previous  $3\sigma$  at low  $\Delta m^2$ .

The combination yields a consistent result.

Limits from fits to open data



10%-30% improvement in  
90% C.L. limit below  $\sim 1 \text{ eV}^2$ .



## Global data analysis

Combine results of MiniBooNE, LSND, KARMEN2, and Bugey.

### Compatibility:

- How probable is it that all experimental results come from the same underlying 2- $\nu$  oscillation hypothesis?
- Assessed by combining the  $\Delta\chi^2$  surface of each experiment.

LSND	KARMEN2	MB	Bugey	Max. Comp. (%)	$\Delta m^2$ (eV <sup>2</sup> )	$\sin^2 2\theta$
✓	✓	✓		25.36	0.072	0.256
✓	✓	✓	✓	3.94	0.242	0.023
✓		✓		16.00	0.072	0.256
✓		✓	✓	2.14	0.253	0.023
	✓	✓		73.44	0.052	0.147
	✓	✓	✓	27.37	0.221	0.012

arXiv: 0805.1764[hep-ex], submitted to Phys. Rev. D

## Global data analysis

Combine results of MiniBooNE, LSND, KARMEN2, and Bugey.

### Compatibility:

- How probable is it that all experimental results come from the same underlying 2- $\nu$  oscillation hypothesis?
- Assessed by combining the  $\Delta\chi^2$  surface of each experiment.


LSND	KARMEN2	MB	Bugey	Max. Comp. (%)	$\Delta m^2$ (eV <sup>2</sup> )	$\sin^2 2\theta$
✓	✓	✓		25.36	0.072	0.256
✓	✓	✓	✓	3.94	0.242	0.023
✓		✓		16.00	0.072	0.256
✓		✓	✓	2.14	0.253	0.023
	✓	✓		73.44	0.052	0.147
	✓	✓	✓	27.37	0.221	0.012

arXiv: 0805.1764[hep-ex], submitted to Phys. Rev. D

## Global data analysis, Allowed regions

### Allowed Regions:

- Indicate where oscillation parameters would lie, at a given CL, assuming all experimental results can arise in a framework of 2- $\nu$  oscillations.
- The compatibility is the measure of this assumption.



### LSND, KARMEN2 & MiniBooNE:

- 25.36% compatibility at  $\Delta m^2 = 0.072 \text{ eV}^2$ ,  $\sin^2 2\theta = 0.256$ .


### LSND, KARMEN2, MiniBooNE & Bugey:

- 3.94% compatibility at  $\Delta m^2 = 0.242 \text{ eV}^2$ ,  $\sin^2 2\theta = 0.023$ .



## We observe an excess of events below 475 MeV

- $96 \pm 17 \pm 20$  evts. above background for  $300 < E_{\nu}^{QE} < 475$  MeV.
- Opened bin from 200-300 MeV.
- Calculated full systematic errors.
- Excess persists below 300 MeV



	Reconstructed $\nu$ energy bin (MeV)		
	200-300	300-475	475-1250
<b>total BG</b>	<b>284±25</b>	<b>274±21</b>	<b>358±35</b>
$\nu_e$ intrinsic	26	67	229
$\nu_{\mu}$ induced	258	207	129
NC $\pi^0$	115	76	62
NC $\Delta \rightarrow N\gamma$	20	51	20
Dirt	99	50	17
other	24	30	30
<b>DATA</b>	<b>375±19</b>	<b>369±19</b>	<b>380±19</b>

### New Bin


- $\nu_{\mu}$  mis-ID BG dominates the new bin even more.

## Investigating the low E excess ( $E_\nu < 475$ MeV)


### No Reconstruction problems found

All low-E electron candidate events have been examined via event displays, consistent with 1-ring events.

*Could be electrons or photons.*



event/POT vs day,  $300 < E < 475$  MeV



### No Detector anomalies found

Example: rate of electron candidate events is constant (within errors) over course of run

## Possible Sources of Single Gamma Backgrounds


**MiniBooNE cannot tell an electron from a single gamma.**

Processes that remove/absorb one of the gammas from a  $\nu_\mu$  induced NC  $\pi^0 \rightarrow \gamma\gamma$

- Should be in the GEANT detector Monte Carlo. Might be exceptions or inaccurate rates.
  - **Example: photo-nuclear absorption**

⇒ Under active investigation


photo nuclear cross section



$\nu$  processes that produce a final state single gamma

- **Example: Anomaly mediated photon production** (Harvey, Hill, and Hill, arXiv:0708.1281[hep-ex])

⇒ Under active investigation






## Advances in understanding the Low Energy excess:

- Included photo-nuclear effect (reduces excess)
  - Absent from GEANT3 – creates background from  $\pi^0$ 's
- More comprehensive hadronic errors.
  - e.g. uncertainties from final states in photo-nuclear interactions
- Better handling of beam  $\pi^+$  production uncertainties
  - Errors propagated in a model-independent way
- Improved measurement of  $\nu$  induced  $\pi^0$ 's (increases excess)
  - e.g. finer momentum binning
- Incorporation of MiniBooNE  $\pi^0$  coherent/resonant measurement (increases excess)
  - No longer rely on more uncertain past results
- Better handling of radiative decay of  $\Delta$  resonance (reduces excess)
  - As inferred from the measured  $\pi^0$  rate.

Nearing a the end of a comprehensive review of the  $\nu_e$  appearance backgrounds and uncertainties. Not ready for release yet.


## Check with neighboring neutrino source: NuMI $\rightarrow$ MINOS




Test of principle of a horn-focused off-axis beam.

Joint Exp. Theor. Phys. Seminar Dec. 11, 2007

NuMI flux composition:  
 $\nu_\mu$  (66%),  $\nu_e$  (2%),  $\bar{\nu}_\mu$  (31%),  $\bar{\nu}_e$  (1%)



Enhanced in  $\nu_e$  from K decay because of the off-axis position ( $\sim 111$  mrad off axis).





## MiniBooNE Present and Future

- Collected  $\sim 6.6 \times 10^{20}$  POT in neutrino mode.
  - Making various cross section measurements.
  - Searching for various neutrino oscillations.
  - Publications being produced.
- Collected  $\sim 2.5 \times 10^{20}$  POT in anti-neutrino mode.
  - Making various cross section measurements.
  - Searching for  $\bar{\nu}_\mu$  disappearance.
- In Nov 2007 request for extra running in anti-neutrino mode granted.
  - LSND was an indication of  $\bar{\nu}_e$  appearance.
  - Extra  $\sim 2.5 \times 10^{20}$  for a grand total of  $\sim 5 \times 10^{20}$  POT.
  - Will take data during FY2008 and FY2009.



## Summary

- MiniBooNE observes no evidence for  $\nu_{\mu} \rightarrow \nu_e$  2 $\nu$  oscillations.
- Combined BDT and TBL analysis sets tighter limit below  $\Delta m^2 < 1 \text{ eV}^2$ .
- High  $\Delta m^2$  experiments (LSND, KARMEN2, MB & Bugey) compatible only at the 3.94% level.
- Low energy excess under active investigation. Expect full update this summer.
- NuMI beam data is complementary to MiniBooNE flux. Only a small significance excess in the  $\nu_e$  sample is seen with current uncertainties (will constrain them using  $\nu_{\mu}$  sample as done with booster beam data).
- More analysis of more data in progress.