


Title: Thermodynamics of Galaxy Clusters and Beyond

Date: Apr 27, 2009 04:45 PM

URL: <http://pirsa.org/09040037>

Abstract: Clusters of galaxies are unique probes of cosmology and astrophysics, promising to provide new insights into both the nature of dark energy and dark matter and the physics of galaxy formation. One of the key challenges facing this approach lies in our understanding of cluster physics and their impact on cluster structure and evolution. In this talk, I will present numerical simulations of galaxy clusters and their comparisons to recent Chandra X-ray observations, with focus on thermodynamics of intracluster plasma. Numerical simulations including gas cooling and star formation reproduce global properties of the intracluster medium (ICM) and observable-mass relations with an accuracy of ~10%. I will further show that non-thermal processes, such as turbulence, cosmic-rays, and ICM plasma physics, will become important for understanding the remaining systematic uncertainty in the cluster mass estimate and cosmological constraints derived using galaxy clusters.

Thermodynamics of Galaxy Clusters and Beyond



Daisuke Nagai



Yale University

Perimeter Institute

April 27th 2009

Cosmology and Astrophysics with Galaxy Clusters

Clusters of galaxies provide important insights into the nature of dark energy and dark matter.



The most massive galaxies and black holes in the universe form and evolve in cores of galaxy clusters.

Cosmology and Astrophysics with Galaxy Clusters

- The majority of baryons in clusters are in the form of hot, X-ray emitting intracluster plasma.
- Understanding thermodynamics (e.g., heating and cooling) of the intracluster plasma is important for the use of galaxy clusters as cosmological probes as well as understanding the physics of the most massive galaxies and black holes.
- **Main Challenges for Cluster Cosmology:** understanding cluster gas physics (e.g., gas cooling and heating by energy feedback) and calibrate the relationship between X-ray and SZE observables and mass ($\Delta=500$).

$$\begin{aligned} T_{\text{gas}} &\propto GM_{\Delta} / R_{\Delta} \propto M_{\Delta}^{2/3} \\ \text{SZ flux} &\propto \int P_{\text{gas}} dl d\Omega \propto f_{\text{gas}} M_{\Delta}^{5/3} \end{aligned}$$

$$M_{\Delta} \equiv (4\pi/3) R_{\Delta}^3 \Delta \rho_{\text{crit}}(z)$$

How does galaxy formation physics affect global cluster properties?

How do current simulations compare with observations?

How well cluster observables correlate with mass?

High-Resolution Cluster Simulations

Dark Matter



Stars




Box Size $\sim 80 h^{-1} \text{Mpc}$
Peak Resolution $\sim 2 h^{-1} \text{kpc}$


Gas Density




Entropy



Temperature



Metals



N-body+Gasdynamics with ART code


- Collisionless dynamics of DM and stars
- Gasdynamics: Eulerian Adaptive Mesh Refinement
- Radiative cooling and heating of gas: metallicity dependent net cooling/heating rates
- Star Formation using the Kennicutt (1998) recipe
- Thermal stellar feedback
- Metal enrichment by SNIa/Ia
- No AGN feedback, thermal conduction, cosmic-rays, magnetic field & physical viscosity

Cluster Samples

- High-resolution allows us to actually simulate clusters of galaxies
- Effects of galaxy formation on the ICM
 - ▶ Sample of 16 clusters in Λ CDM model
 - ▶ Two sets of runs with cooling & SF (CSF) and with non-radiative gasdynamics
 - ▶ Comparison with Chandra X-ray observations of nearby relaxed clusters


Testing Chandra measurements with mock observations of simulated clusters

- generate “Chandra data” for clusters from cosmological simulations
- reduce with real data analysis pipeline
 - ▶ gas mass accurate to ~3%, temperatures are accurate to <~10%
 - ▶ but, hydrostatic mass is biased low by ~10% due to turbulence




Testing Chandra measurements with mock observations of simulated clusters

- generate “Chandra data” for clusters from cosmological simulations
- reduce with real data analysis pipeline
 - ▶ gas mass accurate to ~3%, temperatures are accurate to <~10%
 - ▶ but, hydrostatic mass is biased low by ~10% due to turbulence



Intracluster Gas Profiles: Effects of gas cooling and star formation





red line:
mean profile for relaxed
clusters in non-radiative
("adiabatic") simulations

blue band:
mean profile for relaxed
clusters in simulations with
cooling and star formation
width = rms scatter

dotted line : $T_x < 3\text{keV}$
dashed line : $T_x > 3\text{keV}$



Nagai, Kravtsov, Vikhlinin
2007, ApJ, 668, 1

Intracluster Gas Profiles: Comparison with observations




red line:
mean profile for relaxed
clusters in non-radiative
("adiabatic") simulations

blue band:
mean profile for relaxed
clusters in simulations with
cooling and star formation
width = rms scatter



Thin dashed lines: profiles
of Chandra observations of
nearby, relaxed clusters of
different temperature

Intracluster Gas Profiles: Effects of gas cooling and star formation





red line:
mean profile for relaxed clusters in non-radiative ("adiabatic") simulations

blue band:
mean profile for relaxed clusters in simulations with cooling and star formation
width = rms scatter

dotted line : $T_x < 3\text{keV}$
dashed line : $T_x > 3\text{keV}$



Nagai, Kravtsov, Vikhlinin
2007, ApJ, 668, 1

Intracluster Gas Profiles: Comparison with observations





red line:
mean profile for relaxed clusters in non-radiative (“adiabatic”) simulations

blue band:
mean profile for relaxed clusters in simulations with cooling and star formation
width = rms scatter



Thin dashed lines: profiles of Chandra observations of nearby, relaxed clusters of different temperature

Entropy scaling with cluster mass & T_x



Simulations with cooling+SF reproduce both the amplitude and scaling with temperature (i.e., mass) exhibited by observed clusters at $r > 0.1 r_{200}$, but not in the core

Solid black pts: CSF simulations

Open black pts: non-radiative sim.

magenta pts: Chandra data
Vikhlinin et al. 2006


red pts: XMM-Newton data
Pratt et al. 2006

blue pts: ROSAT+ASCA data
Ponman et al. 2003

Nagai, Kravtsov, Vikhlinin
2007, ApJ, 668, 1

additional physical processes affect properties of intracluster gas in cores

example: heating by Active Galactic Nuclei of the central cluster galaxy in the Perseus cluster (Talks by A. Babul, P. Bode, & P. Oh)




these effects, however, appear to be confined to the core

=> outer regions of clusters can be used to reliably

estimate their total masses

Mass – ICM temperature relation



~10% agreement in the amplitude between observed and model M-Tx relation -- improvements are in both sim. and obs.

Scatter in M-Tx is ~20% in mass at a given Tx - the scatter is primarily driven by unrelaxed systems


Unrelaxed systems have systematically lower Tx

black pts: Simulated clusters with cooling+SF

magenta pts: Chandra data Vikhlinin et al. 2006 ApJ, 640, 691

additional physical processes affect properties of intracluster gas in cores

example: heating by Active Galactic Nuclei of the central cluster galaxy in the Perseus cluster (Talks by A. Babul, P. Bode, & P. Oh)





these effects, however, appear to be confined to the core

=> outer regions of clusters can be used to reliably



estimate their total masses

Intracluster Gas Profiles: Comparison with observations



red line:
mean profile for relaxed clusters in non-radiative (“adiabatic”) simulations


blue band:
mean profile for relaxed clusters in simulations with cooling and star formation
width = rms scatter




Thin dashed lines: profiles of Chandra observations of nearby, relaxed clusters of different temperature

Testing Chandra measurements with mock observations of simulated clusters

- generate “Chandra data” for clusters from cosmological simulations
- reduce with real data analysis pipeline
 - ▶ gas mass accurate to ~3%, temperatures are accurate to <~10%
 - ▶ but, hydrostatic mass is biased low by ~10% due to turbulence



Intracluster Gas Profiles: Effects of gas cooling and star formation




red line:
mean profile for relaxed
clusters in non-radiative
("adiabatic") simulations

blue band:
mean profile for relaxed
clusters in simulations with
cooling and star formation
width = rms scatter


dotted line : $T_x < 3\text{keV}$
dashed line : $T_x > 3\text{keV}$

Nagai, Kravtsov, Vikhlinin
2007, ApJ, 668, 1

Mass – Y_x relation a new X-ray mass proxy



Mass – ICM temperature relation



~10% agreement in the amplitude between observed and model M - T_x relation -- improvements are in both sim. and obs.


Scatter in M - T_x is ~20% in mass at a given T_x - the scatter is primarily driven by unrelaxed systems

Unrelaxed systems have systematically lower T_x

black pts: Simulated clusters with cooling+SF


magenta pts: Chandra data Vikhlinin et al. 2006 ApJ, 640, 691

Mass – Y_x relation a new X-ray mass proxy



X-ray "pressure" = gas mass x temperature

Mass – ICM temperature relation



~10% agreement in the amplitude between observed and model M - T_x relation -- improvements are in both sim. and obs.


Scatter in M - T_x is ~20% in mass at a given T_x - the scatter is primarily driven by unrelaxed systems

Unrelaxed systems have systematically lower T_x

black pts: Simulated clusters with cooling+SF

magenta pts: Chandra data Vikhlinin et al. 2006 ApJ, 640, 691

Mass – ICM temperature relation



~10% agreement in the amplitude between observed and model M - T_x relation -- improvements are in both sim. and obs.


Scatter in M - T_x is ~20% in mass at a given T_x - the scatter is primarily driven by unrelaxed systems

Unrelaxed systems have systematically lower T_x

black pts: Simulated clusters with cooling+SF


magenta pts: Chandra data Vikhlinin et al. 2006 ApJ, 640, 691

Mass – Y_x relation a new X-ray mass proxy




X-ray "pressure" = gas mass x temperature

Mass – Y_x relation comparison with observations




X-ray "pressure" = gas mass x temperature

Mass – Y_x relation a new X-ray mass proxy




X-ray "pressure" = gas mass x temperature

Mass – Y_x relation comparison with observations




X-ray "pressure" = gas mass x temperature

Accretion, Mergers → Shocks, Turbulence




Norman & Bryan 1999, Nagai, Kravtsov & Kosowsky 2003
Sunyaev, Norman & Bryan 2003; Rasia et al. 2004, 2006;
Delag et al. 2005; Nagai et al. 2007; Lau et al. 2009

Mass – Y_x relation comparison with observations




Accretion, Mergers → Shocks, Turbulence



Norman & Bryan 1999, Nagai, Kravtsov & Kosowsky 2003
Sunyaev, Norman & Bryan 2003; Rasia et al. 2004, 2006;
Delag et al. 2005; Nagai et al. 2007; Lau et al. 2009


Effect of random gas motions on mass measurements

$$M_{\text{tot}}(< r) = \frac{-r^2}{G\rho} \left(\frac{dP_{\text{ther}}}{dr} + \frac{dP_{\text{turb}}}{dr} \right)$$



Effect of random gas motions on mass measurements


$$M_{\text{tot}}(< r) = \frac{-r^2}{G\rho} \left(\frac{dP_{\text{ther}}}{dr} + \frac{dP_{\text{turb}}}{dr} \right)$$



Lau, Kravtsov, Nagai 2009
 astro-ph/0903.4895



Mass – Y_x relation

using mass derived from the hydrostatic equilibrium analysis
both in observations and simulations



Dark energy constraints from the evolution of cluster mass function

The 400 sq. deg. X-ray cluster survey (Vikhlinin et al. 2009)
 Also talk by M. Hilton for the XMM cluster survey



M_{500} – total mass within radius enclosing
 overdensity of $500 \times \rho_{crit}$



Dark energy density, Ω_x

Cosmology with Sunyaev-Zel'dovich Effect

Upcoming SZE cluster surveys will produce large statistical samples
(e.g., from AMI, AMiBA, APEX, SZA to ACT, Planck, and SPT)

SZ Effect directly probes the
integrated pressure

Sample of 38 BIMA/OVRO SZE+Chandra
X-ray clusters at $0.14 < z < 0.89$





Gas mass within r_{2500}

Simulations: Nagai 2006, also Mott et al. 2005, Hallman et al. 2007

Data: Bonamente, Joy, LaRoque, Carlstrom, Nagai, Marrone 2008

Dark energy constraints from the evolution of cluster mass function

The 400 sq. deg. X-ray cluster survey (Vikhlinin et al. 2009)
 Also talk by M. Hilton for the XMM cluster survey



M_{500} – total mass within radius enclosing
 overdensity of $500 \times \rho_{\text{crit}}$



Dark energy density, Ω_x

Cosmology with Sunyaev-Zel'dovich Effect

Upcoming SZE cluster surveys will produce large statistical samples
(e.g., from AMI, AMiBA, APEX, SZA to ACT, Planck, and SPT)

SZ Effect directly probes the
integrated pressure

Sample of 38 BIMA/OVRO SZE+Chandra
X-ray clusters at $0.14 < z < 0.89$





Gas mass within r_{2500}

Simulations: Nagai 2006, also Mott et al. 2005, Hallman et al. 2007

Data: Bonamente, Joy, LaRoque, Carlstrom, Nagai, Marrone 2008

Dark energy constraints from the evolution of cluster mass function

The 400 sq. deg. X-ray cluster survey (Vikhlinin et al. 2009)
 Also talk by M. Hilton for the XMM cluster survey



M_{500} – total mass within radius enclosing
 overdensity of $500 \times \rho_{\text{crit}}$



Dark energy density, Ω_x

Cosmology with Sunyaev-Zel'dovich Effect

Upcoming SZE cluster surveys will produce large statistical samples
(e.g., from AMI, AMiBA, APEX, SZA to ACT, Planck, and SPT)

SZ Effect directly probes the
integrated pressure

Sample of 38 BIMA/OVRO SZE+Chandra
X-ray clusters at $0.14 < z < 0.89$



Gas mass within r2500


Simulations: Nagai 2006, also Mott et al. 2005, Hallman et al. 2007

Data: Bonamente, Joy, LaRoque, Carlstrom, Nagai, Marrone 2008

Probing cosmic-rays pressure with Fermi

$$M_{\text{tot}}(< r) = \frac{-r^2}{G\rho} \left(\frac{dP_{\text{ther}}}{dr} + \frac{dP_{\text{turb}}}{dr} + \frac{dP_{\text{cr}}}{dr} \right)$$

Fermi will provide stringent constraints (~1%) on the cosmic-ray protons in nearby, rich clusters





Ando & Nagai 2008; also

Cosmology with Sunyaev-Zel'dovich Effect

Upcoming SZE cluster surveys will produce large statistical samples
(e.g., from AMI, AMiBA, APEX, SZA to ACT, Planck, and SPT)

SZ Effect directly probes the
integrated pressure

Sample of 38 BIMA/OVRO SZE+Chandra
X-ray clusters at $0.14 < z < 0.89$



Gas mass within r_{2500}



Simulations: Nagai 2006, also Mott et al. 2005, Hallman et al. 2007

Data: Bonamente, Joy, LaRoque, Carlstrom, Nagai, Marrone 2008

Probing cosmic-rays pressure with Fermi

$$M_{\text{tot}}(< r) = \frac{-r^2}{G\rho} \left(\frac{dP_{\text{ther}}}{dr} + \frac{dP_{\text{turb}}}{dr} + \frac{dP_{\text{cr}}}{dr} \right)$$

Fermi will provide stringent constraints (~1%) on the cosmic-ray protons in nearby, rich clusters





Ando & Nagai 2008; also

Cosmology with Sunyaev-Zel'dovich Effect

Upcoming SZE cluster surveys will produce large statistical samples
(e.g., from AMI, AMiBA, APEX, SZA to ACT, Planck, and SPT)

SZ Effect directly probes the
integrated pressure

Sample of 38 BIMA/OVRO SZE+Chandra
X-ray clusters at $0.14 < z < 0.89$



Gas mass within r_{2500}



Simulations: Nagai 2006, also Mott et al. 2005, Hallman et al. 2007

Data: Bonamente, Joy, LaRoque, Carlstrom, Nagai, Marrone 2008

Probing cosmic-rays pressure with Fermi

$$M_{\text{tot}}(< r) = \frac{-r^2}{G\rho} \left(\frac{dP_{\text{ther}}}{dr} + \frac{dP_{\text{turb}}}{dr} + \frac{dP_{\text{cr}}}{dr} \right)$$

Fermi will provide stringent constraints (~1%) on the cosmic-ray protons in nearby, rich clusters




Ando & Nagai 2008; also


He sedimentation in X-ray Clusters

$$M_{\text{tot}}(< r) = \frac{-r^2}{G\rho} \frac{dP_{\text{ther}}}{dr} \propto \frac{1}{\mu} = \frac{8 + 3(Y/X)}{4 + 4(Y/X)}$$

Intracluster plasma consists of ~75% hydrogen and ~25% helium by mass




Solving the diffusion equations for the fully ionized H-He plasma in the NFW potential



He sedimentation can introduce systematic uncertainty in X-ray measurements of galaxy clusters at the level of <5-10%.

Bias in X-ray + SZE derived Hubble parameter

H_0 measurements at $r = r_{2500}$



Enhancement in He-to-H number density ratio relative to the primordial value

X-ray + SZE-derived Hubble parameter

$$\begin{cases} S_X \propto n_e^2 d_A \left(\frac{1+4x}{1+2x} \right) \\ y_{SZE} \propto n_e T_e d_A \end{cases} \Rightarrow d_A \propto \frac{y_{SZE}^2}{S_X T_e^2} \left(\frac{1+4x}{1+2x} \right)$$

Our model predicts an overestimate of the X-ray+SZE H_0 ($\propto d_A^{-1}$) by about 5%.

This can explain the offset seen in the latest X-ray+SZE-derived $H_0 = 77.6 \pm 4.5$ $\text{km s}^{-1} \text{Mpc}^{-1}$ (Bonamante et al. 2007) relative to $H_0 = 73$ $\text{km s}^{-1} \text{Mpc}^{-1}$ of the WMAP and/or Hubble Key projects.

Summary

■ Successes

- ▶ Modern cosmological cluster simulations with cooling+SF reproduce observed thermodynamic properties of real clusters outside cores.
- ▶ Observable-mass relations of simulated clusters and recent X-ray observations agree to about 10%.
- ▶ Robust, low-scatter mass proxies (Y_x and Y_{sz}) are accessible for *both* X-ray and SZE cluster surveys.

■ Problems & Challenges

- ▶ *But, there is a remaining offset of ~10% between simulations and observations. Likely, due to non-thermal pressure components (e.g., turbulence, cosmic-rays, ICM plasma physics).*
- ▶ *Also, cluster cores are not well-reproduced in simulations.*

■ Future Prospect

- ▶ Upcoming cluster surveys will produce large statistical samples of clusters (X-ray: eROSITA; SZE: ACT, AMI, APEX, Planck, SPT, SZA)
- ▶ Further advances in numerical simulations are also underway
 - ❖ Larger sample of simulated clusters to study the scatter
 - ❖ Detailed understanding of cluster gas physics (e.g., AGN feedback, turbulence, cosmic-rays, ICM plasma physics)