Title: Ekpyrotic/Cyclic Cosmology - Lecture 4

Date: Jun 29, 2009 11:30 AM

URL: http://pirsa.org/09060081

Abstract: TBA


IV Higher-Dimensional Point of View


IV Higher-Dimensional Point of View
Approach to Big crunch / Sol Milne
after ek phase V turns off
Finale phase $a = a_0(-t)^{1/3}$


Approach to Big crunch / 5d Milne
after ek phase
V turns off
kinetic phase
$a = a_0 (-t)^{1/3}$
gravity in 5d
$S_5 = \frac{1}{2} \int \sqrt{-g_5} R_5$


Approach to Big crunch / 5d Milne
after ek phase V turns off
linear phase $a = a_0(-t)^{1/3}$
$S_5 = \frac{1}{2}\int \sqrt{-g}\, R_5$
orbifold fixed pts


compactify on $S^1/\mathbb{Z}_2$
orbifold fixed pts
y
$ds_5^2 = e^{-\frac{1}{\sqrt{3}}\phi} ds_4^2 + e^{2\sqrt{3}\phi} dy^2$


Compactify on $S^1/\mathbb{Z}_2$
orbifold fixed
$\mathbb{Z}_2$
y
$ds_5^2 = e^{-\frac{1}{\sqrt{3}}\phi} ds_4^2 + e^{2/\sqrt{3}\phi} dy^2$
$\rightarrow S_4 = \frac{1}{2}\int \sqrt{-g}[R - (\partial\phi)^2]$
$ds_4^2 = -dt^2 + a^2(t)\, ds^2$
$6(1-\epsilon)^2$
$ds_5^2 = (1-\epsilon)^{-1/3}[-dt^2 + (1-\epsilon)^{1/3} ds^2] + A(1-\epsilon)^{1/3} dy^2$


$T_f \propto (-\bar{t})^{2/3}$
$ds_5^2 = -dT^2 + T^2 dy^2 + ds^2$
Milne
↳ Fiscale factors in 5d finite!


$T_1 \propto (-\bar{t})^{2/3}$
$ds_5^2 = -dT^2 + T^2 dy^2 + ds^2$
Milne
↳ Physical region is $T \neq 0$!
$\begin{cases} u = T \cosh y \\ v = T \sinh y \end{cases} \Rightarrow ds_1^2 = -du^2 + dv^2 + ds^2$
Minkowski

$T_1 \propto (-\bar{t})^{2/3}$

$ds_5^2 = -dT^2 + T^2 dy^2 + ds^2$

Milne

↳ fiscale factors is for finite!

$\begin{cases} u = T \cosh y \\ v = T \sinh y \end{cases} \Rightarrow ds_1^2 = -du^2 + dv^2 + ds^2$ Minkowski




Horava; Witten
IIA SUGRA ← E8 × E8 heterotic string


Horava; Witten
11d SUGRA
6 M10 x S1/Z2


Horčava; Witten
11d SUGRA
6 M10 x S1/2
10d
10d


Horizon; Whitten
11d SUORA ⟶ $E_1' < E_2$ luotettavuus
6 M_{10} × $5^9/_{22}$
y
F_1
10d
F_2
12d


Horčava; Witten
11d SUGRA → $E_8' \times E_8$ holonomy
$M_{10} \times S^1/\mathbb{Z}_2$
y
E_8
10d
E_8
10d
$A_{11} = g^{2/3}$




$M_{10} \times S^1/\mathbb{Z}_2 \to M_4 \times CY_3 \times S^1/\mathbb{Z}_2$
CY_3
CY_3
M_4
M_4
$d_{11} \sim 30$ Planck


$M_{10} \times S^1/\mathbb{Z}_2 \to M_4 \times CY_6 \times S^1/\mathbb{Z}_2$
CY
CY
4d
4d
$d_H \sim 30$ Hubbles CY
4d → 5d → 11d


4d
4d
$d_{11} \sim 30$ Hubble cy
$4d \rightarrow (5d) \rightarrow 11d$
$t \rightarrow 0$ Branes collide


$ds_{11}^2 = V_{CY}^{-2/3} ds_5^2 + V_{CY}^{1/3} ds_{CY}^2$
in 6d get 2 scalars: volume of CY
radion (distance between orbifolds)


Potential M-theory
$S_F \sim \int \left(R - \frac{(\partial V_{CY}^4)}{V_{CY}^4} - \frac{G\alpha'}{V_{CY}^4} \right)$


Heterotic M-theory
$S_5 \sim \int \left(R - \frac{(\partial V_{CY}^2)}{V_{CY}^2} - \frac{G_4^2}{V_{CY}^2} \right)$
4-potential
$ds_5^2 = h^{2/3}(y)\left[A^2(-dt^2 + dx_1^2 + dx_2^2) + B^2 dy^2\right]$
$V_{CY} = B h^{4/3}$
$h = 5Ay + C$

Intrinsic M-theory

$$S_F \sim \int \left(R - \frac{(\partial V_{cy}^2)}{V_{cy}^2} - \frac{6\alpha'}{V_{cy}^4} \right)$$

hypersurface

Sol

$$ds_5^2 = h^{2/5}(y)\left[A^2(-dt^2 + dz^2) + B^2 dy^2\right]$$

$$V_{cy} = B h^{4/5}$$

$$h = 5\alpha y + C$$


primate A, B, C — fold of C


11d SUGRA ⟷ $E_8 \times E_8$ heterotic string
$M_{10} \times S^1/\mathbb{Z}_2$
-1
$y+1$
E_8
10d
E_8
10d
$M_{11} = g_s^{2/3}$


promote A, B, C — fcts of τ
$I_n \equiv \int_{-1}^{+1} h^n \, dy$
$a^2 = A^2 B I_{1/2}$
$e^{\Phi/\sqrt{2}} = B (I_{1/2})^{3/2}$
$\Phi_2 = -\frac{\sqrt{2}}{20} \int d\zeta \, \frac{1}{I_{1/2}} \sqrt{9(I_{-1/2})^2 + 16 I_{-3/2} I_{1/2}}$

promote A, B, C — fcts of τ

$$I_n \equiv \int_{-1}^{+1} h^n \, dy$$

$$\begin{cases} a^2 = A^2 B I_{1/2} \\ e^{\phi/\sqrt{2}} = B (I_{1/2})^{3/2} \\ \phi_2 = -\frac{\sqrt{2}}{20} \int d\zeta \, \frac{1}{I_{1/2}} \sqrt{9 (I \cdot \dot{u}_4)^2 + 16 I_{-3/5} I_{1/2}} \end{cases}$$

$$\Rightarrow \boxed{S_{11} = \int -6a'' + a^2 (\phi_1'^2 + \phi_2'^2)}$$

promote $A, B, C \rightarrow$ fcts of τ

$$I_n \equiv \int_{-1}^{+1} h^n \, dy$$

$$\begin{cases} a^2 = A^2 B I_{1/2} \\ e^{\phi/\sqrt{2}} = B (I_{1/2})^{3/2} \\ \phi_2 = -\frac{\sqrt{2}}{20} \int d\zeta \, \frac{1}{I_{1/2}} \sqrt{9 (I_{-1/2})^2 + 16 I_{-3/2} I_{1/2}} \end{cases}$$

$$\Rightarrow \boxed{S_{\text{eff}} = \int -6 a a'' + a^2 (\phi_1'^2 + \phi_2'^2)}$$


Horava; Witten
at y = -1 $V_{cy-} = e^{\Phi/4} (-\sinh \sqrt{8}\,\phi_2)^{3/2}$


Witten
at $y=-1$ $V_{cy-} = e^{\frac{\phi_1}{\hbar}} (-\sinh \sqrt{\hbar}\phi_2)^{\chi}$
$\phi_2=0 \Rightarrow V_{cy-}=0$
$\phi_2>0$ unphysical
ϕ_2
forbidden
ϕ_1


promote A, B, C → fcts of τ
$I_n \equiv \int_{-1}^{+1} h^n \, dy$
$\begin{cases} a^2 = A^2 B I_{1/2} \\ e^{\phi/\sqrt{12}} = B (I_{1/2})^{3/2} \\ \phi_2 = -\frac{\sqrt{12}}{20} \int d\tau \frac{1}{I_{1/2}} \sqrt{9 (I_{-1/2})^2 + 16 I_{-3/2} I_{1/2}} \end{cases}$
moduli space approximation
$S_{11} = \int -6a'^2 + a^2 (\phi_1'^2 + \phi_2'^2)$


at y=-1
$\phi_2 = 0 \Rightarrow V_{cy-} = 0$
$\phi_2 > 0$ unphysical
ϕ_2
ϕ_1
$a = |2c|^{1/2}$
$e^{\phi_1} = |2c|^{\frac{3}{2\sqrt{2}}}$
$e^{\phi_1} = |4\alpha c|^{\frac{\sqrt{3}}{2\sqrt{\alpha}}}$


at y=-1
$\phi_1 = 0 \Rightarrow V_{cy-} = 0$
$\phi_1 > 0$ unphysical
ϕ_2
ϕ_1
sol to MSA:
$d \sim 2\sigma$


wide
variety of matter on - ve tension


wide
variety of matter on - ve tension


wide
variety of matter on - ve tension


wide
variety of matter on - ve tension
~ T matter + Very dyn. corrections


-ve tension branes bounce!