Title: Locally inertial coordinates for self-force calculations

Date: Jun 21, 2010 09:10 AM

URL: http://pirsa.org/10060043

Abstract: TBA


R ~ length or Timescale of background
Einstein Tensor
G(g+h) = G(g) +
0
TT+L coords
Locally inert. ∅
Cartesian
Harmonic
G'(g, h) = □h + R...h
G'(g, h) = h∇∇h + ∇h × h
+ L - L
G'(g, h^R) = -G^2(g, h^R) + S_eff


R ~ length or Timescale of background
THz coords
Locally inertial Cartesian Harmonic
Einstein Tensor
G(g+h) =
G'(g,h) = □h + R..h
G²(g,h) = h∇∇h + ∇h×h
+ L-L
G³(g,h) = Lmkn


R ~ length or Timescale of background
THZ coords
Locally inertial Cartesian Harmonic
Einstein Tensor
G(g+h) =
G'(g,h) = □h + R...h
G²(g,h) = h∇∇
G³(g,h) =
G'(g,h^R) = -G²(g,h^R) + S_eff
G/r + O(1/r²)


R ~ length or Timescale of background
THZ coords
Locally inert. ∅
Cartesian
Harmonic
Einstein Tensor
G(g+h) =
G'(g,h) = □h + R..h
+L-L
= unkn
O(h¹) O(h²) O(h³)
G'(g) + G'(g,h) + G²(g,h) + G³(g,h) + ... = 16πT
0
G'(g,h^R + h^S) = 16πT
G'(g,h^R) + G''(g,h^S) = 16πT
G'(g,h^R) = 16πT − G'(h^S) = S_eff
~ 4π/R⁴
h^R = C² at M
C^∞ elsewhere


R ~ length or Timescale of background
THz coords
Locally inertial
Cartesian
Harmonic
Einstein Tensor
G(g+h) =
G¹(g,h) = □h + R...h
G²(g,h) = h∇∇h
G³(g,h) =
G¹(g,h^R) = −G²(g,h^R) + S_eff


Einstein Tensor
$G(g+h) = \underbrace{G^0(g)}_{0} + G^1(g,h) + G^2(g,h) + G^3(g,h) + \ldots = 16\pi T$
$G^1(g,h) = \Box h + \partial R \ldots h$
$G^2(g,h) = h\nabla\nabla h + \nabla h \times h + L - L\Gamma_{ab}$
$G^3(g,h) = \text{cubic}$
$G^1(g,h^R + h^S) = 16\pi T$
$G^1(g,h^R) + G^1(g,h^S) = 16\pi T$
$G^1(g,h^R) = 16\pi T - G^1(h^S) = S_{eff}$
$\sim \frac{m}{r^3}$
$h^R = C^2$ at m
C^∞ elsewhere


Einstein Tensor
$G(g+h) = \underbrace{G^0(g)}_{0} + G^1(g,h) + G^2(g,h) + G^3(g,h) + \cdots = 16\pi T$
$O(h^1)$ $O(h^2)$ $O(h^3)$
$G^1(g,h) = \Box h + \partial R \ldots h$
$G^2(g,h) = h \ldots \nabla h \times h$
$G^3(g,h) = \ldots$
$G^1(g,h^R) = 16\pi T$
$G^1(g,h^R) + G^1(g,h^S) = 16\pi T$
$G^1(g,h^R) = 16\pi T - G^1(h^S) = S_{eff}$
$\sim \frac{M}{r^3}$
$h^R = C^2$ at m
C^∞ elsewhere


Einstein Tensor
$G(g+h) = G^0(g) + G^1(g,h) + G^2(g,h) + G^3(g,h) + \ldots = 16\pi T$
$O(h^1)$ $O(h^2)$ $O(h^3)$
$G^1(g,h) = \Box h + \partial R \ldots h$
$G^2(g,h) = h\nabla\nabla h + \nabla h \times h + \Gamma - \Gamma\Gamma_{av}$
$G^3(g,h) = \ldots$
$G^1(g, h^R + h^S) = 16\pi T$
$G^1(g, h^R) + G^1(h^S) = S_{eff}$
h^R
$\sim \frac{Mr}{R^4}$


Einstein Tensor
$G(g+h) = \underbrace{G^0(g)}_{0} + G^1(g,h) + G^2(g,h) + G^3(g,h) + \cdots = 16\pi T$
$O(h^1)$ $O(h^2)$ $O(h^3)$
$G^1(g,h) = \Box h + \partial R \ldots h$
$G^2(g, h) = h\nabla\nabla h + \nabla h \times h + L\text{-}L\,\Gamma_{\alpha\beta}$
$G^3(g, h) = \text{unkn}$
$G^1(g, h^R + h^S) = 16\pi T$
$G^1(g, h^R) + G^{\geq 1}(g, h^S) = 16\pi T$
$G^1(g, h^R) = 16\pi T - G^1(h^S) = S_{eff}$
$\sim \frac{Mr}{R^9}$
$h^R = C^2$ at m
C^∞ elsewhere

Einstein Tensor

$G(g+h) = \underbrace{G^0(g)}_{0} + G^1(g,h) + G^2(g,h) + G^3(g,h) + \ldots = 16\pi T$

$O(h^1)$ $O(h^2)$ $O(h^3)$

$G^1(g,h) = \Box h + \partial R \ldots h$

$G^2(g,h) = h\nabla\nabla h + \nabla h \times h + \Gamma - \Gamma \Gamma_{\alpha\beta}$

$G^3(g,h) = \text{unkn}$

$G^1(g, h^R + h^S) = 16\pi T$

$G^1(g, h^R) + G^1(g, h^S) = 16\pi T$

$G^1(g, h^R) = 16\pi T - G^1(h^S) = S_{\text{eff}} \sim \frac{m r}{r^3}$

$h^R = C^2$ at m

C^∞ elsewhere


Einstein tensor
$G(g+h) = G^1(g) + G^1(g,h) + G^2(g,h) + G^3(g,h) + \ldots = 16\pi T$
$O(h^1)$ $O(h^2)$ $O(h^3)$
$G^1(g,h) = \Box h + R \ldots h$
$G^2(g,h) = h\nabla\nabla h + \nabla h \times h + L\text{-}C\,\Gamma_{ab}$
$G^3(g,h) = \ldots$
$G^1(g,h^R+h^S)$
$16\pi T$
$G^1(g,h^R) + G^1(g,h^S) = 16\pi T$
$G^1(g,h^R) = 16\pi$
$h^R = C^2$ at
C^∞ el
S_{eff}
$\frac{Mr}{R^3}$


Einstein Tensor
$G(g+h) = G^1(g) + G^1(g,h) + G^2(g,h) + G^3(g,h) + \ldots = 16\pi T$
$O(h^1)$ $O(h^2)$ $O(h^3)$
$G^1(g,h) = \Box h + \partial R \ldots h$
$G^2(g,h) = h\nabla\nabla h + \nabla h \times h + L\text{-}L\Gamma_{ab}$
$G^3(g,h) = \text{cubic}$
$G^1(g,h^R + h^S) = 16\pi T$
$G^1(g,h^R) + G^1(g,h^S) = 16\pi T$
$G^1(g,h^R) = 16\pi T - G^1(h^S) =$
$h^R = C^2$ at Γ
C^∞ elsewhere

Einstein Tensor

$G(g+h) = \underbrace{G^0(g)}_{0} + G^1(g,h) + G^2(g,h) + G^3(g,h) + \dots = 16\pi T$

$O(h^1)$ $O(h^2)$ $O(h^3)$

$G^1(g,h) = \Box h + \partial R \dots h$

$G^2(g,h) = h\nabla\nabla h + \nabla h \times h + L - L\Gamma_{ab}$

$G^3(g,h) = \text{cubic}$

$G^1(g, h^R + h^S) = 16\pi T$

$G^1(g,h^R) + G^1(g,h^S) = 16\pi T$

$G^1(g,h^R) = 16\pi T - G^1(h^S) = S_{eff} \sim \frac{Mr}{R^9}$

$h^R = C^2$ at m

C^∞ elsewhere


Einstein Tensor
$G(g+h) = \underbrace{G^0(g)}_{0} + G^1(g,h) + G^2(g,h) + G^3(g,h) + \ldots = 16\pi T$
$G^1(g,h) = \Box h + R \ldots h$
$G^2(g,h) = h\nabla\nabla h + \nabla h \times h + L - L\Gamma_{ab}$
$G^3(g,h) = \ldots$
$G^1(g,h^R + h^S) = 16\pi T$
$G^1(g,h^R) + G^1(g,h^S) = 16\pi T$
$G^1(g,h^R) = 16\pi T - G^1(g,h^S) = S_{eff}$
$h^R = \ldots$

Einstein Tensor

$G(g+h) = \underbrace{G^0(g)}_{0} + G^1(g,h) + G^2(g,h) + G^3(g,h) + \dots = 16\pi T$

$O(h^1)$ $O(h^2)$ $O(h^3)$

$G^1(g,h) = \Box h + R \dots h$

$G^2(g,h) = h\nabla\nabla h + \nabla h \times h + L - L\,\Gamma_{ab}$

$G^3(g,h) = \text{cubic}$

$G^1(g, h^R + h^S) = 16\pi T$

$G^1(g,h^R) + G^1(g,h^S) = 16\pi T$

$G^1(g,h^R) = 16\pi T - G^1(h^S) = S_{\text{eff}}$

$h^R = C^2$ at m

C^∞ elsewhere

Einstein Tensor

$$G(g+h) = \underbrace{G^0(g)}_{0} + G^1(g,h) + G^2(g,h) + G^3(g,h) + \cdots = 16\pi T$$

$O(h^1)$ $O(h^2)$ $O(h^3)$

$$G^1(g,h) = \Box h + R \cdots h$$

$$G^2(g,h) = h\nabla\nabla h + \nabla h \times h + L - L\Gamma_{ab}$$

$$G^3(g,h) = \text{unkn.}$$

$$G^1(g, h^R + h^S) = 16\pi T$$

$$G^1(g, h^R) + G^1(g, h^S) = 16\pi T$$

$$G^1(g, h^R) = 16\pi T - G^1(h^S) = S_{eff}$$

$$\sim \frac{Mr}{R^4}$$

h^R = C^2 at m

C^∞ elsewhere


From Grains of Pollen to Evidence for Atoms
$h^S = h^{S_1} + h^{S_2}$
$G(g) + G^1(g, h^R) + G^2(g,$
$G^1(g, h^R) = - G^2(g, h^R) -$
$G^1(g, h^R) = - G^2(g, h^R) - [G^1($
$G^1(g, h^R) = - G^2(g, h^R)$
$+ G^2(g, h^S) = 16\pi T$
$+ G^2(g, h^S) - 16\pi T]$
$+ G^1(g + h^R, h^{S_2}) + G^2(g, h^{S_1})$
$O(M^2/R^4)$
discontinuous


From Grains of Pollen to Evidence for Atoms
$h^S = h^{S_1} + h^{S_2}$
$G(g + h^R + h^S) = 16\pi T$
$G(g) + G^1(g, h^R) + G^2(g, h^R) + G^1(g + h^R, h^S) + G^2(g, h^S) = 16\pi T$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^S) + G^2(g, h^S) - 16\pi T]$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^{S_1}) - 16\pi T + G^1(g + h^R, h^{S_2}) + G^2(g, h^{S_1})]$
$G^1(g, h^R) = -G^2(g, h^R) + S_{eff}$
$\mathcal{O}\left(\frac{Mr}{R^4}\right)$
C^0
$\mathcal{O}(M^2/R^4)$
discontinuous


$h^S = h^{S_1} + h^{S_2}$
$G(g_{\mu\nu} + h^R + h^S) = 16\pi T$
$G(g) + G^1(g, h^R) + G^2(g, h^R) + G^1(g + h^R, h^S) + G^2(g, h^S) = 16\pi T$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^S) + G^2(g, h^S) - 16\pi T]$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^{S_1}) - 16\pi T + G^1(g + h^R, h^{S_2}) + G^2(g, h^{S_1})]$
$G^1(g, h^R) = -G^2(g, h^R) + S_{eff}$
$\mathcal{O}\left(\frac{M^4}{R^4}\right)$
C^0
$\mathcal{O}(M^2/R^4)$
discontinuous


$h^S = h^{S1} + h^{S2}$
$= 16\pi T$
$G^1(g, h^R) + G^2(g, h^S) = 16\pi T$
$G^1(g, h^R) + G^2(g, h^S) - 16\pi T$
$G^1(g+h^R, h^{S2}) + G^2(g, h^{S1})$
$O(M^2/R^4)$
discontinuous


$h^S = h^{S1} + h^{S2}$
$= 16\pi T$
$G^1(g, h^R) + G^2(g, h^R)$
$+ G^2(g, h^S) = 16\pi T$
$G^1(g, h^R) = -G^2(g, h^R) - [$
$+ G^2(g, h^S) - 16\pi T]$
$+ G^1(g+h^R, h^{S2}) + G^2(g, h^{S1})$
$O(M^2/R^4)$
discontinuous
$G^1(g, h^R) = -G^2(g, h^R)$


$h^S = h^{S_1} + h^{S_2}$
$G(g+h^R+h^S) = 16\pi T$
$G(g) + G^1(g,h^R) + G^2(g,h^R) + G^1(g+h^R,h^S)$
$= 16\pi T$
$G^1(g,h^R) = -G^2(g,h^R) - [$
$- 16\pi T]$
$G^1(g,h^R) = -G^2(g,h^R) - [G^1(g+h^R,h^{S_1}) - 16$
$+ G^2(g,h^{S_1})$
$G^1(g,h^R) = -G^2(g,h^R) + S_{eff}$


$h^S = h^{S_1} + h^{S_2}$
$G(g + h^R + h^S) = 16\pi T$
$G(g) + G^1(g, h^R) + G^2(g, h^R) + G^1(g + h^R, h^S) + G^2(g, h^S) = 16\pi$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^S) + G^2(g, h^S) - 16$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^{S_1}) - 16\pi T + G^1(g + h^R, h$
$G^1(g, h^R) = -G^2(g, h^R) + S_{eff}$
$\mathcal{O}\left(\frac{M^2}{R^4}\right)$
C^0
$\mathcal{O}(M^2/R^4)$
discontinuous


$h^S = h^{S_1} + h^{S_2}$
$G(g + h^R + h^S) = 16\pi T$
$G(g) + G^1(g, h^R) + G^2(g, h^R) + G^1(g + h^R, h^S) + G^2(g, h^S) = 16\pi T$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^S) + G^2(g, h^S) - 16\pi T]$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^{S_1}) - 16\pi T + G^1(g + h^R, h^{S_2}) + G^2(g, h^S)]$
$G^1(g, h^R) = -G^2(g, h^R) + S_{eff}$
$O\left(\frac{M^2}{R^4}\right)$
C^0
$O(M^2/R^4)$
discontinuous


$h^S = h^{S_1} + h^{S_2}$
$G(g + h^R + h^S) = 16\pi T$
$G(g) + G^1(g, h^R) + G^2(g, h^R) + G^1(g + h^R, h^S) + G^2(g, h^S) = 16\pi T$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^S) + G^2(g, h^S) - 16\pi T]$
$G^1(g, h^R) = -G^2(g, h^R) - [G^1(g + h^R, h^{S_1}) - 16\pi T + G^1(g + h^R, h^{S_2}) + G^2(g, h^{S_1})]$
$G^1(g, h^R) = -G^2(g, h^R) + S_{EH}$
$O\left(\frac{M^2}{R^4}\right)$
C^0
$O(M^2/R^4)$
discontinuous