Title: Condensed Matter Review - Lecture 15

Date: Jan 21, 2011 10:15 AM

URL: http://pirsa.org/11010097

Abstract:


perimeter scholars
international


T
300
200
100
AF
Pseudo gap
Strange Metal
(Less Strange)
F.L.
d-wave sc
0.1
0.2
0.3
x (holes)

1.) T vs x ∈ fraction
of holes/cell x=0
→ 1 el/cell

2) Around x=0 AFM
quickly destroyed by
doping with holes


gap
F.L.
d-wave
SC
0.1
x (holes)


AF
gap
F.L.
d-wave
SC
0.1
0.2
0.3
X (holes)
CuO_2
2) Around
quick
dop

2) Around x=0 AFM quickly destroyed by doping with holes

- x (holes)

CuO_2



$YBaCuO_{6+x}$

1.) T_c as x ← function of holes/cell $x=0$ → 1 el/cell

2.) Around $x=0$ AFM quickly destroyed by doping with holes

3. D-wave SC
- $T_c > 0$ for some small x
- Optimal T_c
- T_c disappears at large x

(... strange)

L.

x (holes)

CuO_2


T
300
200
100
AF
Pseudo gap
Strange Metal
(Less Strange)
F.L.
d-wave SC
0.1
0.2
0.3
x (holes)
CuO_2

strange)

X (holes)



1.) T vs x ∈ fraction of holes/site x=0
→ 1 el/site

2.) Around x=0 AFM
quickly destroyed by
doping with holes

3. D-wave SC
- $T_c > 0$ for some small x
- Optimal T_c
- T_c disappears at large x

4. Pseudogap region with


R
T
AFM
SC


R
T
AFM
SC


R
X=0.2
T


R
X=0.2
T


300
00
100
AF
Pseudo gap
Strange Metal
(Less Strange)
F.L.
d-wave SC
0.1
0.2
0.3
1,1 T
2) A
3.
4, Pseu


R
X=0.2
T
Questions:
How is AFM destroyed by doping
What's going on in P.G. phase?


R
X=0.2
T_c(0.2)
T
Questions:
How is AFM destroyed by doping
What's going on in P.G. phase?
How does d-wave S.C. arise from P.G. phase?


Doping an AFM Mott Insulator
with Holes.


Doping an AFM Mott Insulator
with Holes.


Doping an AFM Mott Insulator
with Holes.


Doping an AFM Mott Insulator
with Holes.

Doping an AFM Mott Insulator with Holes.



nce Bond State

Anderson 1973,

ekas + PWA 1974



Questions:

- How is AFM destroyed by dop
- What's going on in P.G. phas
- How does d-wave S.C. arise from P.G. phase?

Valence Bond States

P.W. Anderson 1973,

Fazekas + PWA 1974

Neel State

Valence Bond Stat




Valence Bond States
P.W. Anderson 1973,
Fazekas + PWA 1974
Neel State
Valence Bond Stat
$E = -JS^z S^z$
For Spin S
$\frac{E_N}{N} = -zJS^2$
Notes for Square latt


Valence Bond States
P.W. Anderson 1973,
Fazekas + PWA 1974
Neel State
Valence Bond State


Valence Bond States
P.W. Anderson 1973,
Fazekas + PWA 1974
Neel State
Valence Bond Stat


nce Bond States
Anderson 1973,
gekas + PWA 1974
leel State
Valence Bond State
$J\vec{S}_i \cdot \vec{S}_j = \frac{J}{2}\left[(S_i+S_j)^2 - S_i^2\right.$
$= \frac{J}{2}$
$\frac{1}{2}JS^2$
for Square Latt


ond States
m 1973,
PWA 1974
ite
Valence Bond State
$J\vec{S}_i\cdot\vec{S}_j = \frac{J}{2}\left[(S_i+S_j)^2 - S_i^2 - S_j^2\right]$
0 for singlet
$= \frac{J}{2} 2S(S+1)\frac{N}{2}$


ond States
m 1973,
PWA 1974
ete
Valence Bond State
$J\vec{S}_i\cdot\vec{S}_j = \frac{J}{2}\left[(\vec{S}_i+\vec{S}_j)^2 - \vec{S}_i^2 - \vec{S}_j^2\right]$
0 for singlet
$E_{VB} = -\frac{J}{2}z\,S(S+1)\frac{N}{2}$
$\frac{E_{VB}}{N} = -\frac{J}{2}z\,S(S+1)$


Doping an AFM Mott Insulator
with Holes.
$E_{Gb} \leq E_{Neel}$
$2S^z \leq S(S+1)/2$

Doping an AFM Mott Insulator with Holes.



So $E_{Gb} \leq E_{Neel}$

$6 \cdot 2S^2 \leq 8(S+1)/2$

$\frac{3}{2} S^2 \leq \frac{1}{2} S(S+1) \rightarrow S \leq \frac{1}{3}$


But
So $E_{GS} \leq E_{Neel}$
$-6 \cdot 2S^2 \leq -8(S+1)/2$
$\frac{3}{2} \leq \frac{1}{2} \rightarrow S \leq \frac{1}{3}$

But - Frustration
(e.g. 2nd neighbor AF int's

So $E_{GS} \leq E_{Neel}$

$\langle \quad \rangle \; 2S^2 \leq 8(S+1)/2$

$\frac{3}{2} \leq \frac{1}{2} \rightarrow S \leq \frac{1}{3}$

But - Frustration
(e.g. 2nd neighbor AF int)

- Quantum Effects

So $E_{GS} \leq E_{Neel}$

$6 \cdot 2S^2 \leq 8(S+1)/2$

$\frac{3S}{2} \leq \frac{1}{2} \rightarrow S \leq \frac{1}{3}$


1973,
OA 1974
Valence Bond State
of a singlet


1973,
A 1974
Valence Bond State
$J\vec{S}_i\cdot\vec{S}_j = \frac{J}{2}\left[(\vec{S}_i+\vec{S}_j)^2 - S_i^2 - S_j^2\right]$
of a singlet
$E_{VB} = -\frac{J}{2}z\,S(S+1)\frac{N}{2}$
$\frac{E_{VB}}{N} = -\frac{J}{2}S(S+1)$
RVB


But
- Frustration
(e.g. 2nd neighbor AF inter
- Quantum Effects
- Holes (confined for AFM
So $E_{Gb} \leq E_{Neel}$
$\frac{1}{2} 2S^2 \leq S(S+1)/2$
$\frac{3S}{2} \leq \frac{1}{2} \rightarrow S \leq \frac{1}{3}$
Valenc
P.W. And
Fazeka
Neel
$E = -JS^z S^z$
For spin S
$\frac{E_N}{N_{bds}} = -ZJS^2$
for Spi


But
- Frustration
(e.g. 2nd neighbor AF int
- Quantum Effects
- Holes (confined for AFM
but not for RVB)
So $E_{Gb} \leq E_{Neel}$
$\frac{3S}{2} \leq \frac{1}{2} \rightarrow S \leq \frac{1}{3}$
Valenc
P.W. And
Fazeka
Neel
$E_N = -2JS^2$


Spin-Charge
Valenc
P.W. And
Farekha
Neel
$H = JS^zS^z$
For spins
$\frac{E_N}{N_{sites}} = -.25J$


Spin-Charge Separation
Valenc
P.W. And
Farjeha
Neel
$H = -JS^zS^z$
$\frac{E_N}{N_{sites}} = -2JS^2$


Spin-Charge Separation
Valenc
P.W. And
Farjeha
Neel


Spin-Charge Separation
hole
Valenc
P.W. And
Farekha
Neel
for spins


Spin-Charge Separation
hole
spinon


P.W.A et al J. Phys Cond Matt
16 R755 (2004) "Plain Vanilla RVB"