Title: Explorations in Numerical Relativity - Lecture 3

Date: Apr 06, 2011 11:30 AM

URL: http://pirsa.org/11040042

Abstract:

perimeter scholars
international

Richardson Extrapolation

- When it works, Richardson extrapolation has an almost magical quality about it
- However, generally have to start with fairly accurate (on the order of a few %) solutions in order to see the dramatic improvement in accuracy suggested by (84).
- Still a struggle to achieve that sort of accuracy (i.e. a few %) for *any* computation in many areas of numerical relativity/astrophysics *and* keep the error smooth (which is necessary for Richardson extrapolation to be effective)
- Thus, techniques based on Richardson extrapolation have not had a major impact in this context, although higher-order $O(h^4)$, $O(h^6)$ etc. finite difference methods *are* increasingly common for the vacuum Einstein equations

Richardson Extrapolation

- When it works, Richardson extrapolation has an almost magical quality about it
- However, generally have to start with fairly accurate (on the order of a few %) solutions in order to see the dramatic improvement in accuracy suggested by (84).
- Still a struggle to achieve that sort of accuracy (i.e. a few %) for *any* computation in many areas of numerical relativity/astrophysics *and* keep the error smooth (which is necessary for Richardson extrapolation to be effective)
- Thus, techniques based on Richardson extrapolation have not had a major impact in this context, although higher-order $O(h^4)$, $O(h^6)$ etc. finite difference methods *are* increasingly common for the vacuum Einstein equations

Richardson Extrapolation

- When it works, Richardson extrapolation has an almost magical quality about it
- However, generally have to start with fairly accurate (on the order of a few %) solutions in order to see the dramatic improvement in accuracy suggested by (84).
- Still a struggle to achieve that sort of accuracy (i.e. a few %) for *any* computation in many areas of numerical relativity/astrophysics *and* keep the error smooth (which is necessary for Richardson extrapolation to be effective)
- Thus, techniques based on Richardson extrapolation have not had a major impact in this context, although higher-order $O(h^4)$, $O(h^6)$ etc. finite difference methods *are* increasingly common for the vacuum Einstein equations


$u_{,t} = u_{,x}$
D


$u_{,t} = u_{,x}$
$D u|_j \; \frac{u_{j+1} - u_{j-1}}{2\Delta x}$


$u_{,t} = u_{,x}$
$D u|_j \; \frac{u_{j+1} - u_{j-1}}{2\Delta x}$
$u_{,t} = 2u_{,x}$

Independent Residual Evaluation

- Question that often arises in convergence testing: is the following:

  "OK, you've established that u^h is converging as $h \to 0$, but how do you know you're converging to u, the solution of the continuum problem?"

- Here, notion of an independent residual evaluation is very useful.

- Idea is as follows: have continuum PDE

$$Lu - f = 0 \tag{85}$$

  and FDA

$$L^h u^h - f^h = 0 \tag{86}$$

- Assume that u^h is apparently converging from, for example, computation of convergence factor (78) that looks like it tends to 4 as h tends to 0.

- However, do not know if we have derived and/or implemented our discrete operator L^h correctly.

Independent Residual Evaluation

- Question that often arises in convergence testing: is the following:

  "OK, you've established that u^h is converging as $h \to 0$, but how do you know you're converging to u, the solution of the continuum problem?"

- Here, notion of an independent residual evaluation is very useful.

- Idea is as follows: have continuum PDE

$$Lu - f = 0 \tag{85}$$

  and FDA

$$L^h u^h - f^h = 0 \tag{86}$$

- Assume that u^h is apparently converging from, for example, computation of convergence factor (78) that looks like it tends to 4 as h tends to 0.

- However, do not know if we have derived and/or implemented our discrete operator L^h correctly.

Independent Residual Evaluation

- Note that implicit in the "implementation" is the fact that, particularly for multi-dimensional and/or implicit and/or multi-component FDAs, considerable "work" (i.e. analysis and coding) may be involved in setting up and solving the algebraic equations for u^h.

- As a check that solution *is* converging to u, consider a *distinct* (i.e. independent) discretization of the PDE:

$$\tilde{L}^h \tilde{u}^h - f^h = 0 \tag{87}$$

- Only thing needed from this FDA for the purposes of the independent residual test is the new FD operator $\tilde{L}^h$.

- As with L^h, can expand $\tilde{L}^h$ in powers of the mesh spacing:

$$\tilde{L}^h = L + h^2 E_2 + h^4 E_4 + \cdots \tag{88}$$

where E_2, E_4, $\cdots$ are higher order (involve higher order derivatives than L) differential operators


$u_{,t} = u_{,x}$
$D u|_j \approx \frac{u_{j+1} - u_{j-1}}{2\Delta x}$
$u_{,t} = 2 u_{,x}$
D_x


$u_{,t} = u_{,x}$
D_x
$Du|_j \approx \frac{u_{j+1} - u_{j-1}}{2\Delta x}$
$u_{,t} = 2u_{,x}$


for Atoms
Molecule!
$u_{,t} = \ddot{u}_{,x}$
D_x
$D u|_j$
$\frac{u_{j+1} - u_{j-1}}{2\Delta x}$
$u_{,t} = 2u_{,x}$


for Atoms
Molecules
$u_{,t} = u_{,x}$
D_x
$D u|_j$
$\frac{u_{j+1} - u_{j-1}}{2\Delta x}$
$u_{,t} = 2u_{,x}$

Independent Residual Evaluation

- Note that implicit in the "implementation" is the fact that, particularly for multi-dimensional and/or implicit and/or multi-component FDAs, considerable "work" (i.e. analysis and coding) may be involved in setting up and solving the algebraic equations for u^h.

- As a check that solution *is* converging to u, consider a *distinct* (i.e. independent) discretization of the PDE:

$$\tilde{L}^h \tilde{u}^h - f^h = 0 \tag{87}$$

- Only thing needed from this FDA for the purposes of the independent residual test is the new FD operator $\tilde{L}^h$.

- As with L^h, can expand $\tilde{L}^h$ in powers of the mesh spacing:

$$\tilde{L}^h = L + h^2 E_2 + h^4 E_4 + \cdots \tag{88}$$

where E_2, E_4, $\cdots$ are higher order (involve higher order derivatives than L) differential operators

Pirsa: 11040042

Independent Residual Evaluation

- Conversely, assume there is a problem in the derivation and/or implementation of $L^h u^h = f^h = 0$, but there is still convergence; i.e. for example,

$$u^{2h} - u^h \to 0 \quad \text{as} \quad h \to 0 \tag{93}$$

- Then must have something like

$$u^h = u + e_0 + h e_1 + h^2 e_2 + \cdots \tag{94}$$

where crucial fact is that the error must have an $O(1)$ component, e_0.

- In this case, will compute

$$\begin{aligned} \tilde{L}^h u^h &= \left(L + h^2 E_2 + O(h^4)\right)\left(u + e_0 + h e_1 + h^2 e_2 + O(h^4)\right) \\ &= Lu + Le_0 + hLe_1 + O(h^2) \\ &= Le_0 + O(h) \end{aligned}$$

Independent Residual Evaluation

- Conversely, assume there is a problem in the derivation and/or implementation of $L^h u^h = f^h = 0$, but there is still convergence; i.e. for example,

$$u^{2h} - u^h \to 0 \quad \text{as} \quad h \to 0 \tag{93}$$

- Then must have something like

$$u^h = u + e_0 + h e_1 + h^2 e_2 + \cdots \tag{94}$$

where crucial fact is that the error must have an $O(1)$ component, e_0.

- In this case, will compute

$$\begin{aligned} \tilde{L}^h u^h &= \left(L + h^2 E_2 + O(h^4)\right)\left(u + e_0 + h e_1 + h^2 e_2 + O(h^4)\right) \\ &= Lu + Le_0 + hLe_1 + O(h^2) \\ &= Le_0 + O(h) \end{aligned}$$

Applications Places System Wed Apr
obe Reader
Tools Window Help
62.6% 72 (73 of 83) Find

Independent Residual Evaluation

- Conversely, assume there is a problem in the derivation and/or implementation of $L^h u^h = f^h = 0$, but there is still convergence; i.e. for example,

$$u^{2h} - u^h \to 0 \quad \text{as} \quad h \to 0 \tag{93}$$

- Then must have something like

$$u^h = u + e_0 + h e_1 + h^2 e_2 + \cdots \tag{94}$$

where crucial fact is that the error must have an $O(1)$ component, e_0.

- In this case, will compute

$$\begin{aligned} \tilde{L}^h u^h &= \left(L + h^2 E_2 + O(h^4)\right)\left(u + e_0 + h e_1 + h^2 e_2 + O(h^4)\right) \\ &= Lu + Le_0 + hLe_1 + O(h^2) \\ &= Le_0 + O(h) \end{aligned}$$

- Unless one is *extraordinarily* (un) lucky, and $L\,e_0$ vanishes, will *not* observe the

Applications Places System Wed Apr
obe Reader
Tools Window Help
138% 1 / 29 Find

PHY 3E7M THE 3+1 (ADM / HAMILTONIAN) FORM of GR

REFERENCES

(1) MTW CHAPTER 21

(2) YORK, "KINEMATICS & DYNAMICS of GR" IN "SOURCES of GRAV. RADIATION" (L. SMARR ed)

(3) WALD, APP. E.2 ; CHAPTER 10

(4) ARNOWITT, DESER & MISNER (1962) "THE DYN of GR" IN "GRAVITATION: AN INTRODUCTION TO CURRENT RESEARCH" (L. WITTEN ed)



Applications Places System
Lec2.pdf - Adobe Reader
/ 29

PHY 3E7M THE 3+1

REFERENCES

(1) ATMI CHART

(2) YORK, "KLYS


Applications Places System Wed Apr 6, 11:44 AM
Lec2.pdf - Adobe Reader
Edit View Document Tools Window Help
1.pdf
Lec2.pdf
1 / 29
Lec2.pdf
CHAPTER 21
"KINEMATICS & DYNAMICS of G.R." IN
"SOURCES of GRAV. RADIATION" (L. SMARR ed)
(3) WALD, APP. E.2 ; CHAPTER 10
(4) ARNOWITT, DESER & MISNER (1962) "THE DYNAMICS
of GR" IN "GRAVITATION: AN INTRODUCTION TO
CURRENT RESEARCH" (L. WITTEN ed)





Applications Places System Wed Apr
obe Reader
Tools Window Help
138% 1 / 29 Find

PHY 3E7M THE 3+1 (ADM / HAMILTONIAN) FORM of GR

REFERENCES

(1) MTW CHAPTER 21

(2) YORK, "KINEMATICS & DYNAMICS of GR" IN "SOURCES of GRAV. RADIATION" (L. SMARR ed)

(3) WALD, APP. E.2 ; CHAPTER 10

(4) ARNOWITT, DESER & MISNER (1962) "THE DYN of GR" IN "GRAVITATION: AN INTRODUCTION TO CURRENT RESEARCH" (L. WITTEN ed)

Applications Places System Wed Apr
obe Reader
Tools Window Help
66.7% 1 / 29 Find

PHYSICS THE 3+1 (ADM/HAMILTONIAN) FORM of GR ⑥

REFERENCES

(1) MTW CHAPTER 21

(2) YORK, "KINEMATICS & DYNAMICS of GR" IN "SOURCES of GRAV. RADIATION" (L. SMARR ed)

(3) WALD, APP. E.2 ; CHAPTER 10

(4) ARNOWITT, DESER & MISNER (1962) "THE DYNAMICS of GR" IN "GRAVITATION: AN INTRODUCTION TO CURRENT RESEARCH" (L. WITTEN ed)

• ADM MOTIVATION WAS, AGAIN, PREPARATION FOR QUANTIZATION of GR; FORMALISM TURNED OUT TO BE EXCELLENT BASIS FOR COMPUTATIONAL (NUMERICAL) ASSAULT ON EINSTEIN EQUATIONS

APPROACHES

(1) "COORDINATE-FUL" (MTW) : INTUITIVE, CONNECTS MORE DIRECTLY TO FORM of E.O.M USED IN PRACTICE

(2) "COORDINATE-FREE" (YORK/WALD) : PREFERABLE

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 1 / 29 Find

PHY 3674 THE 3+1 (ADM / HAMILTONIAN) FORM of GR

REFERENCES

(1) MTW CHAPTER 21

(2) YORK, "KINEMATICS & DYNAMICS of GR" IN "SOURCES of GRAV. RADIATION" (L. SMARR ed)

(3) WALD, APP. E.2 ; CHAPTER 10

(4) ARNOWITT, DESER & MISNER (1962) "THE DYNAMICS of GR" IN "GRAVITATION: AN INTRODUCTION TO CURRENT RESEARCH" (L. WITTEN ed)

• ADM MOTIVATION WAS, AGAIN, PREPARATION FOR QUANTIZATION of GR; FORMALISM TURNED OUT TO BE EXCELLENT BASIS FOR COMPUTATIONAL (NUMERICAL)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 1 / 29 Find

PHY 3674 THE 3+1 (ADM / HAMILTONIAN) FORM of GR

REFERENCES

(1) MTW CHAPTER 21

(2) YORK, "KINEMATICS & DYNAMICS of GR" IN "SOURCES of GRAV. RADIATION" (L. SMARR ed)

(3) WALD, APP. E.2 ; CHAPTER 10

(4) ARNOWITT, DESER & MISNER (1962) "THE DYNAMICS of GR" IN "GRAVITATION: AN INTRODUCTION TO CURRENT RESEARCH" (L. WITTEN ed)

• ADM MOTIVATION WAS, AGAIN, PREPARATION FOR QUANTIZATION of GR; FORMALISM TURNED OUT TO BE EXCELLENT BASIS FOR COMPUTATIONAL (NUMERICAL)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 1 / 29 Find

PHY 3E7M THE 3+1 (ADM / HAMILTONIAN) FORM of GR

REFERENCES

(1) MTW CHAPTER 21

(2) YORK, "KINEMATICS & DYNAMICS of GR" IN "SOURCES of GRAV. RADIATION" (L. SMARR ed)

(3) WALD, APP. E.2 ; CHAPTER 10

(4) ARNOWITT, DESER & MISNER (1962) "THE DYNAMICS of GR" IN "GRAVITATION: AN INTRODUCTION TO CURRENT RESEARCH" (L. WITTEN ed)

- ADM MOTIVATION WAS, AGAIN, PREPARATION FOR QUANTIZATION of GR; FORMALISM TURNED OUT TO BE EXCELLENT BASIS FOR COMPUTATIONAL (NUMERICAL)


∇B=0




∇B=0


DB=0
∂t B = -Dn E
∂t E = Dn B.


∇·B = 0
∇·E = 0


∇·B=0
∂tB = -∇×E
∇·E=0
∂tE = ∇×B.


∇·B = 0
∇·E = 0


∇·B = 0
∇·E = 0


$\nabla \cdot B = 0$
$\partial_t B = -\nabla \times E$
$\nabla \cdot E = 0$
$\partial_t E = \nabla \times B$
$\nabla_a F^{ab} = 0$; $\nabla_a {}^*F^{ab} = 0$


$\nabla \cdot B = 0$
$\partial_t B = -\nabla \times E$
$\nabla \cdot E = 0$
$\partial_t E = \nabla \times B$
$\nabla_a F^{ab} = 0 \; ; \; \nabla_a {}^{*}F^{ab} = 0$


$\nabla \cdot B = 0$
$\partial_t B = -\nabla \times E$
$\nabla \cdot E = 0$
$\partial_t E = \nabla \times B$
$\partial_a F^{ab} = 0$; $\partial_a {}^*F^{ab} = 0$


$\nabla \cdot B = 0$ $\partial_t B = -\nabla \wedge E$
$\nabla \cdot E = 0$ $\partial_t E = \nabla \wedge B$
$\nabla_a F^{ab} = 0$; $\nabla_a {}^*F^{ab} = 0$
$G_{ab} = 8\pi T_{ab}$


M
$D\cdot B = 0$ $\partial_t B = -D \wedge E$
$D\cdot E = 0$ $\partial_t E = D \wedge B$
$D_a F^{ab} = 0$; $D_a {}^*F^{ab} = 0$
$G_{ab} = 8\pi T_{ab}$


$\{M, g_{ab}\}$
$D \cdot B = 0$
$\partial_t B = -D \wedge E$
$D \cdot E = 0$
$\partial_t E = D \wedge B$
$D_a F^{ab} = 0 \; ; \; D_a {}^*F^{ab} = 0$
$G_{ab} = 8\pi T_{ab}$


$\{M_{ij}^{ab}\}$
$D\vec{B}=0$ $\partial_t \vec{B} = -D\wedge E$
$D\vec{E}=0$ $\partial_t \vec{E} = D\wedge B$
$D_a F^{ab} = 0$; $D_a {}^{*}F^{ab} = 0$
$G_{ab} = 8\pi T_{ab}$


$\{M_{ij}^{ab}\}$
$\nabla \cdot B = 0$
$\partial_t B = -\nabla \times E$
$\nabla \cdot E = 0$
$\partial_t E = \nabla \times B$
$\nabla_a F^{ab} = 0 \; ; \; \nabla_a {}^{*}F^{ab} = 0$
$G_{ab} = 8\pi T_{ab}$
$R = R_{ab} = \sum_i g_{ab} \cdots$


$\{M, g_{ab}\}$
$\nabla \cdot B = 0$
$\partial_t B = -\nabla \times E$
$\nabla \cdot E = 0$
$\partial_t E = \nabla \times B$
$\nabla_a F^{ab} = 0 \; ; \; \nabla_a {}^{*}F^{ab} = 0$
$G_{ab} = 8\pi T_{ab}$
R_{ab}
$\partial_t^2 g_{ab} \cdots$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 1 / 29 Find

PHY 3E7M THE 3+1 (ADM / HAMILTONIAN) FORM of GR

REFERENCES

(1) MTW CHAPTER 21

(2) YORK, "KINEMATICS & DYNAMICS of GR" IN "SOURCES of GRAV. RADIATION" (L. SMARR ed)

(3) WALD, APP. E.2 ; CHAPTER 10

(4) ARNOWITT, DESER & MISNER (1962) "THE DYNAMICS of GR" IN "GRAVITATION: AN INTRODUCTION TO CURRENT RESEARCH" (L. WITTEN ed)

• ADM MOTIVATION WAS, AGAIN, PREPARATION FOR QUANTIZATION of GR; FORMALISM TURNED OUT TO BE EXCELLENT BASIS FOR COMPUTATIONAL (NUMERICAL)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 1 / 29 Find

(4) Ap... of GR" IN "GRAVITATION: AN INTRODUCTION TO CURRENT RESEARCH" (L. WITTEN ed)

- ADM MOTIVATION WAS, AGAIN, PREPARATION FOR QUANTIZATION of GR; FORMALISM TURNED OUT TO BE EXCELLENT BASIS FOR COMPUTATIONAL (NUMERICAL) ASSAULT ON EINSTEIN EQUATIONS

APPROACHES

(1) "COORDINATE-FUL" (MTW): INTUITIVE, CONNECTS MORE DIRECTLY TO FORM of E.O.M USED IN PRACTICE

(2) "COORDINATE FREE" (YORK/WALD): PREFERABLE FOR DERIVATION of E.O.M.

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 1 / 29 Find

STAY COORDINATE FREE FOREVER

- WILL ~~START~~ WITH COORDINATE BASED APPROACH TO INTRODUCE CONCEPTS, THEN WILL GO OVER TO COORDINATE-FREE APPROACH TO DERIVE 3+1 EQNS

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 2 / 29 Find

$$G_{ab} = 8\pi T_{ab}$$

AS SYSTEM of FIRST-ORDER (IN TIME) PDE'S FOR THE GRAV. FIELD VBL'S, WHICH CAN THEN BE SOLVED AS AN "INITIAL-VALUE" or "CAUCHY" PROBLEM

SHIFT IN PERSPECTIVE: UP TO NOW $G_{ab} = 8\pi T_{ab}$ DESCRIBED THE LINKAGE OF THE GEOMETRY of SPACETIME (4-D) TO THE DISTRIBUTION of MATTER-STRESS-ENERGY IN S.T.

NEW VIEW: GEOMETRY of S.T IS "TIME-HISTORY" (EVOLUTION) of GEOMETRY of A SPACELIKE HYPERSURFACE (3-D) =) GEOMETRODYNAMICS; "VIEW" NOT UNIQUE SINCE THERE ARE MANY MANY WAYS of "SLICING UP" GIVEN S.T. INTO A




M


Signature (−,+++)
M

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 2 / 29 Find

$$G_{ab} = \varepsilon_{\pi} T_{ab}$$

AS SYSTEM of FIRST-ORDER (IN TIME) PDE'S FOR THE GRAV. FIELD VBL'S, WHICH CAN THEN BE SOLVED AS AN "INITIAL-VALUE" OR "CAUCHY" PROBLEM

SHIFT IN PERSPECTIVE: UP TO NOW $G_{ab} = \varepsilon_{\pi} T_{ab}$ DESCRIBED THE LINKAGE OF THE GEOMETRY of SPACETIME (4-D) TO THE DISTRIBUTION of MATTER-STRESS-ENERGY IN S.T.

NEW VIEW: GEOMETRY of S.T IS "TIME-HISTORY" (EVOLUTION) of GEOMETRY of A SPACELIKE HYPERSURFACE (3-D) =)

GEOMETRODYNAMICS; "VIEW" NOT UNIQUE SINCE THERE ARE MANY MANY WAYS of "SLICING UP" GIVEN S.T. INTO A

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 2 / 29 Find

THE LINKAGE OF THE GEOMETRY of SPACETIME (4-D)
TO THE DISTRIBUTION of MATTER-STRESS-ENERGY IN S.T

NEW VIEW: GEOMETRY of S.T IS "TIME-HISTORY" (EVOLUTION)
of GEOMETRY of A SPACELIKE HYPERSURFACE (3-D) =)
GEOMETRODYNAMICS; "VIEW" NOT UNIQUE SINCE THERE ARE
CO'NLY MANY WAYS of "SLICING UP" GIVEN S.T. INTO A
FAMILY OF S.L. HYPERSURFACES; SUCH PREFERRED (i.e.
DYNAMICALLY MORE RELEVANT) IN GENERAL

SPLITTING SPACETIME INTO SPACE-PLUS-TIME (THE 3+1 SPLIT)

- SPACETIME IS 4-DIMENSIONAL MANIFOLD M, WITH LORENTZIAN SIGNATURE METRIC g_{ab} (−+++)
- INTRODUCE COORDINATES $\{x^\mu\} = \{t, x^i\}$ (MAY NOT COVER ENTIRE S.T., BUT WILL GENERALLY PRETEND THEY DO)

NOTATION/CONVENTIONS

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 2 / 29 Find

SPLITTING SPACETIME INTO SPACE-PLUS-TIME (THE 3+1 SPLIT)

- SPACETIME IS 4-DIMENSIONAL MANIFOLD M, WITH LORENTZIAN SIGNATURE METRIC g_{ab} $(-+++)$
- INTRODUCE COORDINATES $\{x^a\} = \{t, x^i\}$ (MAY NOT COVER ENTIRE S.T., BUT WILL GENERALLY PRETEND THEY DO)

NOTATION / CONVENTIONS:

GREEK INDICES: $(\mu, \nu$ etc.) $0, 1, 2, 3$

"INTEGER" LATIN " (i, j, k, l, m, n) $1, 2, 3$ (SPATIAL)

WILL ADOPT USUAL EINSTEIN SUMMATION CONVENTION FOR BOTH TYPES!

- DEMAND THAT $t =$ CONST SURFACES (HYPERSURFACES), $\Sigma(t)$, ARE SPACELIKE; I.E. IF DISTINCT EVENTS P, P'


M


M
P(t, x)


ature (−,+++)
M
g_{ab}
θ_{ij}
$P(t, x^i)$


(9) g_{ab}
(3) g_{ij}
M
P(t, x)


(9) g_{ab}
(3) g_{ij}
M

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 3 / 29 Find

- EACH $\Sigma(t)$ IS A DIFFERENTIABLE MANIFOLD IN ITS OWN RIGHT, WITH A 3-METRIC ${}^{(3)}g_{ij} = {}^{(3)}g_{(ij)}$ WHICH IS INDUCED ON $\Sigma(t)$ BY THE 4-METRIC ${}^{(4)}g_{\mu\nu}$ OF THE ENVELOPING SPACETIME

- ANY GIVEN FOLIATION (= CHOICE OF TIME COORD = CHOICE OF SLICING) ALSO DEFINES A NATURAL, UNIT-NORM VECTOR FIELD, n^μ, WHICH IS NORMAL (ORTHOGONAL) TO THE SLICES, AND POINTS "TO THE FUTURE"

$$n^\mu n_\mu = {}^{(4)}g_{\mu\nu} n^\mu n^\nu = -1$$

SPACETIME DISPLACEMENT IN THE 3+1 SPLIT

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 3 / 29 Find

WHICH IS INDUCED ON $\Sigma(t)$ BY THE 4-METRIC
(4) $g_{\mu\nu}$ OF THE ENVELOPING SPACETIME

ANY GIVEN FOLIATION (= CHOICE OF TIME COORD = CHOICE OF SLICING) ALSO DEFINES A NATURAL, UNIT-NORM VECTOR FIELD, n^μ, WHICH IS NORMAL (ORTHOGONAL) TO THE SLICES, AND POINTS "TO THE FUTURE"

$$n^\mu n_\mu \overset{(4)}{=} g_{\mu\nu} n^\mu n^\nu = -1$$

SPACETIME DISPLACEMENT IN THE 3+1 SPLIT



Pirsa: 11040042 Page 65/135

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 3 / 29 Find

$^{(4)}g_{\mu\nu}$ OF THE ENVELOPING SPACETIME

ANY GIVEN FOLIATION (= CHOICE OF TIME COORD = CHOICE OF SLICING) ALSO DEFINES A NATURAL, UNIT-NORM VECTOR FIELD, n^μ, WHICH IS NORMAL (ORTHOGONAL) TO THE SLICES, AND POINTS "TO THE FUTURE"

$$n^\mu n_\mu = {}^{(4)}g_{\mu\nu} n^\mu n^\nu = -1$$

SPACETIME DISPLACEMENT IN THE 3+1 SPLIT



Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 3 / 29 Find

OF SLICING) ALSO DEFINES A NATURAL, UNIT-NORM VECTOR FIELD, n^μ, WHICH IS NORMAL (ORTHOGONAL) TO THE SLICES, AND POINTS "TO THE FUTURE"

$$n^\mu n_\mu = {}^{(4)}g_{\mu\nu} n^\mu n^\nu = -1$$

SPACETIME DISPLACEMENT IN THE 3+1 SPLIT




M
Σ(t)
P(t, x)


M
P(t, x)


ture (−,+++)
M
$\nabla_\mu t = n_\mu$
(1) $g_{\mu\nu}$
(3) g_{iT}
P(t, x^i)


ture (−,+++)
M
(1) $g_{\mu\nu}$
$\nabla_\mu t = (\#) n_\mu$
$n^\mu n_\mu = -1$
P(t, x^i)


ture (−,+++)
M
(1) $g_{\mu\nu}$
(3) g_{iT}
$\nabla_\mu t = (\#) n_\mu$
$n^\mu n_\mu = -1$
P(t, x)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 3 / 29 Find

OF SLICING) ALSO DEFINES A NATURAL, UNIT-NORM VECTOR FIELD, n^μ, WHICH IS NORMAL (ORTHOGONAL) TO THE SLICES, AND POINTS "TO THE FUTURE"

$$n^\mu n_\mu = {}^{(4)}g_{\mu\nu} n^\mu n^\nu = -1$$

SPACETIME DISPLACEMENT IN THE 3+1 SPLIT




M
$\nabla_\mu t = (*) n_\mu$
$n^a n_a = -1$
$ds^2 = g_{\mu\nu} dx^\mu dx^\nu$
P'
Q
$\Sigma(t)$
P(t, x^i)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 4 / 29 Find

$$= (-\alpha^2 + {}^{(3)}g_{ij}\beta^i\beta^j)\,dt^2 + 2\,{}^{(3)}g_{ij}\beta^i dx^j dt + {}^{(3)}g_{ij}dx^i dx^j \qquad (1)$$

NOMENCLATURE:

$\alpha \equiv \alpha(t, x^i)$: LAPSE FUNCTION ("LAPSE") : GIVES LAPSE OF PROPER TIME PER UNIT COORDINATE TIME FOR AN OBSERVER MOVING NORMAL TO THE SLICES

$\beta^i \equiv \beta^i(t, x^i)$: SHIFT VECTOR ("SHIFT") : 3-VECTOR DESCRIBING

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 4 / 29 Find

$$\text{distance}(P,Q)^2 = {}^{(4)}ds^2$$

$$= -\alpha^2 dt^2 + {}^{(3)}g_{ij}(dx^i + \beta^i dt)(dx^j + \beta^j dt)$$

$$= (-\alpha^2 + {}^{(3)}g_{ij}\beta^i\beta^j)dt^2 + 2\,{}^{(3)}g_{ij}\beta^i dx^j dt$$

$$+ {}^{(3)}g_{ij}dx^i dx^j \qquad (1)$$

NOMENCLATURE:

$\alpha \equiv \alpha(t, x^i)$: LAPSE FUNCTION ("LAPSE") : GIVES LAPSE OF PROPER TIME PER UNIT COORDINATE TIME FOR AN OBSERVER MOVING NORMAL TO THE SLICES

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 4 / 29 Find

"SPACETIME PYTHAGOREAN THM" ($\lim dt \to 0$) $\Rightarrow$

$$\text{distance}(P,Q)^2 \overset{(a)}{=} ds^2$$

$$= -\alpha^2 dt^2 + {}^{(3)}g_{ij}(dx^i + \beta^i dt)(dx^j + \beta^j dt)$$

$$= (-\alpha^2 + {}^{(3)}g_{ij}\beta^i\beta^j)dt^2 + 2\,{}^{(3)}g_{ij}\beta^i dx^j dt + {}^{(3)}g_{ij}dx^i dx^j \qquad (1)$$

NOMENCLATURE:

$\alpha \equiv \alpha(t, x^i)$: LAPSE FUNCTION ("LAPSE") : GIVES LAPSE OF PROPER TIME PER UNIT COORDINATE TIME FOR AN OBSERVER

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 4 / 29 Find

"SPACETIME PYTHAGOREAN THM" ($\lim dt \to 0$) $\Rightarrow$

$$\text{distance}(P,Q)^2 \overset{(a)}{=} ds^2$$

$$= -\alpha^2 dt^2 + {}^{(3)}g_{ij}(dx^i + \beta^i dt)(dx^j + \beta^j dt)$$

$$= (-\alpha^2 + {}^{(3)}g_{ij}\beta^i\beta^j)dt^2 + 2\,{}^{(3)}g_{ij}\beta^i dx^j dt + {}^{(3)}g_{ij}dx^i dx^j \qquad (1)$$

NOMENCLATURE:

$\alpha \equiv \alpha(t, x^i)$: LAPSE FUNCTION ("LAPSE") : GIVES LAPSE OF PROPER TIME PER UNIT COORDINATE TIME FOR AN OBSERVER


M
$ds^2 = g_{\mu\nu} dx^\mu dx^\nu$
P'
Q
$\Sigma(t)$
P(t, x^i)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 4 / 29 Find

"SPACETIME PYTHAGOREAN THM" ($\lim dt \to 0$) $\Rightarrow$

$$\text{distance}(P,Q)^2 \overset{(a)}{=} ds^2$$

$$= -\alpha^2 dt^2 + {}^{(3)}g_{ij}(dx^i + \beta^i dt)(dx^j + \beta^j dt)$$

$$= (-\alpha^2 + {}^{(3)}g_{ij}\beta^i\beta^j)dt^2 + 2\,{}^{(3)}g_{ij}\beta^i dx^j dt + {}^{(3)}g_{ij}dx^i dx^j \qquad (1)$$

NOMENCLATURE:

$\alpha \equiv \alpha(t, x^i)$: LAPSE FUNCTION ("LAPSE") : GIVES LAPSE OF PROPER TIME PER UNIT COORDINATE TIME FOR AN OBSERVER

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 4 / 29 Find

$$\cdots + g_{ij}\beta^i\beta^j)\,dt^2 + 2\,g_{ij}\beta^i dx^j dt$$

$$+ {}^{(3)}g_{ij}\,dx^i dx^j \qquad (1)$$

NOMENCLATURE:

$\alpha \equiv \alpha(t, x^i)$: LAPSE FUNCTION ("LAPSE"): GIVES LAPSE OF PROPER TIME PER UNIT COORDINATE TIME FOR AN OBSERVER MOVING NORMAL TO THE SLICES

$\beta^i \equiv \beta^i(t, x^i)$: SHIFT VECTOR ("SHIFT"): 3-VECTOR DESCRIBING STRETCH OF SPATIAL COORDINATES RELATIVE TO "NORMAL PROPAGATION"

TOGETHER $\{\alpha, \beta^i\}$ CONSTITUTE 4-FOLD COORD. FREEDOM

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 4 / 29 Find

NOMENCLATURE:

$\alpha \equiv \alpha(t, x^i)$: LAPSE FUNCTION ("LAPSE") : GIVES LAPSE OF PROPER TIME PER UNIT COORDINATE TIME FOR AN OBSERVER MOVING NORMAL TO THE SLICES

$\beta^i \equiv \beta^i(t, x^i)$: SHIFT VECTOR ("SHIFT") : 3-VECTOR DESCRIBING SHIFTING OF SPATIAL COORDINATES RELATIVE TO "NORMAL PROPAGATION"

* TOGETHER $\{\alpha, \beta^i\}$ CONSTITUTE 4-FOLD COORD. FREEDOM of GR

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 4 / 29 Find

NOMENCLATURE:

$\alpha \equiv \alpha(t, x^i)$: LAPSE FUNCTION ("LAPSE"): GIVES LAPSE OF PROPER TIME PER UNIT COORDINATE TIME FOR AN OBSERVER MOVING NORMAL TO THE SLICES

$\beta^i \equiv \beta^i(t, x^i)$: SHIFT VECTOR ("SHIFT"): 3-VECTOR DESCRIBING SHIFTING OF SPATIAL COORDINATES RELATIVE TO "NORMAL PROPAGATION"

- TOGETHER $\{\alpha, \beta^i\}$ CONSTITUTE 4-FOLD COORD. FREEDOM of GR

DUAL VIEWS

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 4 / 29 Find

(COORDINATE FREEDOM)

(2) SOME PRESCRIPTION FOR α, β^i MUST BE GIVEN "FROM OUTSIDE" - I.E. E.O.M. (EINSTEIN EQ^{N}) ALONE WILL NOT, IN GENERAL, DETERMINE THEM ("GAUGE FIXING")

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

NOTE: TENSORS SUCH AS β^i ARE DEFINED ON $\Sigma(t)$, AND ARE CALLED SPATIAL TENSORS

CLEARLY, ${}^{(3)}g_{ij}$ IS A SPATIAL TENSOR; IT HAS AN ASSOCIATED INVERSE ${}^{(3)}g^{ij}$ SATISFYING

$$ {}^{(3)}g^{ij}\,{}^{(3)}g_{jk} = \delta^i{}_k $$

INDICES OF 3-TENSORS ARE RAISED/LOWERED WITH ${}^{(3)}g^{ij}$, ${}^{(3)}g_{ij}$; THUS, WE CAN REWRITE (1) AS

$$ {}^{(4)}ds^2 = (-\alpha^2 + \beta^i\beta_i)\,dt^2 + 2\beta_j\,dx^j\,dt + {}^{(3)}g_{ij}\,dx^i\,dx^j $$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

NOTE: TENSORS SUCH AS β^i ARE DEFINED ON $\Sigma(t)$, AND ARE CALLED SPATIAL TENSORS

CLEARLY, ${}^{(3)}g_{ij}$ IS A SPATIAL TENSOR; IT HAS AN ASSOCIATED INVERSE ${}^{(3)}g^{ij}$ SATISFYING

$$ {}^{(3)}g^{ij}\,{}^{(3)}g_{jk} = \delta^i{}_k $$

INDICES OF 3-TENSORS ARE RAISED/LOWERED WITH ${}^{(3)}g^{ij}$, ${}^{(3)}g_{ij}$; THUS, WE CAN REWRITE (1) AS

$$ (4)\quad ds^2 = (-\alpha^2 + \beta^i\beta_i)\,dt^2 + 2\beta_i\,dx^i\,dt + {}^{(3)}g_{ij}\,dx^i\,dx^j $$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

$$ {}^{(3)}g^{ij}\,{}^{(3)}g_{jk} = \delta^{i}{}_{k} $$

INDICES OF 3-TENSORS ARE RAISED / LOWERED WITH ${}^{(3)}g^{ij}$, ${}^{(3)}g_{ij}$; THUS, WE CAN REWRITE (1) AS

$$ {}^{(4)}ds^2 = (-\alpha^2 + \beta^i\beta_i)\,dt^2 + 2\beta_j\,dx^j\,dt + {}^{(3)}g_{ij}\,dx^i dx^j $$

SO, WE HAVE

$$ {}^{(4)}g_{\mu\nu} = \begin{bmatrix} {}^{(4)}g_{00} & {}^{(4)}g_{0j} \\ {}^{(4)}g_{i0} & {}^{(4)}g_{ij} \end{bmatrix} = \begin{bmatrix} -\alpha^2 + \beta^k\beta_k & \beta_j \\ \beta_i & {}^{(3)}g_{ij} \end{bmatrix} \qquad (2) $$

IN PARTICULAR, NOTE THAT

$$ {}^{(4)}g^{00} = {}^{(3)}g^{ij}\,\frac{\beta_i}{\alpha}\,\frac{\beta_j}{\alpha} - \frac{1}{\alpha^2} $$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

$$ {}^{(3)}g^{ij}\,{}^{(3)}g_{jk} = \delta^{i}{}_{k} $$

INDICES OF 3-TENSORS ARE RAISED / LOWERED WITH ${}^{(3)}g^{ij}$, ${}^{(3)}g_{ij}$; THUS, WE CAN REWRITE (1) AS

$$ {}^{(4)}ds^2 = (-\alpha^2 + \beta^i\beta_i)\,dt^2 + 2\beta_j\,dx^j\,dt + {}^{(3)}g_{ij}\,dx^i dx^j $$

SO, WE HAVE

$$ {}^{(4)}g_{\mu\nu} = \begin{bmatrix} {}^{(4)}g_{00} & {}^{(4)}g_{0j} \\ {}^{(4)}g_{i0} & {}^{(4)}g_{ij} \end{bmatrix} = \begin{bmatrix} -\alpha^2 + \beta^k\beta_k & \beta_j \\ \beta_i & {}^{(3)}g_{ij} \end{bmatrix} \qquad (2) $$

IN PARTICULAR, NOTE THAT

$$ {}^{(4)}g^{\mu\nu} = \begin{bmatrix} -\frac{1}{\alpha^2} & \frac{\beta^j}{\alpha^2} \\ \frac{\beta^i}{\alpha^2} & {}^{(3)}g^{ij} - \frac{\beta^i\beta^j}{\alpha^2} \end{bmatrix} $$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

SO, WE HAVE

$$ {}^{(4)}g_{\mu\nu} = \begin{bmatrix} {}^{(4)}g_{00} & {}^{(4)}g_{0j} \\ {}^{(4)}g_{i0} & {}^{(4)}g_{ij} \end{bmatrix} = \begin{bmatrix} -\alpha^2 + \beta^k\beta_k & \beta_j \\ \beta_i & {}^{(3)}g_{ij} \end{bmatrix} \qquad (2) $$

IN PARTICULAR, NOTE THAT

$$ {}^{(4)}g_{ij} = {}^{(3)}g_{ij} $$

I.E. SPATIAL COVARIANT COMPONENTS OF 4- AND 3-METRICS ARE IDENTICAL

THIS IS A GENERAL RESULT; GIVEN ANY 4-TENSOR OF TYPE $(0,k)$ (COVARIANT TENSOR), THE SPATIAL COMPONENTS OF THAT TENSOR CAN BE IDENTIFIED AS THE COMPONENTS OF A TYPE $(0,k)$ 3-TENSOR

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

$$ {}^{(4)}g_{\mu\nu} = \begin{bmatrix} {}^{(4)}g_{00} & {}^{(4)}g_{0j} \\ {}^{(4)}g_{i0} & {}^{(4)}g_{ij} \end{bmatrix} = \begin{bmatrix} -\alpha^2 + \beta^k \beta_k & \beta_j \\ \beta_i & {}^{(3)}g_{ij} \end{bmatrix} \qquad (2) $$

IN PARTICULAR, NOTE THAT

$$ {}^{(4)}g_{ij} = {}^{(3)}g_{ij} $$

I.E. SPATIAL COVARIANT COMPONENTS OF 4- AND 3-METRICS ARE IDENTICAL

• THIS IS A GENERAL RESULT; GIVEN ANY 4-TENSOR OF TYPE $(0,k)$ (COVARIANT TENSOR), THE SPATIAL COMPONENTS OF THAT TENSOR CAN BE IDENTIFIED AS THE COMPONENTS OF A TYPE $(0,k)$ 3-TENSOR

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

$$\begin{bmatrix} {}^{(4)}g_{i0} & {}^{(4)}g_{ij} \end{bmatrix} \quad \begin{bmatrix} \beta_i & {}^{(3)}g_{ij} \end{bmatrix} \qquad (2)$$

IN PARTICULAR, NOTE THAT

$${}^{(4)}g_{ij} = {}^{(3)}g_{ij}$$

I.E. SPATIAL COVARIANT COMPONENTS OF 4- AND 3-METRICS ARE IDENTICAL

- THIS IS A GENERAL RESULT: GIVEN ANY 4-TENSOR OF TYPE $(0,k)$ (COVARIANT TENSOR), THE SPATIAL COMPONENTS OF THAT TENSOR CAN BE IDENTIFIED AS THE COMPONENTS OF A TYPE $(0,k)$ 3-TENSOR

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 6 / 29 Find

• THIS IS A GENERAL RESULT: GIVEN ANY 1-TENSOR OF TYPE $(0,k)$ (COVARIANT TENSOR), THE SPATIAL COMPONENTS OF THAT TENSOR CAN BE IDENTIFIED AS THE COMPONENTS OF A TYPE $(0,k)$ 3-TENSOR

WHY? RECALL THAT COVARIANT TENSOR COMPONENTS CAN BE DEFINED IN TERMS OF THE ACTION OF THE TENSOR ON THE COORDINATE BASIS VECTORS $\vec{e}_\mu = {}^{(4)}\vec{e}_\mu$, $\mu = 0, 1, 2, 3$

E.G. $T_{\mu\nu} = T(\vec{e}_\mu, \vec{e}_\nu)$

AND $T_{ij} = T(\vec{e}_i, \vec{e}_j)$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 6 / 29 Find

OF A TYPE $(0,k)$ 3-TENSOR

WHY? RECALL THAT COVARIANT TENSOR COMPONENTS CAN BE DEFINED IN TERMS OF THE ACTION OF THE TENSOR ON THE COORDINATE BASIS VECTORS $^{(a)}\vec{e}_\mu$, $\mu = 0, 1, 2, 3$

E.G. $^{(a)}t_{\mu\nu} = {}^{(a)}t(\,{}^{(a)}\vec{e}_\mu, {}^{(a)}\vec{e}_\nu)$

AND $^{(a)}t_{ij} = {}^{(a)}t(\,{}^{(a)}\vec{e}_i, {}^{(a)}\vec{e}_j)$

BUT CLEARLY THE $\{{}^{(a)}\vec{e}_i\}$ ARE DIRECTLY THE COORDINATE BASIS VECTORS $\{{}^{(3)}\vec{e}_i\}$ FOR $\Sigma(t)$ WITH COORDINATES $\{x^i\}$; THUS INTERPRETING $^{(a)}t(\ldots)$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 6 / 29 Find

$$ {}^{(2)}t_{ij} = {}^{(4)}t_{ij} $$

ON THE OTHER HAND, THE SPATIAL METRICS $\{ {}^{(4)}\underline{\omega}^i \}$ OF THE DUAL BASIS $\{ {}^{(4)}\underline{\omega}^m \}$, WILL NOT, IN GENERAL COINCIDE WITH $\{ {}^{(3)}\underline{\omega}^i \}$ - THE DUAL BASIS ON $\Sigma(t)$ DEPENDS ON HOW $\Sigma(t)$ IS EMBEDDED IN THE S.T.

• THUS, IN GENERAL

$$ {}^{(3)}t_{ij} \neq {}^{(4)}t_{ij} $$

EXAMPLE: CONSIDER THE INVERSE 4-METRIC COMPONENTS FROM (2)

$$ \begin{vmatrix} -1/N^2 & N^j/N^2 \\ \cdots & \cdots \end{vmatrix} $$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 6 / 29 Find

• THUS, IN GENERAL

$$ {}^{(3)}\Gamma_{ij} \neq {}^{(a)}\Gamma_{ij} $$

EXAMPLE : CONSIDER THE INVERSE 4-METRIC COMPONENTS FROM (2)

$$ {}^{(a)}g^{\mu\nu} = \begin{bmatrix} -1/\alpha^2 & \beta^j/\alpha^2 \\ \beta^i/\alpha^2 & {}^{(3)}g^{ij} - \beta^i\beta^j/\alpha^2 \end{bmatrix} \qquad (3) $$

(EXERCISE: VERIFY)


for Atoms
Molecules
Signature (−,+,+,+)
M
$\nabla_\mu t = (*)\, n_{\mu a}$
$n^a n_a = -1$
$ds^2 = g_{\mu\nu}\, dx^\mu dx^\nu$


for Atoms
Molecule!
Signature (−,+++)
M
$\nabla_\mu t = (*) n_{\mu\nu}$
$n^\nu n_{\nu a} = -1$
$ds^2 = g_{\mu\nu} dx^\mu dx^\nu$
$n_\mu = \alpha \nabla_\mu t$
$= (\alpha, 0, 0, 0)$
P(t,

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 6 / 29 Find

• THUS, IN GENERAL

$$ {}^{(3)}\Gamma_{ij} \neq {}^{(4)}\Gamma_{ij} $$

EXAMPLE : CONSIDER THE INVERSE 4-METRIC COMPONENTS FROM (2)

$$ {}^{(4)}g^{\mu\nu} = \begin{bmatrix} -1/\alpha^2 & \beta^j/\alpha^2 \\ \beta^i/\alpha^2 & {}^{(3)}g^{ij} - \beta^i\beta^j/\alpha^2 \end{bmatrix} \qquad (3) $$

(EXERCISE: VERIFY)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 7 / 29 Find

FROM (2)

$$ {}^{(4)}g^{\mu\nu} = \begin{bmatrix} -1/\alpha^2 & \beta^j/\alpha^2 \\ \beta^i/\alpha^2 & {}^{(3)}g^{ij} - \beta^i\beta^j/\alpha^2 \end{bmatrix} \qquad (3) $$

(EXERCISE: VERIFY)

DAY 387M THE 3+1 FORMULATION OF GR ④

- FROM (2) WE CAN ALSO COMPUTE THE USEFUL RESULT

$$ \sqrt{-{}^{(4)}g} = \alpha\sqrt{{}^{(3)}g} \qquad (4) $$

THE NORMAL VECTOR FIELD n^μ

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 7 / 29 Find

PHY 357H THE 3+1 FORMULATION of GR ④

- FROM (2) WE CAN ALSO COMPUTE THE USEFUL RESULT

$$\sqrt{-{}^{(4)}g} = \alpha \sqrt{{}^{(3)}g} \qquad (4)$$

THE NORMAL VECTOR FIELD n^μ

- EASIEST TO START WITH ASSOC. DUAL-VECTOR (ONE-FORM) FIELD, n_μ

GEOM. INTERP of DUAL VECTOR FIELD: LEVEL SURFACES of SCALAR FUNCTION → DUAL MOTION TO "INFINITESIMAL DISPLACEMENT" (VECTOR)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 7 / 29 Find

FROM (2) WE CAN ALSO COMPUTE THE USEFUL RESULT

$$\sqrt{-{}^{(4)}g} = \alpha \sqrt{{}^{(3)}g} \qquad (4)$$

THE NORMAL VECTOR FIELD n^μ

EASIEST TO START WITH ASSOC. DUAL-VECTOR (ONE-FORM) FIELD, n_μ

CLEAR, INSTEAD OF DUAL VECTOR FIELD: LEVEL SURFACES OF SCALAR FUNCTION → DUAL MOTION TO "INFINITESIMAL DISPLACEMENT" (VECTOR)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 7 / 29 Find

GEOM. INTERP. OF DUAL-VECTOR FIELD: LEVEL SURFACES of SCALAR FUNCTION ⇒ DUAL MOTION TO "INFINITESIMAL DISPLACEMENT" (VECTOR)

$\Rightarrow \langle \vec{v}, \underline{df} \rangle = v^\mu (df)_\mu$: # of level surfaces "pierced" by $\vec{v}$

• HERE OUR SCALAR FUNCTION IS THE TIME COORDINATE t, WITH ASSOCIATED DUAL-VECTOR FIELD $\underline{dt}$, THEN

$$\underline{\sim} \propto \underline{dt}$$

OR IN COMPONENT FORM

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 7 / 29 Find

$$n_\mu = (-\alpha, 0, 0, 0) \qquad (5)$$

AND THEN

$$n^\mu = {}^{(4)}g^{\mu\nu} n_\nu = \left(\frac{1}{\alpha}, -\frac{\beta^i}{\alpha} \right) \qquad (6)$$

EXTRINSIC CURVATURE

- THE INTRINSIC GEOMETRY of $\Sigma(t)$ IS DESCRIBED BY
  $^{(3)}g_{ij}$ WHICH ENCODES ALL GEOM. INFO. WHICH MAY BE DEDUCED BY MAKING MEASUREMENTS ON $\Sigma(t)$ ALONE

- HOWEVER, A GIVEN 3-GEOMETRY (SLICE, HYPERSURFACE) MAY BE EMBEDDED IN SPACETIME IN INFINITELY MANY


Applications Places System
Wed Apr
obe Reader
Tools Window Help
100%
7 / 29
Find

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 8 / 29 Find

• HOWEVER, A GIVEN 3-GEOMETRY (SLICE, HYPERSURFACE) MAY BE EMBEDDED IN SPACETIME IN INFINITELY MANY DISTINGUISHABLE (BY 4-D MEASUREMENTS) WAYS

EXAMPLE: 2D EMBEDDED IN 3D; FLAT SURFACE EMBEDDED WITH / WITHOUT EXTRINSIC CURVATURE

• THE MANNER IN WHICH $\Sigma(t)$ IS EMBEDDED CAN BE CHARACTERIZED BY INVESTIGATING THE CHANGE IN THE DIRECTION OF THE NORMAL FIELD AS A FCN OF POS ON $\Sigma(t)$ — THIS DEFINES THE EXTRINSIC CURVATURE TENSOR (AKA THE SECOND FUNDAMENTAL FORM)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 8 / 29 Find

AND THEN

$$n^\mu = {}^{(4)}g^{\mu\nu}\, l_\nu = \left(\frac{1}{\alpha}, -\frac{\beta^i}{\alpha} \right) \qquad (6)$$

EXTRINSIC CURVATURE

- THE INTRINSIC GEOMETRY of $\Sigma(t)$ IS DESCRIBED BY ${}^{(3)}g_{ij}$ WHICH ENCODES ALL GEOM. INFO. WHICH MAY BE DEDUCED BY MAKING MEASUREMENTS ON $\Sigma(t)$ ALONE

- HOWEVER, A GIVEN 3-GEOMETRY (SLICE, HYPERSURFACE) MAY BE EMBEDDED IN SPACETIME IN INFINITELY MANY DISTINGUISHABLE (BY 4-D MEASUREMENTS) WAYS

EXAMPLE: 2D EMBEDDED IN 3D; FLAT SURFACE EMBEDDED

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 8 / 29 Find

WE HAVE

SIGN CHOSEN SO THAT n^μ IS "FUTURE-DIRECTED"

$$n_\mu = (-\alpha, 0, 0, 0) \qquad (5)$$

AND THEN

$$n^\mu = {}^{(4)}g^{\mu\nu} n_\nu = \left(\frac{1}{\alpha}, -\frac{\beta^i}{\alpha} \right) \qquad (6)$$

EXTRINSIC CURVATURE

• THE INTRINSIC GEOMETRY of $\Sigma(t)$ IS DESCRIBED BY
${}^{(3)}g_{ij}$ WHICH ENCODES ALL GEOM. INFO. WHICH MAY
BE DEDUCED BY MAKING MEASUREMENTS ON $\Sigma(t)$ ALONE


for Atoms
Molecule?
Signature (−,+,+,+)
M
$\nabla_\mu t = (*) n_\mu$
$n^a n_a = -1$
$ds^2 = g_{\mu\nu} dx^\mu dx^\nu$
$n_\mu = \alpha \nabla_\mu t$
$= (-\alpha, 0, 0, 0)$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 8 / 29 Find

WE HAVE

SIGN CHOSEN SO THAT n^μ IS "FUTURE-DIRECTED"

$$n_\mu = (-\alpha, 0, 0, 0) \qquad (5)$$

AND THEN

$$n^\mu = {}^{(4)}g^{\mu\nu} n_\nu = \left(\frac{1}{\alpha}, -\frac{\beta^i}{\alpha} \right) \qquad (6)$$

EXTRINSIC CURVATURE

- THE INTRINSIC GEOMETRY of $\Sigma(t)$ IS DESCRIBED BY ${}^{(3)}g_{ij}$ WHICH ENCODES ALL GEOM. INFO. WHICH MAY BE DEDUCED BY MAKING MEASUREMENTS ON $\Sigma(t)$ ALONE

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 8 / 29 Find

$$n_\mu = (-\alpha, 0, 0, 0) \qquad (5)$$

AND THEN

$$n^\mu = {}^{(4)}g^{\mu\nu} n_\nu = \left(\frac{1}{\alpha}, -\frac{\beta^i}{\alpha}\right) \qquad (6)$$

EXTRINSIC CURVATURE

- THE INTRINSIC GEOMETRY of $\Sigma(t)$ IS DESCRIBED BY ${}^{(3)}g_{ij}$ WHICH ENCODES ALL GEOM. INFO. WHICH MAY BE DEDUCED BY MAKING MEASUREMENTS ON $\Sigma(t)$ ALONE

- HOWEVER, A GIVEN 3-GEOMETRY (SLICE, HYPERSURFACE) MAY BE EMBEDDED IN SPACETIME IN INFINITELY MANY

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 8 / 29 Find

AND THEN

$$n^\mu = {}^{(4)}g^{\mu\nu}\, l_\nu = \left(\frac{1}{\alpha}, -\frac{\beta^i}{\alpha} \right) \qquad (6)$$

EXTRINSIC CURVATURE

- THE INTRINSIC GEOMETRY of $\Sigma(t)$ IS DESCRIBED BY ${}^{(3)}g_{ij}$ WHICH ENCODES ALL GEOM. INFO. WHICH MAY BE DEDUCED BY MAKING MEASUREMENTS ON $\Sigma(t)$ ALONE

- HOWEVER, A GIVEN 3-GEOMETRY (SLICE, HYPERSURFACE) MAY BE EMBEDDED IN SPACETIME IN INFINITELY MANY DISTINGUISHABLE (BY 4-D MEASUREMENTS) WAYS

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 8 / 29 Find

- THE INTRINSIC GEOMETRY of $\Sigma(t)$ IS DESCRIBED BY $^{(3)}g_{ij}$ WHICH ENCODES ALL GEOM. INFO. WHICH MAY BE DEDUCED BY MAKING MEASUREMENTS ON $\Sigma(t)$ ALONE

- HOWEVER, A GIVEN 3-GEOMETRY (SLICE, HYPERSURFACE) MAY BE EMBEDDED IN SPACETIME IN INFINITELY MANY DISTINGUISHABLE (BY 4-D MEASUREMENTS) WAYS

EXAMPLE: 2D EMBEDDED IN 3D; FLAT SURFACE EMBEDDED WITH / WITHOUT EXTRINSIC CURVATURE

- THE MANNER IN WHICH $\Sigma(t)$ IS EMBEDDED CAN BE CHARACTERIZED BY INVESTIGATING THE CHANGE IN THE

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 8 / 29 Find

EXAMPLE : 2D EMBEDDED IN 3D ; FLAT SURFACE EMBEDDED WITH / WITHOUT EXTRINSIC CURVATURE

- THE MANNER IN WHICH $\Sigma(t)$ IS EMBEDDED CAN BE CHARACTERIZED BY INVESTIGATING THE CHANGE IN THE DIRECTION of THE NORMAL FIELD AS A FCN of POS ON $\Sigma(t)$ - THIS DEFINES THE EXTRINSIC CURVATURE TENSOR (aka THE SECOND FUNDAMENTAL FORM)

$$K_{ij} = -\nabla_i n_j = -\nabla(_i n_{j)} \qquad (7)$$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 8 / 29 Find

• THE MANNER IN WHICH $\Sigma(t)$ IS EMBEDDED CAN BE CHARACTERIZED BY INVESTIGATING THE CHANGE IN THE DIRECTION of THE NORMAL FIELD AS A FCN of POS on $\Sigma(t)$ - THIS DEFINES THE EXTRINSIC CURVATURE TENSOR (aka THE SECOND FUNDAMENTAL FORM)

$$K_{ij} = -\nabla_i n_j = -\nabla(_i n_{j)} \qquad (7)$$

$$\nabla_i n_j = \partial_i n_j - \Gamma^m{}_{ij} n_m$$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 9 / 29 Find

PART 3 CONT THE 3+1 FORMULATION of GR ⑤

$$\nabla_i n_j = \partial_i n_j - \Gamma^m{}_{ij} n_m$$

BUT $n_m = (-\alpha, 0, 0, 0)$ SO

$$\nabla_i n_j = \alpha^{(4)}\Gamma^0{}_{ij}$$

$$= \alpha\left({}^{(4)}g^{0\alpha}\,{}^{(4)}\Gamma_{\alpha ij} + {}^{(4)}g^{\alpha\beta}\,{}^{(4)}\Gamma_{\beta ij}\right)$$

$$\Gamma_{\alpha ij} = \frac{1}{2}\left({}^{(4)}g_{\alpha i,j} + {}^{(4)}g_{\alpha j,i} - {}^{(4)}g_{ij,\alpha}\right)$$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 9 / 29 Find

$$K_{ij} = -\nabla_i n_j = -\nabla(_i n_{j)} \qquad (7)$$

ADM 3+1 THE 3+1 FORMULATION of GR (5)

$$\nabla_i n_j = \partial_i n_j - \Gamma^{\mu}{}_{ij} n_\mu$$

BUT $n_\mu = (-\alpha, 0, 0, 0)$ SO

$$\nabla_i n_j = \alpha^{(4)}\Gamma^0{}_{ij}$$

$$= \alpha\left({}^{(4)}g^{0\alpha}\,{}^{(4)}\Gamma_{\alpha ij} + {}^{(4)}g^{0k}\,{}^{(4)}\Gamma_{kij}\right)$$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 9 / 29 Find

PART 3 (2+1) THE 3+1 FORMULATION of GR

$$\nabla_i n_j = \partial_i n_j - \Gamma^m{}_{ij} n_m$$

BUT $n_m = (-\alpha, 0, 0, 0)$ SO

$$\nabla_i n_j = \alpha^{(4)}\Gamma^0{}_{ij}$$

$$= \alpha\left({}^{(4)}g^{00}\,{}^{(4)}\Gamma_{0ij} + {}^{(4)}g^{0k}\,{}^{(4)}\Gamma_{kij}\right)$$

$$\Gamma_{0ij} = \tfrac{1}{2}\left({}^{(4)}g_{0i,j} + {}^{(4)}g_{0j,i} - {}^{(4)}g_{ij,0}\right)$$

THUS

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 9 / 29 Find

ANY 3+1 THE 3+1 FORMULATION and GR

$$\nabla_i n_j = \partial_i n_j - \Gamma^m{}_{ij} n_m$$

BUT $n_m = (-\alpha, 0, 0, 0)$ SO

$$\nabla_i n_j = \alpha^{(a)}\Gamma^0{}_{ij}$$

$$= \alpha\left({}^{(a)}g^{0\alpha}\,{}^{(a)}\Gamma_{\alpha ij} + {}^{(a)}g^{\alpha\beta}\,{}^{(a)}\Gamma_{\beta ij}\right)$$


Applications Places System
Wed Apr
obe Reader
Tools Window Help
100%
9 / 29
Find


Applications Places System
Wed Apr
obe Reader
Tools Window Help
100%
9 / 29
Find

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 6 / 29 Find

COORDINATES $\{x^i\}$, THUS INTERPRETING $t(\ldots)$ AS A 3-TENSOR $^{(2)}t(\ldots)$, WE NECESSARILY HAVE

$$^{(3)}t_{ij} = {}^{(A)}t_{ij}$$

ON THE OTHER HAND, THE SPATIAL VECTORS $\{{}^{(A)}\underline{\omega}^i\}$ OF THE DUAL BASIS $\{{}^{(A)}\underline{\omega}^m\}$, WILL NOT, IN GENERAL COINCIDE WITH $\{{}^{(3)}\underline{\omega}^i\}$ - THE DUAL BASIS ON $\Sigma(t)$ DEPENDS ON HOW $\Sigma(t)$ IS EMBEDDED IN THE S.T.

THUS, IN GENERAL

$$^{(3)}t_{ij} \neq {}^{(A)}t_{ij}$$

EXAMPLE: CONSIDER THE INVERSE 4-METRIC COMPONENTS

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

I.E. SPATIAL COVARIANT COMPONENTS OF 4- AND 3-METRICS ARE IDENTICAL

- THIS IS A GENERAL RESULT: GIVEN ANY 4-TENSOR OF
- THIS IS A GENERAL RESULT: GIVEN ANY 4-TENSOR OF TYPE (0, k) (COVARIANT TENSOR), THE SPATIAL COMPONENTS OF THAT TENSOR CAN BE IDENTIFIED AS THE COMPONENTS OF A TYPE (0,k) 3-TENSOR

WHY? RECALL THAT COVARIANT TENSOR COMPONENTS CAN BE DEFINED IN TERMS OF THE ACTION OF THE TENSOR ON THE COORDINATE BASIS VECTORS $e_{(\mu)} = e_\mu$, $\mu = 0, 1, 2, 3$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

$$ds^2 = (-\alpha^2 + \beta^i \beta_i)dt^2 + 2\beta_j dx^j dt + {}^{(3)}g_{ij} dx^i dx^j$$

SO, WE HAVE

$$ {}^{(4)}g_{\mu\nu} = \begin{bmatrix} {}^{(4)}g_{00} & {}^{(4)}g_{0j} \\ {}^{(4)}g_{i0} & {}^{(4)}g_{ij} \end{bmatrix} = \begin{bmatrix} -\alpha^2 + \beta^k \beta_k & \beta_j \\ \beta_i & {}^{(3)}g_{ij} \end{bmatrix} \qquad (2)$$

IN PARTICULAR, NOTE THAT

$$ {}^{(4)}g_{ij} = {}^{(3)}g_{ij}$$

I.E. SPATIAL COVARIANT COMPONENTS OF 4- AND 3-METRICS ARE IDENTICAL

THIS IS A GENERAL RESULT; GIVEN ANY 4-TENSOR OF TYPE $(0,k)$ (COVARIANT TENSOR)

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 5 / 29 Find

THIS IS A GENERAL RESULT: GIVEN ANY 1-TENSOR OF TYPE (0, k) (COVARIANT TENSOR), THE SPATIAL COMPONENTS OF THAT TENSOR CAN BE IDENTIFIED AS THE COMPONENTS OF A TYPE (0,k) 3-TENSOR


Applications Places System
Wed Apr
obe Reader
Tools Window Help
100%
5 / 29
Find


Applications Places System
Wed Apr
obe Reader
Tools Window Help
100%
5 / 29
Find


Applications Places System
Wed Apr
obe Reader
Tools Window Help
100%
5 / 29
Find


Applications Places System
Wed Apr
obe Reader
Tools Window Help
100%
5 / 29
Find

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 10 / 29 Find

$$g^{ab}\Omega_a\Omega_b = -\alpha^{-2}\,, \tag{3}$$

where α is the lapse function, as previously. Thus we can construct the unit-norm dual-vector field, n_a, via

$$n_a = -\alpha\Omega_a = -\alpha\nabla_a\tau\,, \tag{4}$$

where the sign is chosen so that the associated unit-norm, hypersurface-orthogonal vector field, n^a,

$$n^a = g^{ab}n_b\,, \tag{5}$$

is future-directed. Note that we can view n^a as the 4-velocity field of a congruence of observers moving orthogonally to the slices (not necessarily coordinate-stationary). Such observers will have a 4-acceleration, a^b given by

$$a^b = n^a\nabla_a n^b\,. \tag{6}$$

2) The Projection Tensor and the Spatial Metric

In the derivation of the 3+1 form of the Einstein equations, we will necessarily be interested in decomposing various spacetime tensors into hypersurface-tangential ("spatial") and hypersurface-orthogonal ("temporal") pieces. Determining the "temporal" part of a tensor is straightforward, we simply contract with n^a. York throws in a slight twist by introducing a relative minus depending on whether a vector or covector index is being projected. Thus, for a vector field, W^a, we define

$$W^{\hat{n}} = -W^a n_a\,, \tag{7}$$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 9 / 29 Find

THE 3+1 EINSTEIN EQUATIONS

These notes rework the calculation of the 3+1 equations as presented in *Kinematics and Dynamics of General Relativity*, by J. W. York, Jr., which itself is contained in the volume *Sources of Gravitational Radiation*, edited by L. Smarr. Many calculational details omitted from that source are included here.

(**Note:** A colleague of mine, who will remain anonymous this week, has used these notes to demonstrate the power of his symbolic manipulation software and has found at least one error

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 9 / 29 Find

$$\nabla_i u_j = \alpha^{(4)}{}_{|0} \, {}^{(4)}_{ij}$$

$$= \alpha \left({}^{(4)}g^{ou} \, {}^{(4)}\Gamma_{oij} + {}^{(4)}g^{ol} \, {}^{(4)}\Gamma_{lij} \right)$$

$$\Gamma_{oij} = \frac{1}{2} \left({}^{(4)}g_{oi,j} + {}^{(4)}g_{oj,i} - {}^{(4)}g_{ij,o} \right)$$

THUS

$$K_{ij} = -\frac{1}{2\alpha} \frac{\partial {}^{(3)}g_{ij}}{\partial t} + \ldots$$

SO WE CAN VIEW THE EXTRINSIC CURVATURE AS THE "VELOCITY" OF THE 3-METRIC ${}^{(3)}g_{ij}$ ("CONJUGATE MOMENTA")


for Atoms
Molecule!

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 9 / 29 Find

$$\nabla_i n_j = \alpha \, {}^{(4)}\Gamma_{ij}$$

$$= \alpha \left({}^{(4)}g^{00}\, {}^{(4)}\Gamma_{0ij} + {}^{(4)}g^{ab}\, {}^{(4)}\Gamma_{aij} \right)$$

$$\Gamma_{0ij} = \frac{1}{2}\left({}^{(4)}g_{0i,j} + {}^{(4)}g_{0j,i} - {}^{(4)}g_{ij,0} \right)$$

THUS

$$K_{ij} = -\frac{1}{2\alpha} \frac{\partial\, {}^{(3)}g_{ij}}{\partial t} + \ldots$$

SO WE CAN VIEW THE EXTRINSIC CURVATURE AS THE "VELOCITY" OF THE 3-METRIC ${}^{(3)}g_{ij}$ ("CONJUGATE MOMENTA")

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 10 / 29 Find

THE 3+1 EINSTEIN EQUATIONS

These notes rework the calculation of the 3+1 equations as presented in *Kinematics and Dynamics of General Relativity*, by J. W. York, Jr., which itself is contained in the volume *Sources of Gravitational Radiation*, edited by L. Smarr. Many calculational details omitted from that source are included here.

(**Note:** A colleague of mine, who will remain anonymous this week, has used these notes to demonstrate the power of his symbolic manipulation software and has found at least one error amongst the many "intermediate" results that are derived below. There is a $20 reward for the PSI student who first identifies such an error and who submits a hand-written explanation of the gaffe, along with the corrected expression, to the tutor.)

1) Foliations and Normals

As before, we consider a spacetime M with metric g_{ab} which is sliced into a foliation $\{\Sigma\}$ defined by the isosurfaces of a scalar field τ (the time parameter). Then the spacelike hypersurfaces are, at least locally, described by a *closed* one-form (dual vector field), Ω_a:

$$\Omega_a = \nabla_a \tau \,. \tag{1}$$

Note that since Ω_a is the gradient of a scalar function, and ∇_a is torsion-free, we have

$$\nabla_{[a}\Omega_{b]} = \nabla_{[a}\nabla_{b]}\tau = 0 \,. \tag{2}$$

The norm of Ω_a is given by

$$g^{ab}\Omega_a\Omega_b = -\alpha^{-2} \,, \tag{3}$$

Applications Places System Wed Apr
obe Reader
Tools Window Help
100% 10 / 29 Find

THE 3+1 EINSTEIN EQUATIONS

These notes rework the calculation of the 3+1 equations as presented in *Kinematics and Dynamics of General Relativity*, by J. W. York, Jr., which itself is contained in the volume *Sources of Gravitational Radiation*, edited by L. Smarr. Many calculational details omitted from that source are included here.

(**Note:** A colleague of mine, who will remain anonymous this week, has used these notes to demonstrate the power of his symbolic manipulation software and has found at least one error amongst the many "intermediate" results that are derived below. There is a $20 reward for the PSI student who first identifies such an error and who submits a hand-written explanation of the gaffe, along with the corrected expression, to the tutor.)

1) Foliations and Normals

As before, we consider a spacetime M with metric g_{ab} which is sliced into a foliation $\{\Sigma\}$ defined by the isosurfaces of a scalar field τ (the time parameter). Then the spacelike hypersurfaces are, at least locally, described by a *closed* one-form (dual vector field), Ω_a:

$$\Omega_a = \nabla_a \tau \,. \tag{1}$$

Note that since Ω_a is the gradient of a scalar function, and ∇_a is torsion-free, we have

$$\nabla_{[a}\Omega_{b]} = \nabla_{[a}\nabla_{b]}\tau = 0 \,. \tag{2}$$

The norm of Ω_a is given by

$$g^{ab}\Omega_a\Omega_b = -\alpha^{-2} \,, \tag{3}$$