Title: Explorations in Numerical Relativity - Lecture 12

Date: Apr 19, 2011 11:30 AM

URL: http://pirsa.org/11040055

Abstract:

perimeter scholars
international

1) Toy model


(x, z)
1) Toy model
$\frac{dx}{dt} = a +$


(x, z)
1) Toy model
$\frac{dx}{dt} = a + (z - z_0)\,x$
$z_0 = z(t=0)$
$\frac{dz}{dt} = (\Pi - z_0)$
$z(t=0) = \Pi$


(x, z)
1) Toy model
dx/dt = a + (z − z₀) x
z₀ = z(t = 0)
dz/dt = (π − z₀)
z(t) = (π − z₀) t
I.C.
z(t = 0) = π
x(t = 0) =
08

(x, z)

1) Toy model

$$\frac{dx}{dt} = \alpha + (z - z_0)\, x$$

$$z_0 = z(t=0) \qquad \frac{dz}{dt} = (\Pi - z_0)$$

$$\Rightarrow z(t) = (\Pi - z_0)\, t + z_0$$

I.C.: $z(t=0) = \Pi$

$x(t=0) = 0$

Soln.: $z(t) = \Pi$

$x(t) = \alpha t$


$\pi - z_0 = \delta$
$\frac{dx}{dt} = a + \delta x$
$\frac{dz}{dt} =$
$z_0) t + z_0$
$= \pi$
$= at$


$\pi - z_0 = \delta$
$\frac{dx}{dt} = a + \delta x$
$\rightarrow x(t) = \frac{a}{\delta}\left[-1 + e^{\delta t}\right]$
$\frac{dz}{dt} = \delta$
$z_0)t + z_0$
$= \pi$
$= at$

$\pi - z_0 = \delta$

$\frac{dx}{dt} = a + \delta x \rightarrow x(t) = \frac{a}{\delta}\left[-1 + e^{\delta t}\right]$

$\frac{dz}{dt} = \delta$

$z_0)t + z_0$

$= \pi$

$= at$



$\pi - z_- = \delta$

$\frac{dx}{dt} = \alpha + \delta x \quad \Rightarrow \quad x(t) = \frac{\alpha}{\delta}\left[-1 + e^{\delta t}\right]$

$+ z_0$ $\quad \frac{dz}{dt} = \delta$

(x, z)

1) Toy model

$$\frac{dx}{dt} = a + (z - z_0)\, x$$

$$\frac{dz}{dt} = (\pi - z_0) - \beta (z - \pi)$$

$z_0 = z(t=0)$

$$\Rightarrow z(t) = (\pi - z_0)\, t + z_0$$

I.C.: $z(t=0) = \pi$

$x(t=0) = 0$

Soln.: $z(t) = \pi$

$x(t) = at$

(E_i, A_i, ϕ)

$\partial_t A_i = -E_i - \partial_i \phi$

$\partial_t E_i = -\partial^j \partial_j A_i + \partial_i \partial^j A_j$


(E_i, A_i, ϕ)
$\partial_t A_i = -E_i - \partial_i \phi$
$\partial_t E_i = -\partial^j \partial_j A_i + \partial_i \partial^j A_j$
$\hookrightarrow \begin{cases} \partial_t^2 A = -\partial_t E \\ \quad \partial^T \partial_j A = \partial_t \end{cases}$

(E_i, A_i, ϕ) $\partial_t A_i = -E_i - \partial_i \phi$

$\partial_t E_i = -\partial^j \partial_j A_i + \partial_i \partial^j A_j$

$\hookrightarrow \begin{cases} \partial_t^2 A = -\partial_t E \\ \partial_t^2 A_i = \partial^j \partial_j A_i - \partial_i \partial^j A_j \end{cases}$


(E_i, A_i, ϕ)
$\partial_t A_i = -E_i - \partial_i \phi$
$\partial_t E_i = -\partial^j \partial_j A_i + \partial_i \partial^j A_j$
$\hookrightarrow \begin{cases} \partial_t^2 A = -\partial_t E \\ \partial_t^2 A_i = \partial^j \partial_j A_i - \partial_i \partial^j A_j \end{cases}$


(E_i, A_i, ϕ)
$\partial_t A_i = -E_i - \partial_i \phi$
$\partial_t E_i = -\partial^j \partial_j A_i + \partial_i \partial^j A_j$
$\hookrightarrow \begin{cases} \partial_t^2 A = -\partial_t E \\ \partial_t^2 A_i = \partial^j \partial_j A_i - \partial_i \partial^j A_j \end{cases}$

Assume $\rho = 0$; 2D

$\partial_x A_1 = a$ $\quad \partial_y A_1 = \alpha$

$\partial_x A_2 = \beta$ $\quad \partial_y A_2 = b$

Assume $\phi = 0$; 2D

$\partial_x A_1 = a$ $\quad \partial_y A_1 = \alpha$

$\partial_x A_2 = \beta$ $\quad \partial_y A_2 = b$

$\partial_t a = \partial_t \partial_x A_1$

$= \partial_x \partial_t A_1 = -$

Assume $\rho = 0$; 2D

$\partial_x A_1 = a$ $\partial_y A_1 = \alpha$

$\partial_x A_2 = \beta$ $\partial_y A_2 = b$

$\partial_t a = \partial_t \partial_x A_1$

$= \partial_x \partial_t A_1 = -\partial_x E_1$

$\partial_t \alpha = \partial_t \partial_y A_1$

$= \partial_y \partial_t A_1 = -\partial_y E_1$

$\partial_t \beta = \partial_x \partial_t A_2 = -\partial_x E_2$

$\partial_t \beta = -\partial_y \partial_t A_1 = -\partial_y E_2$

Assume $\rho = 0$; 2D

$\partial_x A_1 = a$ $\partial_y A_1 = \alpha$

$\partial_x A_2 = \beta$ $\partial_y A_2 = b$

$\partial_t A_x = -E_1$

$\partial_t E_1 = -$

$\partial_t a = \partial_t \partial_x A_1$

$= \partial_x \partial_t A_1 = -\partial_x E_1$

$\partial_t \alpha = \partial_t \partial_y A_1$

$= \partial_y \partial_t A_1 = -\partial_y E_1$

$\partial_t \beta = \partial_x \partial_t A_2 = -\partial_x E_2$

$\partial_t \beta = -\partial_y \partial_t A_2 = -\partial_y E_2$


Assume β = 0. ; 2D
∂x A1 = α
∂y A1 = α
∂x A2 = β
∂y A2 = b
∂t A1 = −E1
E1 = − ∂x a − ∂y α + ∂x (a + b)
E2 = −
∂t a = ∂t ∂x A1
= ∂x ∂t A1 = − ∂x
∂t α = ∂t ∂y A1
= ∂y ∂t A1 = −∂y E
∂t β = ∂x ∂t A2 = −∂x E
∂t β = − ∂y ∂t A2 = −∂y

Assume $\beta = 0$; 2D

$\partial_x A_1 = a$ $\quad \partial_y A_1 = \alpha$

$\partial_x A_2 = \beta$ $\quad \partial_y A_2 = b$

$\partial_t A_x = -E_i$

$\partial_t E_1 = -\partial_x a - \partial_y \alpha + \partial_x (a+b)$

$\partial_t E_2 = -\partial_x \beta - \partial_y b + \partial_y (a+b)$

$\partial_t a = \partial_t \partial_x A_1$

$= \partial_x \partial_t A_1 = -\partial_x$

$\partial_t \alpha = \partial_t \partial_y A_1$

$= \partial_y \partial_t A_1 = -\partial_y E$

$\partial_t \beta = \partial_x \partial_t A_2 = -\partial_x E$

$\partial_t \beta = -\partial_y \partial_t A_2 = -\partial_y$


$\partial_t E_1 = -\partial_x a + \partial_x$
∂_t
∂_t
∂_t


$\partial_t E_1 = -\partial_x a + \partial_x a + \partial_x b$
$\partial_t E_2 =$
08


$\partial_t E_1 = -\partial_x \alpha + \partial_x a + \partial_x b$
$\partial_t E_2 = -\partial_x \beta$
08

$$\partial_t E_1 = -\partial_x a + \partial_x a + \partial_x b$$

$$\partial_t E_2 = -\partial_x \beta$$

$$\partial_t \beta = -\partial_x E_2$$

$$\partial_t b = -\partial_x E_2$$


$\partial_t E_1 = -\partial_x a + \partial_x a + \partial_x b$
$\partial_t E_2 = -\partial_x \beta$
$\partial_t \beta = -\partial_x E_2$
$\partial_t b = 0$
E_1
E_2
β
b
$/t$
$=$
E_1
E_2
β
b
$/x$


$\partial_t E_1 = -\partial_x a + \partial_x a + \partial_x b$
$\partial_t E_2 = -\partial_x \beta$
$\partial_t \beta = -\partial_x E_2$
$\partial_t b = 0$
$= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & & \end{pmatrix} \begin{pmatrix} E_1 \\ E_2 \\ \beta \\ b \end{pmatrix}$


$\partial_t E_1 = -\partial_x \alpha + \partial_x \alpha + \partial_x b$
$\partial_t E_2 = -\partial_x \beta$
$\partial_t \beta = -\partial_x E_2$
$\partial_t b = 0$
$\begin{pmatrix} E_1 \\ E_2 \\ \beta \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} E_1 \\ E_2 \\ \beta \\ b \end{pmatrix}$
08



Pirsa: 11040055

Assume $\rho = 0$; 2D

$\partial_x A_1 = a$ $\partial_y A_1 = \alpha$

$\partial_x A_2 = \beta$ $\partial_y A_2 = b$

$\partial_t A_i = -E_i$

$\partial_t E_1 = -\partial_x a - \partial_y \alpha + \partial_x (a + b)$

$\partial_t E_2 = -\partial_x \beta - \partial_y b + \partial_y (a + b)$

$\partial_t E_i = -\partial^j \partial_j A_i + \partial^i \partial_j A^j$

$\partial_t a = \partial_t \partial_x A_1$

$= \partial_x \partial_t A_1 = -\partial$

$\partial_t \alpha = \partial_t \partial_y A_1$

$= \partial_y \partial_t A_1 = -\partial_y$

$\partial_t \beta = -\partial_x \partial_t A_2 = -\partial_x$

$\partial_t b = -\partial_y \partial_t A_2 = -\partial$


$\partial_t E_1 = -\partial_x a + \partial_x a + \partial_x b$
$\partial_t E_2 = -\partial_x \beta$
$\partial_t \beta = -\partial_x E_2$
$\partial_t b = +$
$\partial_j E^j = 0$
$\begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$
$\begin{pmatrix} E_1 \\ E_2 \\ \beta \\ b \end{pmatrix}$


$\partial_j E^j = 0$
$\partial_t E_1 = -\partial_x \alpha + \partial_x a + \partial_x b$
$\partial_t E_2 = -\partial_x \beta$
$\partial_t \beta = -\partial_x E_2$
$\partial_t b = +k(\partial_j E^j)$
$\begin{pmatrix} E_1 \\ E_2 \\ \beta \\ b \end{pmatrix}_t = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} E_1 \\ E_2 \\ \beta \\ b \end{pmatrix}_x$


$\partial_j E^j = 0$
$\partial_t E_1 = -\partial_x a + \partial_x a + \partial_x b$
$\partial_t E_2 = -\partial_x \beta$
$\partial_t \beta = -\partial_x E_2$
$\partial_t b = +k\, \partial_x E_1$
$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} E_1 \\ E_2 \\ \beta \\ b \end{pmatrix}$


$\partial_t E_1 = -\partial_x \alpha + \partial_x a + \partial_x b$
$\partial_t E_2 = -\partial_x \beta$
$\partial_t \beta = -\partial_x E_2$
$\partial_t b = +k\, \partial_x E_1$
$\partial_j E^j = 0$
$\begin{pmatrix} E_1 \\ E_2 \\ \beta \\ \end{pmatrix}_{/t} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ k & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} E_1 \\ E_2 \\ \beta \\ b \end{pmatrix}_{/x}$
08

$$\partial_j E^j = 0$$

$$\partial_t E_1 = -\partial_x \alpha + \partial_x a + \partial_x b$$

$$\partial_t E_2 = -\partial_x \beta$$

$$\partial_t \beta = -\partial_x E_2$$

$$\partial_t b = +k\, \partial_x E_1$$

$$\begin{pmatrix} E_1 \\ E_2 \\ \beta \\ b \end{pmatrix}_{lt} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ k & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} E_1 \\ E_2 \\ \beta \\ b \end{pmatrix}_{lt}$$


Grains of
Pollen to
Evidence
for Atoms
How
Big Is A
Molecule?
Hyperbolic


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Hyperbolic
weakly
strongly
Symmetric


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Example
$\dot{\pi}_{t+\epsilon} = \pi_{,x} + P_{,x}$
$P_{,t} = P_{,x}$
$\Rightarrow \begin{pmatrix} \pi \\ P \end{pmatrix}_{,t} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \pi \\ P \end{pmatrix}_{,x}$
Hyperbolic { weakly strongly Symmetric }


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Hyperbolic
weakly
strongly
symmetric
Fourier expansion


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Example
Hypebolic
weakly
strongly
Symmetric
Fourier expansion


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Example
Hypebolic
weakly
strongly
Symmetric
Fourier expansion


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Example
Hyperbolic
weakly
strongly
Symmetric
Fourier expansion


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Example
Hyperbolic
weakly
strongly
symmetric
Fourier expansion


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Example
Hyperbolic
weakly
strongly
symmetric
expansion
08


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Example
Hyperbolic
weakly
strongly
Symmetric
Fourier expansion


Grains of Pollen to Evidence for Atoms
How Big Is A Molecule?
Example
Hyperbolic
weakly
strongly
Symmetric
Fourier expansion


ADM: $\{\gamma_{ij}, K_{ij}, \alpha, \beta^i\}$
$\beta^i = 0$
$\partial_t^2 \gamma_{ij} = \alpha^2 \gamma^{kl} \gamma^{mn} (\partial_k \partial_m \gamma_{ij} + \partial_i \partial_j \gamma_{kl} - \partial$


ADM: $\{\gamma_{ij}, K_{ij}; \alpha, \beta^i\}$
$\beta^i \in 0$
$\partial_t^2 \gamma_{ij} = \alpha^2 \gamma^{lm}\left(\partial_l \partial_m \gamma_{ij} + \partial_i \partial_j \gamma_{lm} - \partial_i \partial_l \gamma_{mj} - \partial_j \partial_l \gamma_{mi}\right) + 2\alpha \partial_i \partial_j \alpha +$

ADM: $\{\gamma_{ij}; K_{ij}; \alpha, \beta^i\}$

$\beta^i = 0$

$$\partial_t^2 \gamma_{ij} = \alpha^2 \gamma^{lm}\left(\partial_l \partial_m \gamma_{ij} + \partial_i \partial_j \gamma_{lm} - \partial_i \partial_l \gamma_{mj} - \partial_j \partial_l \gamma_{mi}\right) + 2\alpha \partial_i \partial_j \alpha + \ldots$$




$\partial_t^2 A_i = - \partial^j \partial_j A_i + \partial_i \partial^j A_j$
$\partial_i \partial_k \gamma_{mj} - \partial_j \partial_k \gamma_{mi}) + 2 \alpha \partial_i \partial_j \alpha + \ldots$


ADM: $\{\gamma_{ij}; K_{ij}; \alpha, \beta^i\}$
Constraints
$\beta^i = 0$
$\partial_t^2 \gamma_{ij} = \alpha^2 \gamma^{kl} \gamma^{mn} \left(\partial_k \partial_l \gamma_{jn} + \partial_i \partial_j \gamma \right) - \partial_i \partial_k \gamma_{jn}$

Constraints

1) $\gamma^{ir}\gamma^{lm}(2\,\partial_e\gamma_{jm} - \partial_j\gamma_{ie}) + \text{l.o.} = 0$

$\gamma_{ir} + 2\,\partial_j\gamma_{lm} - 2\,\partial_e\gamma_{mj} - \partial_j\partial_e\gamma_{mi}) +$


for Atoms
Molecule?


for Atoms
Molecule?
08


for Atoms
Molecule?
$h_{ij} = \delta_{ij} + \epsilon\, h_{ij}$
$\gamma^{ik} = \delta^{ik} - \epsilon\, \delta^{ip}\delta^{kq} h_{pq} + \mathcal{O}(\epsilon^2)$
$tr\, h = \mathrm{trace}\,(h_{ij})$


for Atoms
Molecule?
$g_{ij} = \delta_{ij} + \varepsilon h_{ij}$
$\gamma^{ik} = \delta^{ik} - \varepsilon \delta^{ip} \delta^{kq} h_{pq} + \mathcal{O}(\varepsilon^2)$
$\partial_t (\partial_{t+1} - \delta$
$+1 = \text{trace}(h_{ij})$
$= \delta^{ij} h_{ij}$


for Atoms
Molecule?
constraints
$h_{ij} = \delta_{ij} + \varepsilon\, h_{ij}$
$\delta^{ik}\delta^{lm} = \delta^{ik}\delta^{lm} - \varepsilon\,\delta^{ip}\delta^{mq}h_{mq} + \mathcal{O}(\varepsilon^2)$
$\partial_t(\partial_i H - \delta^{lm}\partial_l h_{im})$
$H = \mathrm{trace}(h_{ij})$
$= \delta^{ij} h_{ij}$
$\delta^{ij}\delta^{lm}\partial_i\partial_l h$
$= 0$


for Atoms
Molecule?
$\gamma^{ij} = \delta^{ij} - \varepsilon\, h^{ij} + \varepsilon^2 h^{ik} h_k{}^j$
$\gamma^{ij} = \delta^{ij} - \varepsilon\, \delta^{ip}\delta^{jq} h_{pq} + \mathcal{O}(\varepsilon^2)$
constraints
$\partial_t(\partial_i H - \delta^{lm}\partial_l h_{im}) = 0$
$\gamma^{ij}\gamma^{lm}\partial_i\partial_l h_{jm} - \Delta H = 0 \quad (\Delta = \delta^{ij}\partial_i\partial_j)$


for Atoms
Molecule?
$g_{ij} = \delta_{ij} + \varepsilon h_{ij}$
$g^{ij} = \delta^{ij} - \varepsilon \delta^{ip} \delta^{jq} h_{pq} + O(\varepsilon^2)$
straints
$\partial_t (\partial_i H - \delta^{lm} \partial_l h_{im}) = 0$
$H = \text{trace}(h_{ij}) = \delta^{ij} h_{ij}$
$g^{ij} g^{lm} \partial_i \partial_l h_{jm} - \Delta H = 0 \quad (\Delta = \partial^i \partial_i)$

Constraints

$\partial_t^2 A_i = -\partial^J \partial_J A_i + \partial_i \partial^J A_J$

$H_{(m)}$) $\gamma^{ij}\gamma^{kl}(\partial_i \partial_k \gamma_{jlm} - \partial_j \gamma_{ik}) + l.o.t. = 0$

$H_{(m)}$) $\gamma^{jk}\partial_t(\partial_i \gamma_{jk} - \partial_j \gamma_{ik}) + l.o.t. = 0$

$\partial_m \gamma_{ij} + \partial_i \partial_j \gamma_{klm} - \partial_i \partial_k \gamma_{mj} - \partial_j \partial_k \gamma_{mi}) + 2\alpha \partial_i \partial_j \alpha +$ lower order

$\{\partial_k \gamma_{ij}, \gamma_{ij}, \partial_i$


ADM: $\{\gamma_{ij}, K_{ij}, \alpha, \beta^i\}$
Constraints
$\beta^i = 0$
$\partial_t^2 \gamma_{ij} = \alpha^2 \gamma^{lm}(\partial_l \partial_m \gamma_{ij} + \partial_i \partial_j \gamma_{lm} - \partial_i \partial_l \gamma_{mj} - \partial_j \partial_l \gamma_{mi}) + 2\alpha \partial_i \partial_j \alpha +$ lower
① $\alpha = 1 \Rightarrow \partial_t h_{ij} = \Delta h_{ij} - 2\partial_i \partial_j H$
where we use that $H_{ADM} + M_{ADM} \Rightarrow$


ADM: $\{\gamma_{ij}; K_{ij}; \alpha, \beta^i\}$
Constraints
$\beta^i = 0$
$\partial_t^2 \gamma_{ij} = \alpha^2 \gamma^{lm}(\partial_l \partial_m \gamma_{ij} + \partial_i \partial_j \gamma_{lm} - \partial_i \partial_l \gamma_{mj} - \partial_j \partial_l$
① $\alpha = 1 \Rightarrow \partial_t^2 h_{ij} = \Delta h_{ij} - \partial_i \partial_j H$
② $\alpha = \frac{1}{3} \delta^{lm} h_{lm} \Rightarrow \partial_t^2$


ADM: $\{\gamma_{ij}; K_{ij}; \alpha, \beta^i\}$
Constraints
$\beta^i = 0$
$\partial_t^2 \gamma_{ij} = \alpha^2 \gamma^{lm}(\partial_l \partial_m \gamma_{ij} + \partial_i \partial_j \gamma_{lm} - \partial_i \partial_l \gamma_{mj} - \partial_j \partial_l$
① $\alpha = 1 \Rightarrow \partial_t^2 h_{ij} = \Delta h_{ij}$
② $\alpha = \frac{1}{3} \delta^{lm} h_{lm} \Rightarrow$
where we use

ADM: $\{\gamma_{ij}; K_{ij}; \alpha, \beta^i\}$ Constraints

$\beta^i = 0$.

$H_{(m)}$) $\gamma^{ij}\gamma^{lm}(\partial_i \partial_l \gamma_{jm} - \partial_j \gamma_{il}) +$

$M_{(m)}$) $\gamma^{ij}\partial_t(\partial_i \gamma_{jl} - \partial_j \gamma_{il}) + 1 \cdot 0 =$

$\rightarrow \partial_t^2 \gamma_{ij} = \alpha^2 \gamma^{lm}\left(\partial_l \partial_m \gamma_{ij} + \partial_i \partial_j \gamma_{lm} - \partial_i \partial_l \gamma_{mj} - \partial_j \partial_l\right.$

① $\alpha = 1 \Rightarrow \partial_t^2 h_{ij} = \Delta h_{ij} - \partial_i \partial_j H$ where we use the

② $\alpha = \frac{1}{3}\delta^{lm} h_{lm} \Rightarrow \partial_t^2 h_{ij} = \left(\Delta h_{ij} - \frac{1}{3}\partial_i \partial_j H\right)$

ADM: $\{\gamma_{ij}; K_{ij}; \alpha, \beta^i\}$ Constraints

$\beta^i \in 0$.

$H_{(m)}$) $\gamma^{ij}\gamma^{lm}(\partial_i \partial_e \gamma_{jm} - \partial_j \gamma_{ie}) +$

$M_{(m)}$) $\gamma^{jl}\partial_t(\partial_i \gamma_{jl} - \partial_j \gamma_{il}) + 1 \circ =$

$\rightarrow \partial_t^2 \gamma_{ij} = \alpha^2 \gamma^{lm}\left(\partial_e \partial_m \gamma_{ij} + \partial_i \partial_j \gamma_{lm} - \partial_i \partial_e \gamma_{mj} - \partial_j \partial_e\right.$

① $\alpha = 1 \Rightarrow \partial_t^2 h_{ij} = \left(\Delta h_{ij} - \partial_i \partial_j H\right)$ where we use the

② $\alpha = \frac{1}{3}\delta^{lm} h_{lm} \Rightarrow \partial_t^2 h_{ij} = \left(\Delta h_{ij} - \frac{1}{3}\partial_i \partial_j H\right)$


Evidence for Atoms
Big Is A Molecule?
$R_{ab} = -\frac{1}{2} g^{\alpha\beta}$


Evidence for Atoms
Big Is A Molecule?
08


Evidence for Atoms
Big Is A Molecule?
$R_{ab} = -\frac{1}{2} g^{\alpha\beta} \partial_\alpha \partial_\beta g_{ab} + \nabla_{(a}\Gamma_{b)} + g^{\delta\delta} g^{\varepsilon f} (\partial_\varepsilon g_{\gamma a} \partial_f g_{\delta b}$
$- \Gamma_{a\gamma\varepsilon} \Gamma$


Evidence for Atoms
Big Is A Molecule?


Evidence for Atoms
Big Is A Molecule?
hyperbolic


$\partial_t^2 A_i - \partial^j \partial_j A_i + \partial_i \partial^j A_j$
$\partial_t (\partial_i \gamma_{jk} - \partial_j \gamma_{ik}) + \dots = 0$
$\partial_i \partial_j \gamma_{km} - \partial_i \partial_k \gamma_{mj} - \partial_j \partial_k \gamma_{mi}) + 2\alpha \partial_i \partial_j \alpha +$ lower order terms
$\{\partial_t \gamma, \gamma, \partial_i \gamma\}$
$H_{ADM} + M_{SW} \Rightarrow \partial_i H - \delta^{kl} \partial_k h_{im} = 0$


Evidence
for Atoms
Big Is A
Molecule?
$R_{ab} = \partial_\alpha \Gamma^\alpha_{ab} - \partial_b \Gamma^\alpha_{a\alpha} + \Gamma^\alpha_{\beta\alpha}\Gamma^\beta_{ab} - \Gamma^\alpha_{\beta b}\Gamma^\beta_{a\alpha}$
$= \partial_\alpha \partial_\beta g_{ab} + \nabla_{(a}\Gamma_{b)} + g^{\gamma\delta} g^{\varepsilon f}(\partial_\varepsilon g_{\gamma a}\partial_f g_{\delta b}$
$- \Gamma_{a\gamma\varepsilon}\Gamma_{b\delta f}$
(de Donder)
symmetric
hyperbolic
$\Gamma_b \rightarrow$ 4 pieces
08


Evidence for Atoms
Big Is A Molecule?
$R_{ab} = -\frac{1}{2} g^{\alpha\beta} \partial_\alpha \partial_\beta g_{ab} + \nabla_{(a}\Gamma_{b)} + g^{\gamma\delta} g^{\varepsilon f} (\partial_\varepsilon g_{\gamma a} \partial_f g_{\delta b} - \Gamma_{a\gamma\varepsilon}\Gamma_{b\delta f})$
(de Donder)
symmetric hyperbolic
$\Gamma_b \to$ 4 piece
$\Box x^i_b = \Gamma^i_b$


Evidence
for Atoms
Big Is A
Molecule?
(de Donder)
symmetric
hyperbolic
"Harmonic


Evidence for Atoms
Big La A Molecule!
(de Donder)
$\Gamma_b \to$ 4 pieces
"Harmonic Gradation"
$\Gamma_b = 0$


Evidence for Atoms
Big Is A Molecule!
$R_{ab} = -\frac{1}{2} g^{\alpha\beta} \partial_\alpha \partial_\beta g_{ab} + \nabla_{(a}\Gamma_{b)} + g^{\alpha\beta} g^{\varepsilon f} (\partial_\varepsilon g_{\gamma\alpha} \partial_f g_{\delta\beta} - \Gamma_{a\gamma\varepsilon}\Gamma_{b\delta f})$
(de Donder)
symmetric hyperbolic
Γ_b -> 4 pieces
$0 = \Box x^i = \Gamma^i$
"Harmonic Gravity"
$\Gamma_b = 0$


Evidence for Atoms
Big Is A Molecule?
$R_{ab} = -\frac{1}{2} g^{\alpha\beta} \partial_\alpha \partial_\beta g_{ab} + \nabla_{(a}\Gamma_{b)} + g^{\delta\delta} g^{\varepsilon f} (\partial_\varepsilon g_{\delta a} \partial_f g_{\delta b}$
symmetric hyperbolic
$\Gamma_b \to$ 4 pieces
$\mathcal{H}_b = \Box x_b = \Gamma_b$
"Harmonic
$\Gamma_b = 0$


evidence
for Atoms
Big Is A
Molecule?
08


Conditions coordinates
$\Rightarrow D X^{\mu}_{\sigma} = \Gamma^{\mu}_{\nu}$
$\Gamma^{\mu}_{\nu} = 0$
$R_{\sigma\nu} = \nabla_{(\sigma} \Gamma_{\nu)}$
where


Conditions coordinates
$\Rightarrow D X^{\mu}_{b} = \Gamma^{a}_{b}$
$\Gamma^{a}_{b} = 0 \Rightarrow C_b = 0$
$R_{ab} = \nabla_{(a} C_{b)}$
where


Conditions coordinates
$R_{ab} = \nabla_{(a} C_{b)}$
Bianchi identities. $(\nabla^a G_{ab} = 0)$
$\nabla^a R_{ab} = \frac{1}{2} \nabla_b R$
take
where

Conditions coordinates $\Rightarrow \Box X^{\mu}_{b} = \Gamma^{\mu}_{b}$

$\Gamma^{\mu}_{b} = 0 \Rightarrow C_b = 0$

$* R_{ab} = \nabla_{(a} C_{b)}$

$\rightarrow$ Bianchi identities. $(\nabla^a G_{ab} = 0)$

$\nabla^a R_{ab} = \frac{1}{2} \nabla_b R$

$\rightarrow$ take $\nabla^a (* \quad)$


Conditions coordinates
Bianchi identities
where


coordinates
$\Rightarrow D X^{\mu}_{b} = \Gamma^{a}_{b}$
$\Gamma^{a}_{b} = 0 \Rightarrow C_b = 0$
$(D^a G_{ab} = 0)$
$\nabla_b R$
$\nabla^b \nabla_b C_{at} + C^b \nabla_a C_b = 0$
$\partial_t^2 A_i = -\partial^j$
$(\partial_i \partial_e \gamma_{jm} - \partial_j \gamma_{ie}) + 1.0. = 0$
$\gamma_{je} - \partial_j \gamma_{ie}) + 1\,0. = 0$
$-\partial_i \partial_e \gamma_{mj} - \partial_j \partial_e \gamma_{mi}) + 2\alpha\, \partial_i \partial_j \alpha$
where we use that $H_{ADM} + M_{gauge} \Rightarrow$
to show


$\partial_t^2 A_i = -\partial^j \partial_j A_i + \partial_i \partial^j A_j$
$(\partial_i \partial_e \gamma_{jm} - \partial_j \gamma_{ie}) + l.o. = 0$
$\gamma_{je} - \partial_j \gamma_{ie}) + l.o. = 0$
$-\partial_i \partial_e \gamma_{mj} - \partial_j \partial_e \gamma_{mi}) + \partial_i \partial_j \alpha +$ lower order terms
$\{\partial_e \gamma, \gamma, \partial_i \alpha\}$
where we use that $H_{ADM} + M_{gauge} \Rightarrow \partial_i H - \delta^{jm} \partial_e h_{im} = 0$
$D^b D_b C_a + C^b D_{(a} C_{b)} = 0$
$\Box C_a - C^2$