Title: Looking for Cosmic Strings in New Observational Windows

Date: Aug 10, 2011 02:00 PM

URL: http://pirsa.org/11080020

Abstract: Cosmic strings are predicted to arise in both inflationary and non-inflationary cosmological models. The signatures of such strings will stand out particularly well at higher redshifts. I will discuss how to look for these signatures in CMB redshift and polarization maps and in 21cm redshift surveys.

Signatures of Cosmic Strings in New Observational Windows

Robert Brandenberger
McGill University

August 10, 2011

Outline

Outline

Cosmic Strings

R. Brandenberger

Plan

1. Introduction
2. Cosmic String Review
3. Kaiser-Stebbins Effect and Cosmic String Wakes
4. Signatures of Cosmic Strings in CMB Polarization
5. Signatures of Cosmic Strings in 21cm Maps
6. Conclusions

Cosmic Strings

T. Kibble, J. Phys. A **9**, 1387 (1976); Y. B. Zeldovich, Mon. Not. Roy. Astron. Soc. **192**, 663 (1980); A. Vilenkin, Phys. Rev. Lett. **46**, 1169 (1981).

- Cosmic string = linear topological defect in a quantum field theory.
- 1st analog: line defect in a crystal
- 2nd analog: vortex line in superfluid or superconductor
- Cosmic string = line of trapped energy density in a quantum field theory.
- Trapped energy density → gravitational effects on space-time → important in cosmology.

Cosmic Strings

T. Kibble, J. Phys. A **9**, 1387 (1976); Y. B. Zeldovich, Mon. Not. Roy. Astron. Soc. **192**, 663 (1980); A. Vilenkin, Phys. Rev. Lett. **46**, 1169 (1981).

- Cosmic string = linear topological defect in a quantum field theory.
- 1st analog: line defect in a crystal
- 2nd analog: vortex line in superfluid or superconductor
- Cosmic string = line of trapped energy density in a quantum field theory.
- Trapped energy density → gravitational effects on space-time → important in cosmology.

Cosmic Strings

T. Kibble, J. Phys. A **9**, 1387 (1976); Y. B. Zeldovich, Mon. Not. Roy. Astron. Soc. **192**, 663 (1980); A. Vilenkin, Phys. Rev. Lett. **46**, 1169 (1981).

- Cosmic string = linear topological defect in a quantum field theory.
- 1st analog: line defect in a crystal
- 2nd analog: vortex line in superfluid or superconductor
- Cosmic string = line of trapped energy density in a quantum field theory.
- Trapped energy density → gravitational effects on space-time → important in cosmology.

Cosmic Strings

T. Kibble, J. Phys. A **9**, 1387 (1976); Y. B. Zeldovich, Mon. Not. Roy. Astron. Soc. **192**, 663 (1980); A. Vilenkin, Phys. Rev. Lett. **46**, 1169 (1981).

- Cosmic string = linear topological defect in a quantum field theory.
- 1st analog: line defect in a crystal
- 2nd analog: vortex line in superfluid or superconductor
- Cosmic string = line of trapped energy density in a quantum field theory.
- Trapped energy density → gravitational effects on space-time → important in cosmology.

Relevance to Particle Physics and Cosmology I

- Cosmic strings are predicted in many particle physics models beyond the “Standard Model".
- Cosmic strings are predicted to form at the end of inflation in many inflationary models.
- Cosmic strings may survive as cosmic superstrings in alternatives to inflation such as string gas cosmology.
- In models which admit cosmic strings, cosmic strings inevitably form in the early universe and persist to the present time.

Relevance to Particle Physics and Cosmology I

Cosmic Strings

R. Brandenberger

- Introduction
- Cosmic String Review
- Kaiser-Stebbins Effect and Cosmic String Wakes
- Signatures of Cosmic Strings in CMB Polarization
- Signatures of Cosmic Strings in 21cm Maps
- Conclusions

- Cosmic strings are predicted in many particle physics models beyond the “Standard Model".
- Cosmic strings are predicted to form at the end of inflation in many inflationary models.
- Cosmic strings may survive as cosmic superstrings in alternatives to inflation such as string gas cosmology.
- In models which admit cosmic strings, cosmic strings inevitably form in the early universe and persist to the present time.

Relevance to Particle Physics and Cosmology I

Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions

- Cosmic strings are predicted in many particle physics models beyond the “Standard Model".
- Cosmic strings are predicted to form at the end of inflation in many inflationary models.
- Cosmic strings may survive as cosmic superstrings in alternatives to inflation such as string gas cosmology.
- In models which admit cosmic strings, cosmic strings inevitably form in the early universe and persist to the present time.

Relevance to Particle Physics and Cosmology I

- Cosmic strings are predicted in many particle physics models beyond the "Standard Model".
- Cosmic strings are predicted to form at the end of inflation in many inflationary models.
- Cosmic strings may survive as cosmic superstrings in alternatives to inflation such as string gas cosmology.
- In models which admit cosmic strings, cosmic strings inevitably form in the early universe and persist to the present time.

Relevance to Particle Physics and Cosmology II

Cosmic Strings

R. Brandenberger

- Cosmic strings are characterized by their tension μ which is associated with the energy scale η at which the strings form ($\mu \sim \eta^2$).
- Searching for the signatures of cosmic strings is a tool to probe physics beyond the Standard Model at energy ranges complementary to those probed by the LHC.
- Cosmic strings are constrained from cosmology: strings with a tension which exceed the value $G\mu \sim 3 \times 10^{-7}$ are in conflict with the observed acoustic oscillations in the CMB angular power spectrum.
- Existing upper bound on the string tension rules out large classes of particle physics models.

It is interesting to find ways to possibly lower the bounds on the string tension.

Relevance to Particle Physics and Cosmology II

- Cosmic strings are characterized by their tension μ which is associated with the energy scale η at which the strings form ($\mu \sim \eta^2$).
- Searching for the signatures of cosmic strings is a tool to probe physics beyond the Standard Model at energy ranges complementary to those probed by the LHC.
- Cosmic strings are constrained from cosmology: strings with a tension which exceed the value $G\mu \sim 3 \times 10^{-7}$ are in conflict with the observed acoustic oscillations in the CMB angular power spectrum.
- Existing upper bound on the string tension rules out large classes of particle physics models.

It is interesting to find ways to possibly lower the bounds on the string tension.

Relevance to Particle Physics and Cosmology II

- Cosmic strings are characterized by their tension μ which is associated with the energy scale η at which the strings form ($\mu \sim \eta^2$).
- Searching for the signatures of cosmic strings is a tool to probe physics beyond the Standard Model at energy ranges complementary to those probed by the LHC.
- Cosmic strings are constrained from cosmology: strings with a tension which exceed the value $G\mu \sim 3 \times 10^{-7}$ are in conflict with the observed acoustic oscillations in the CMB angular power spectrum.
- Existing upper bound on the string tension rules out large classes of particle physics models.

It is interesting to find ways to possibly lower the bounds on the string tension.

Relevance to Particle Physics and Cosmology II

- Cosmic strings are characterized by their tension μ which is associated with the energy scale η at which the strings form ($\mu \sim \eta^2$).
- Searching for the signatures of cosmic strings is a tool to probe physics beyond the Standard Model at energy ranges complementary to those probed by the LHC.
- Cosmic strings are constrained from cosmology: strings with a tension which exceed the value $G\mu \sim 3 \times 10^{-7}$ are in conflict with the observed acoustic oscillations in the CMB angular power spectrum.
- Existing upper bound on the string tension rules out large classes of particle physics models.

It is interesting to find ways to possibly lower the bounds on the string tension.

Relevance to Particle Physics and Cosmology II

- Cosmic strings are characterized by their tension μ which is associated with the energy scale η at which the strings form ($\mu \sim \eta^2$).
- Searching for the signatures of cosmic strings is a tool to probe physics beyond the Standard Model at energy ranges complementary to those probed by the LHC.
- Cosmic strings are constrained from cosmology: strings with a tension which exceed the value $G\mu \sim 3 \times 10^{-7}$ are in conflict with the observed acoustic oscillations in the CMB angular power spectrum.
- Existing upper bound on the string tension rules out large classes of particle physics models.

It is interesting to find ways to possibly lower the bounds on the string tension.

Relevance to Particle Physics and Cosmology III

Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions

Cosmic strings can produce many good things for cosmology:

- String-induced mechanism of baryogenesis (R.B., A-C. Davis and M. Hindmarsh, 1991).
- Explanation for the origin of primordial magnetic fields which are coherent on galactic scales (X.Zhang and R.B. (1999).
- Explanation for cosmic ray anomalies (R.B., Y. Cai, W. Xue and X. Zhang (2009).

It is interesting to find evidence for the possible existence of cosmic strings.

Relevance to Particle Physics and Cosmology II

- Cosmic strings are characterized by their tension μ which is associated with the energy scale η at which the strings form ($\mu \sim \eta^2$).
- Searching for the signatures of cosmic strings is a tool to probe physics beyond the Standard Model at energy ranges complementary to those probed by the LHC.
- Cosmic strings are constrained from cosmology: strings with a tension which exceed the value $G\mu \sim 3 \times 10^{-7}$ are in conflict with the observed acoustic oscillations in the CMB angular power spectrum.
- Existing upper bound on the string tension rules out large classes of particle physics models.

It is interesting to find ways to possibly lower the bounds on the string tension.

Relevance to Particle Physics and Cosmology I

- Cosmic strings are predicted in many particle physics models beyond the "Standard Model".
- Cosmic strings are predicted to form at the end of inflation in many inflationary models.
- Cosmic strings may survive as cosmic superstrings in alternatives to inflation such as string gas cosmology.
- In models which admit cosmic strings, cosmic strings inevitably form in the early universe and persist to the present time.

Relevance to Particle Physics and Cosmology III

Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions

Cosmic strings can produce many good things for cosmology:

- String-induced mechanism of baryogenesis (R.B., A-C. Davis and M. Hindmarsh, 1991).
- Explanation for the origin of primordial magnetic fields which are coherent on galactic scales (X.Zhang and R.B. (1999).
- Explanation for cosmic ray anomalies (R.B., Y. Cai, W. Xue and X. Zhang (2009).

It is interesting to find evidence for the possible existence of cosmic strings.

Relevance to Particle Physics and Cosmology III

Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions

Cosmic strings can produce many good things for cosmology:

- String-induced mechanism of baryogenesis (R.B., A-C. Davis and M. Hindmarsh, 1991).
- Explanation for the origin of primordial magnetic fields which are coherent on galactic scales (X.Zhang and R.B. (1999).
- Explanation for cosmic ray anomalies (R.B., Y. Cai, W. Xue and X. Zhang (2009).

It is interesting to find evidence for the possible existence of cosmic strings.

Preview

Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions

Important lessons from this talk:

- Cosmic strings $\rightarrow$ nonlinearities already at high redshifts.
- Signatures of cosmic strings more pronounced at high redshifts.
- Cosmic strings lead to perturbations which are non-Gaussian.
- Cosmic strings predict specific geometrical patterns in position space.
- 21 cm surveys provide an ideal arena to look for cosmic strings (R.B., R. Danos, O. Hernandez and G. Holder, 2010).

Plan

1. Introduction
2. **Cosmic String Review**
3. Kaiser-Stebbins Effect and Cosmic String Wakes
4. Signatures of Cosmic Strings in CMB Polarization
5. Signatures of Cosmic Strings in 21cm Maps
6. Conclusions

Cosmic String I

A. Vilenkin and E. Shellard, *Cosmic Strings and other Topological Defects* (Cambridge Univ. Press, Cambridge, 1994).

- Cosmic strings form after symmetry breaking phase transitions.
- Prototypical example: Complex scalar field ϕ with "Mexican hat" potential:

$$V(\phi) = \frac{\lambda}{4}\left(|\phi|^2 - \eta^2\right)^2$$

- Vacuum manifold $\mathcal{M}$: set up field values which minimize V.

Scalar Field Potential



Cosmic String I

A. Vilenkin and E. Shellard, *Cosmic Strings and other Topological Defects* (Cambridge Univ. Press, Cambridge, 1994).

- Cosmic strings form after symmetry breaking phase transitions.
- Prototypical example: Complex scalar field ϕ with "Mexican hat" potential:

$$V(\phi) = \frac{\lambda}{4}\left(|\phi|^2 - \eta^2\right)^2$$

- Vacuum manifold $\mathcal{M}$: set up field values which minimize V.

Scalar Field Potential



Cosmic String I

A. Vilenkin and E. Shellard, *Cosmic Strings and other Topological Defects* (Cambridge Univ. Press, Cambridge, 1994).

- Cosmic strings form after symmetry breaking phase transitions.
- Prototypical example: Complex scalar field ϕ with "Mexican hat" potential:

$$V(\phi) = \frac{\lambda}{4}\left(|\phi|^2 - \eta^2\right)^2$$

- Vacuum manifold $\mathcal{M}$: set up field values which minimize V.
- At high temperature: $\phi = 0$.
- At low temperature: $|\phi| = \eta$ - but not at all $\mathbf{x}$.
- Cosmic string core: points with $|\phi| \ll \eta$.
- Criterium for the existence of cosmic strings: $\Pi_1(\mathcal{M}) \neq \infty$.

Cosmic String II

Symmetric cosmic string configuration (uniform along z axis, with core at $\rho = 0$):

$$\begin{aligned} \phi(\rho, \theta) &= f(\rho)\eta e^{i\theta} \\ f(\rho) &\rightarrow 1 \text{ for } \rho > \mathrm{w} \\ f(\rho) &\rightarrow 0 \text{ for } \rho < \mathrm{w} \end{aligned}$$

Important features:

- Width $w \sim \lambda^{-1/2}\eta^{-1}$
- Mass per unit length $\mu \sim \eta^2$ (independent of λ).

Cosmic String I

A. Vilenkin and E. Shellard, *Cosmic Strings and other Topological Defects* (Cambridge Univ. Press, Cambridge, 1994).

- Cosmic strings form after symmetry breaking phase transitions.
- Prototypical example: Complex scalar field ϕ with "Mexican hat" potential:

$$V(\phi) = \frac{\lambda}{4}\left(|\phi|^2 - \eta^2\right)^2$$

- Vacuum manifold $\mathcal{M}$: set up field values which minimize V.
- At high temperature: $\phi = 0$.
- At low temperature: $|\phi| = \eta$ - but not at all **x**.
- Cosmic string core: points with $|\phi| \ll \eta$.
- Criterium for the existence of cosmic strings: $\Pi_1(\mathcal{M}) \neq \infty$.

Scalar Field Potential



Cosmic String II

Symmetric cosmic string configuration (uniform along z axis, with core at $\rho = 0$):

$$\begin{aligned} \phi(\rho, \theta) &= f(\rho)\eta e^{i\theta} \\ f(\rho) &\rightarrow 1 \text{ for } \rho > \mathrm{w} \\ f(\rho) &\rightarrow 0 \text{ for } \rho < \mathrm{w} \end{aligned}$$

Important features:

- Width $w \sim \lambda^{-1/2}\eta^{-1}$
- Mass per unit length $\mu \sim \eta^2$ (independent of λ).

Formation of Strings

T. Kibble, Phys. Rept. **67**, 183 (1980).

- By causality, the values of ϕ in $\mathcal{M}$ cannot be correlated on scales larger than t.
- Hence, there is a probability $\mathcal{O}(1)$ that there is a string passing through a surface of side length t.
- If the field ϕ is in thermal equilibrium above the phase transition temperature, then the actual correlation length of the string network (mean separation and curvature radius of the network of infinite strings) is microscopic, given by the "Ginsburg length" $\lambda^{-1}\eta^{-1}$.

Formation of Strings

T. Kibble, Phys. Rept. **67**, 183 (1980).

- By causality, the values of ϕ in $\mathcal{M}$ cannot be correlated on scales larger than t.
- Hence, there is a probability $\mathcal{O}(1)$ that there is a string passing through a surface of side length t.
- If the field ϕ is in thermal equilibrium above the phase transition temperature, then the actual correlation length of the string network (mean separation and curvature radius of the network of infinite strings) is microscopic, given by the "Ginsburg length" $\lambda^{-1}\eta^{-1}$.

Formation of Strings

T. Kibble, Phys. Rept. **67**, 183 (1980).

- By causality, the values of ϕ in $\mathcal{M}$ cannot be correlated on scales larger than t.
- Hence, there is a probability $\mathcal{O}(1)$ that there is a string passing through a surface of side length t.
- If the field ϕ is in thermal equilibrium above the phase transition temperature, then the actual correlation length of the string network (mean separation and curvature radius of the network of infinite strings) is microscopic, given by the "Ginsburg length" $\lambda^{-1}\eta^{-1}$.

Scaling Solution I

Causality $\rightarrow$ network of cosmic strings persists at all times.

Correlation length $\xi(t) < t$ for all times $t > t_c$.

Dynamics of $\xi(t)$ is governed by a Boltzmann equation which describes the transfer of energy from long strings to string loops



Scaling Solution II

R. H. Brandenberger, Int. J. Mod. Phys. A **9**, 2117 (1994) [arXiv:astro-ph/9310041].

Analysis of the Boltzmann equation shows that $\xi(t) \sim t$ for all $t > t_c$:

- If $\xi(t) << t$ then rapid loop production and $\xi(t)/t$ increases.
- If $\xi(t) >> t$ then no loop production and $\xi(t)/t$ decreases.

Sketch of the scaling solution:



History I

Cosmic Strings

R. Brandenberger

- Cosmic strings were popular in the 1980’s as an alternative to inflation for producing a scale-invariant spectrum of cosmological perturbations.
- Cosmic strings lead to incoherent and active fluctuations (rather than coherent and passive like in inflation).
- Reason: strings on super-Hubble scales are entropy fluctuations which seed an adiabatic mode which is growing until Hubble radius crossing.
- Boomerang CMB data (1999) on the acoustic oscillations in the CMB angular power spectrum rules out cosmic strings as the main source of fluctuations..
- Interest in cosmic strings collapses.

History II

Cosmic Strings

R. Brandenberger

- Supergravity models of inflation typically yield cosmic strings after reheating (R. Jeannerot et al., 2003).
- Brane inflation models typically yield cosmic strings in the form of cosmic superstrings (Sarangi and Tye, 2002; Copeland, Myers and Polchinski, 2004).
- String Gas Cosmology may lead to a remnant scaling network of cosmic superstrings (R.B. and C. Vafa, 1989: A. Nayeri, R.B. and C. Vafa, 2006).
- → renewed interest in cosmic strings as supplementary source of fluctuations.
- Best current limit from angular spectrum of CMB anisotropies: $\sim 10\%$ of the total power can come from strings (see e.g. Wyman, Pogosian and Wasserman, 2005).
- Leads to limit $G\mu < 3 \times 10^{-7}$.

History I

Cosmic Strings

R. Brandenberger

- Introduction
- Cosmic String Review
- Kaiser-Stebbins Effect and Cosmic String Wakes
- Signatures of Cosmic Strings in CMB Polarization
- Signatures of Cosmic Strings in 21cm Maps
- Conclusions

- Cosmic strings were popular in the 1980's as an alternative to inflation for producing a scale-invariant spectrum of cosmological perturbations.
- Cosmic strings lead to incoherent and active fluctuations (rather than coherent and passive like in inflation).
- Reason: strings on super-Hubble scales are entropy fluctuations which seed an adiabatic mode which is growing until Hubble radius crossing.
- Boomerang CMB data (1999) on the acoustic oscillations in the CMB angular power spectrum rules out cosmic strings as the main source of fluctuations..
- Interest in cosmic strings collapses.


S
x


ρ
x




ρ
x



ρ
x


ρ
x


ρ
x

History II

- Supergravity models of inflation typically yield cosmic strings after reheating (R. Jeannerot et al., 2003).
- Brane inflation models typically yield cosmic strings in the form of cosmic superstrings (Sarangi and Tye, 2002; Copeland, Myers and Polchinski, 2004).
- String Gas Cosmology may lead to a remnant scaling network of cosmic superstrings (R.B. and C. Vafa, 1989: A. Nayeri, R.B. and C. Vafa, 2006).
- $\rightarrow$ renewed interest in cosmic strings as supplementary source of fluctuations.
- Best current limit from angular spectrum of CMB anisotropies: $\sim$ 10% of the total power can come from strings (see e.g. Wyman, Pogosian and Wasserman, 2005).
- Leads to limit $G\mu < 3 \times 10^{-7}$.

History II

- Supergravity models of inflation typically yield cosmic strings after reheating (R. Jeannerot et al., 2003).
- Brane inflation models typically yield cosmic strings in the form of cosmic superstrings (Sarangi and Tye, 2002; Copeland, Myers and Polchinski, 2004).
- String Gas Cosmology may lead to a remnant scaling network of cosmic superstrings (R.B. and C. Vafa, 1989: A. Nayeri, R.B. and C. Vafa, 2006).
- $\rightarrow$ renewed interest in cosmic strings as supplementary source of fluctuations.
- Best current limit from angular spectrum of CMB anisotropies: $\sim 10\%$ of the total power can come from strings (see e.g. Wyman, Pogosian and Wasserman, 2005).
- Leads to limit $G\mu < 3 \times 10^{-7}$.

History II

Cosmic Strings

R. Brandenberger

- Supergravity models of inflation typically yield cosmic strings after reheating (R. Jeannerot et al., 2003).
- Brane inflation models typically yield cosmic strings in the form of cosmic superstrings (Sarangi and Tye, 2002; Copeland, Myers and Polchinski, 2004).
- String Gas Cosmology may lead to a remnant scaling network of cosmic superstrings (R.B. and C. Vafa, 1989: A. Nayeri, R.B. and C. Vafa, 2006).
- $\rightarrow$ renewed interest in cosmic strings as supplementary source of fluctuations.
- Best current limit from angular spectrum of CMB anisotropies: $\sim$ 10% of the total power can come from strings (see e.g. Wyman, Pogosian and Wasserman, 2005).
- Leads to limit $G\mu < 3 \times 10^{-7}$.

Plan

Cosmic Strings

R. Brandenberger

Geometry of a Straight String

A. Vilenkin, Phys. Rev. D **23**, 852 (1981).

Space away from the string is locally flat (cosmic string exerts no gravitational pull).

Space perpendicular to a string is conical with deficit angle

$$\alpha = 8\pi G\mu \, ,$$


x

Geometry of a Straight String

A. Vilenkin, Phys. Rev. D **23**, 852 (1981).

Space away from the string is locally flat (cosmic string exerts no gravitational pull).

Space perpendicular to a string is conical with deficit angle

$$\alpha = 8\pi G\mu ,$$

Kaiser-Stebbins Effect

N. Kaiser and A. Stebbins, Nature **310**, 391 (1984).

Cosmic Strings

R. Brandenberger

Photons passing by the string undergo a relative Doppler shift

$$\frac{\delta T}{T} = 8\pi\gamma(v)vG\mu ,$$

Kaiser-Stebbins Effect

N. Kaiser and A. Stebbins, Nature **310**, 391 (1984).

Photons passing by the string undergo a relative Doppler shift

$$\frac{\delta T}{T} = 8\pi\gamma(v)vG\mu \, ,$$



- $\rightarrow$ network of line discontinuities in CMB anisotropy maps.
- *N.B. characteristic scale: comoving Hubble radius at the time of recombination $\rightarrow$ need good angular resolution to detect these edges.*
- Need to analyze position space maps.

- $\rightarrow$ network of line discontinuities in CMB anisotropy maps.
- *N.B. characteristic scale: comoving Hubble radius at the time of recombination $\rightarrow$ need good angular resolution to detect these edges.*
- Need to analyze position space maps.

Signature in CMB temperature anisotropy maps

R. J. Danos and R. H. Brandenberger, arXiv:0811.2004 [astro-ph].

Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions

$10^0 \times 10^0$ map of the sky at 1.5' resolution



- $\rightarrow$ network of line discontinuities in CMB anisotropy maps.
- Characteristic scale: comoving Hubble radius at the time of recombination $\rightarrow$ need good angular resolution to detect these edges.
- Need to analyze position space maps.
- Edges produced by cosmic strings are masked by the "background" noise.

Temperature map Gaussian + strings



Cosmic Strings

R. Brandenberger

- $\rightarrow$ network of line discontinuities in CMB anisotropy maps.
- Characteristic scale: comoving Hubble radius at the time of recombination $\rightarrow$ need good angular resolution to detect these edges.
- Need to analyze position space maps.
- Edges produced by cosmic strings are masked by the "background" noise.
- Edge detection algorithms: a promising way to search for strings
- Application of Canny edge detection algorithm to simulated data (SPT/ACT specification) $\rightarrow$ limit $G\mu < 2 \times 10^{-8}$ may be achievable [S. Amsel, J. Berger and R.B. (2007), A. Stewart and R.B. (2008), R. Danos and R.B. (2008)]


$P_{GS}(k) \sim (G\mu)^2 \sim$

$$P_{CS}(k) \sim (G\mu)^2 \sim 10^{-11}$$

$$P_{cs}(k) \sim (G\mu)^2 \sim 10^{-11}$$


$P_{cs}(k) \sim (G\mu)^2 \sim 10^{-11}$
$\parallel$
$k^3 |\delta_\Phi(k)|^2$


$P_{cs}(k) \sim (G\mu)^2 \sim 10^{-11}$
$= k^3 |\delta_\Phi(k)|^2$

Cosmic String Wake

J. Silk and A. Vilenkin, Phys. Rev. Lett. **53**, 1700 (1984).

Consider a cosmic string moving through the primordial gas:

Wedge-shaped region of overdensity 2 builds up behind the moving string: wake.

Cosmic String Wake

J. Silk and A. Vilenkin, Phys. Rev. Lett. **53**, 1700 (1984).

Consider a cosmic string moving through the primordial gas:

Wedge-shaped region of overdensity 2 builds up behind the moving string: wake.



Cosmic String Wake

J. Silk and A. Vilenkin, Phys. Rev. Lett. **53**, 1700 (1984).

Consider a cosmic string moving through the primordial gas:

Wedge-shaped region of overdensity 2 builds up behind the moving string: wake.



Closer look at the wedge

Cosmic Strings

R. Brandenberger

- Consider a string at time t_i $[t_{rec} < t_i < t_0]$
- moving with velocity v_s
- with typical curvature radius $c_1 t_i$







Closer look at the wedge

Cosmic Strings

R. Brandenberger

- Consider a string at time t_i $[t_{rec} < t_i < t_0]$
- moving with velocity v_s
- with typical curvature radius $c_1 t_i$



Gravitational accretion onto a wake

L. Perivolaropoulos, R.B. and A. Stebbins, Phys. Rev. D **41**, 1764 (1990).

- Initial overdensity → gravitational accretion onto the wake.
- Accretion computed using the Zeldovich approximation.
- Focus on a mass shell a physical distance $w(q,t)$ above the wake:

$$w(q,t) = a(t)\big(q - \psi\big),$$

- Gravitational accretion → ψ grows.
- Turnaround: $\dot{w}(q,t) = 0$ determines $q_{nl}(t)$ and thus the thickness of the gravitationally bound region.

Gravitational accretion onto a wake

L. Perivolaropoulos, R.B. and A. Stebbins, Phys. Rev. D **41**, 1764 (1990).

Cosmic Strings

R. Brandenberger

- Initial overdensity → gravitational accretion onto the wake.
- Accretion computed using the Zeldovich approximation.
- Focus on a mass shell a physical distance $w(q,t)$ above the wake:

$$w(q,t) = a(t)\big(q - \psi\big)\,,$$

- Gravitational accretion → ψ grows.
- Turnaround: $\dot{w}(q,t) = 0$ determines $q_{nl}(t)$ and thus the thickness of the gravitationally bound region.

Plan

Cosmic Strings

R. Brandenberger

1. Introduction
2. Cosmic String Review
3. Kaiser-Stebbins Effect and Cosmic String Wakes
4. **Signatures of Cosmic Strings in CMB Polarization**
5. Signatures of Cosmic Strings in 21cm Maps
6. Conclusions

Signature in CMB Polarization

R. Danos, R.B. and G. Holder, arXiv:1003.0905 [astro-ph.CO].

- Wake is a region of enhanced free electrons.
- CMB photons emitted at the time of recombination acquire extra polarization when they pass through a wake.
- Statistically an equal strength of E-mode and B-mode polarization is generated.
- Consider photons which at time t pass through a string segment laid down at time $t_i < t$.

$$\frac{P}{Q} \simeq \frac{24\pi}{25}\left(\frac{3}{4\pi}\right)^{1/2}\sigma_T f G\mu v_s \gamma_s \times \Omega_B \rho_c(t_0) m_p^{-1} t_0 (z(t)+1)^2 (z(t_i)+1)^{1/2}.$$

Signature in CMB Polarization II

Inserting numbers yields the result:

$$\frac{P}{Q} \sim fG\mu v_s \gamma_s \Omega_B \left(\frac{z(t)+1}{10^3}\right)^2 \left(\frac{z(t_i)+1}{10^3}\right)^3 10^7 .$$

Characteristic pattern in position space:



Is B-mode Polarization the Holy Grail of Inflation?

R.B., arXiv:1104.3581 [astro-ph.CO].

- Cosmic strings produce direct B-mode polarization.
- → gravitational waves not the only source of primordial B-mode polarization.
- Cosmic string loop oscillations produce a scale-invariant spectrum of primordial gravitational waves with a contribution to $\delta T / T$ which is comparable to that induced by scalar fluctuations (see e.g. A. Albrecht, R.B. and N. Turok, 1986).
- → a detection of gravitational waves through B-mode polarization is more likely to be a sign of something different than inflation.

Is B-mode Polarization the Holy Grail of Inflation?

R.B., arXiv:1104.3581 [astro-ph.CO].

- Cosmic strings produce direct B-mode polarization.
- → gravitational waves not the only source of primordial B-mode polarization.
- Cosmic string loop oscillations produce a scale-invariant spectrum of primordial gravitational waves with a contribution to $\delta T/T$ which is comparable to that induced by scalar fluctuations (see e.g. A. Albrecht, R.B. and N. Turok, 1986).
- → a detection of gravitational waves through B-mode polarization is more likely to be a sign of something different than inflation.

Is B-mode Polarization the Holy Grail of Inflation?

R.B., arXiv:1104.3581 [astro-ph.CO].

- Cosmic strings produce direct B-mode polarization.
- $\rightarrow$ gravitational waves not the only source of primordial B-mode polarization.
- Cosmic string loop oscillations produce a scale-invariant spectrum of primordial gravitational waves with a contribution to $\delta T/T$ which is comparable to that induced by scalar fluctuations (see e.g. A. Albrecht, R.B. and N. Turok, 1986).
- $\rightarrow$ a detection of gravitational waves through B-mode polarization is more likely to be a sign of something different than inflation.

Is B-mode Polarization the Holy Grail of Inflation?

R.B., arXiv:1104.3581 [astro-ph.CO].

- Cosmic strings produce direct B-mode polarization.
- → gravitational waves not the only source of primordial B-mode polarization.
- Cosmic string loop oscillations produce a scale-invariant spectrum of primordial gravitational waves with a contribution to $\delta T/T$ which is comparable to that induced by scalar fluctuations (see e.g. A. Albrecht, R.B. and N. Turok, 1986).
- → a detection of gravitational waves through B-mode polarization is more likely to be a sign of something different than inflation.

Is B-mode Polarization the Holy Grail of Inflation? II

R.B., A. Nayeri, S. Patil and C. Vafa, hep-th/0604126.

- N.B. String Gas Cosmology produces a spectrum of gravitational waves with an amplitude larger than in many single field inflation models and with a small blue tilt.
- Inflationary cosmology must produce a red tilt.
- Observing a blue tilt of the gravitational wave spectrum would falsify inflationary cosmology.
- B-mode polarization may be the holy grail of early universe cosmology, but not of inflation.

Is B-mode Polarization the Holy Grail of Inflation? II

R.B., A. Nayeri, S. Patil and C. Vafa, hep-th/0604126.

- N.B. String Gas Cosmology produces a spectrum of gravitational waves with an amplitude larger than in many single field inflation models and with a small blue tilt.
- Inflationary cosmology must produce a red tilt.
- Observing a blue tilt of the gravitational wave spectrum would falsify inflationary cosmology.
- B-mode polarization may be the holy grail of early universe cosmology, but not of inflation.

Is B-mode Polarization the Holy Grail of Inflation? II

R.B., A. Nayeri, S. Patil and C. Vafa, hep-th/0604126.

- N.B. String Gas Cosmology produces a spectrum of gravitational waves with an amplitude larger than in many single field inflation models and with a small blue tilt.
- Inflationary cosmology must produce a red tilt.
- Observing a blue tilt of the gravitational wave spectrum would falsify inflationary cosmology.
- B-mode polarization may be the holy grail of early universe cosmology, but not of inflation.

Is B-mode Polarization the Holy Grail of Inflation? II

R.B., A. Nayeri, S. Patil and C. Vafa, hep-th/0604126.

- N.B. String Gas Cosmology produces a spectrum of gravitational waves with an amplitude larger than in many single field inflation models and with a small blue tilt.
- Inflationary cosmology must produce a red tilt.
- Observing a blue tilt of the gravitational wave spectrum would falsify inflationary cosmology.
- B-mode polarization may be the holy grail of early universe cosmology, but not of inflation.

Plan

Motivation

R.B., D. Danos, O. Hernandez and G. Holder, arXiv:1006.2514; O. Hernandez, Yi Wang, R.B. and J. Fong, arXiv:1104.3337.

Cosmic Strings

R. Brandenberger

- 21 cm surveys: new window to map the high redshift universe, in particular the “dark ages".
- Cosmic strings produce nonlinear structures at high redshifts.
- These nonlinear structures will leave imprints in 21 cm maps.
- 21 cm surveys provide 3-d maps → potentially more data than the CMB.
- → 21 cm surveys is a promising window to search for cosmic strings.

Motivation

R.B., D. Danos, O. Hernandez and G. Holder, arXiv:1006.2514; O. Hernandez, Yi Wang, R.B. and J. Fong, arXiv:1104.3337.

- 21 cm surveys: new window to map the high redshift universe, in particular the "dark ages".
- Cosmic strings produce nonlinear structures at high redshifts.
- These nonlinear structures will leave imprints in 21 cm maps.
- 21 cm surveys provide 3-d maps → potentially more data than the CMB.
- → 21 cm surveys is a promising window to search for cosmic strings.

Motivation

R.B., D. Danos, O. Hernandez and G. Holder, arXiv:1006.2514; O. Hernandez, Yi Wang, R.B. and J. Fong, arXiv:1104.3337.

- 21 cm surveys: new window to map the high redshift universe, in particular the "dark ages".
- Cosmic strings produce nonlinear structures at high redshifts.
- These nonlinear structures will leave imprints in 21 cm maps.
- 21 cm surveys provide 3-d maps $\rightarrow$ potentially more data than the CMB.
- $\rightarrow$ 21 cm surveys is a promising window to search for cosmic strings.

The Effect

Cosmic Strings

R. Brandenberger

- $10^3 > z > 10$: baryonic matter dominated by neutral H.
- Neutral H has hydrogen hyperfine absorption/emission line.
- String wake is a gas cloud with special geometry which emits/absorbs 21cm radiation.
- Whether signal is emission/absorption depends on the temperature of the gas cloud.

The Effect

Cosmic Strings

R. Brandenberger

- $10^3 > z > 10$: baryonic matter dominated by neutral H.
- Neutral H has hydrogen hyperfine absorption/emission line.
- String wake is a gas cloud with special geometry which emits/absorbs 21cm radiation.
- Whether signal is emission/absorption depends on the temperature of the gas cloud.

The Effect

Cosmic Strings

R. Brandenberger

- $10^3 > z > 10$: baryonic matter dominated by neutral H.
- Neutral H has hydrogen hyperfine absorption/emission line.
- String wake is a gas cloud with special geometry which emits/absorbs 21cm radiation.
- Whether signal is emission/absorption depends on the temperature of the gas cloud.

The Effect

Cosmic Strings

R. Brandenberger

- $10^3 > z > 10$: baryonic matter dominated by neutral H.
- Neutral H has hydrogen hyperfine absorption/emission line.
- String wake is a gas cloud with special geometry which emits/absorbs 21cm radiation.
- Whether signal is emission/absorption depends on the temperature of the gas cloud.


t
δv'
δv'

Key general formulas

Cosmic Strings

R. Brandenberger

Brightness temperature:

$$T_b(\nu) = T_S\left(1 - e^{-\tau_\nu}\right) + T_\gamma(\nu) e^{-\tau_\nu} \, ,$$

Spin temperature:

$$T_S = \frac{1 + x_c}{1 + x_c T_\gamma / T_K} T_\gamma \, .$$

T_K: gas temperature in the wake, x_c collision coefficient

Relative brightness temperature:

$$\delta T_b(\nu) = \frac{T_b(\nu) - T_\gamma(\nu)}{1 + z}$$

Cosmic Strings

R. Brandenberger



Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions




t
δv'
δv'

Cosmic Strings

R. Brandenberger



Cosmic Strings

R. Brandenberger



Plan

Cosmic Strings

R. Brandenberger

Is B-mode Polarization the Holy Grail of Inflation? II

R.B., A. Nayeri, S. Patil and C. Vafa, hep-th/0604126.

- N.B. String Gas Cosmology produces a spectrum of gravitational waves with an amplitude larger than in many single field inflation models and with a small blue tilt.
- Inflationary cosmology must produce a red tilt.
- Observing a blue tilt of the gravitational wave spectrum would falsify inflationary cosmology.
- B-mode polarization may be the holy grail of early universe cosmology, but not of inflation.

Motivation

R.B., D. Danos, O. Hernandez and G. Holder, arXiv:1006.2514; O. Hernandez, Yi Wang, R.B. and J. Fong, arXiv:1104.3337.

- 21 cm surveys: new window to map the high redshift universe, in particular the "dark ages".
- Cosmic strings produce nonlinear structures at high redshifts.
- These nonlinear structures will leave imprints in 21 cm maps.
- 21 cm surveys provide 3-d maps → potentially more data than the CMB.
- → 21 cm surveys is a promising window to search for cosmic strings.


t
δv'
δv'

Key general formulas

Brightness temperature:

$$T_b(\nu) = T_S\left(1 - e^{-\tau_\nu}\right) + T_\gamma(\nu) e^{-\tau_\nu},$$

Spin temperature:

$$T_S = \frac{1 + x_c}{1 + x_c T_\gamma / T_K} T_\gamma .$$

T_K: gas temperature in the wake, x_c collision coefficient

Relative brightness temperature:

$$\delta T_b(\nu) = \frac{T_b(\nu) - T_\gamma(\nu)}{1+z}$$

Cosmic Strings

R. Brandenberger

Optical depth:

$$\tau_\nu = \frac{3c^2 A_{10}}{4\nu^2}\left(\frac{\hbar\nu}{k_B T_S}\right)\frac{N_{HI}}{4}\phi(\nu)\,,$$

Frequency dispersion

$$\frac{\delta\nu}{\nu} = 2\sin(\theta)\tan\theta\frac{Hw}{c}\,,$$

Line profile:

$$\phi(\nu) = \frac{1}{\delta\nu} \text{ for } \nu\,\epsilon\,[\nu_{10} - \frac{\delta\nu}{2}, \nu_{10} + \frac{\delta\nu}{2}]\,,$$

Application to Cosmic String Wakes

Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions

Wake temperature T_K:

$$T_K \simeq [20\ \mathrm{K}](G\mu)_6^2 (v_s\gamma_s)^2 \frac{z_i + 1}{z + 1},$$

determined by considering thermalization at the shock which occurs after turnaround when $w = 1/2 w_{max}$ (see Eulerian hydro simulations by A. Sornborger et al, 1997).

Thickness in redshift space:

$$\frac{\delta\nu}{\nu} = \frac{24\pi}{15} G\mu v_s\gamma_s (z_i + 1)^{1/2} (z(t) + 1)^{-1/2}$$
$$\simeq 3 \times 10^{-5} (G\mu)_6 (v_s\gamma_s),$$

using $z_i + 1 = 10^3$ and $z + 1 = 30$ in the second line.

Application to Cosmic String Wakes

Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions

Wake temperature T_K:

$$T_K \simeq [20\ \mathrm{K}](G\mu)_6^2(v_s\gamma_s)^2\frac{z_i+1}{z+1},$$

determined by considering thermalization at the shock which occurs after turnaround when $w = 1/2 w_{max}$ (see Eulerian hydro simulations by A. Sornborger et al, 1997).

Thickness in redshift space:

$$\begin{aligned}\frac{\delta\nu}{\nu} &= \frac{24\pi}{15} G\mu v_s\gamma_s(z_i+1)^{1/2}(z(t)+1)^{-1/2} \\ &\simeq 3\times 10^{-5}(G\mu)_6(v_s\gamma_s),\end{aligned}$$

using $z_i + 1 = 10^3$ and $z + 1 = 30$ in the second line.

Relative brightness temperature:

$$\delta T_b(\nu) = [0.07\,\mathrm{K}]\frac{x_c}{1+x_c}(1-\frac{T_\gamma}{T_K})(1+z)^{1/2}$$
$$\sim 200mK \quad \text{for} \quad z+1=30\,.$$

Signal is emission if $T_K > T_\gamma$ and absorption otherwise.

Critical curve (transition from emission to absorption):

$$(G\mu)_6^2 \simeq 0.1(v_s\gamma_s)^{-2}\frac{(z+1)^2}{z_i+1}$$

Scalings of various temperatures

Cosmic Strings

R. Brandenberger



Top curve: $(G\mu)_6 = 1$, bottom curve: $(G\mu)_6 = 0.3$

Geometry of the signal

Cosmic Strings

R. Brandenberger



Scalings of various temperatures

Cosmic Strings

R. Brandenberger



Top curve: $(G\mu)_6 = 1$, bottom curve: $(G\mu)_6 = 0.3$

Relative brightness temperature:

$$\delta T_b(\nu) = [0.07\,\mathrm{K}]\frac{x_c}{1+x_c}\left(1-\frac{T_\gamma}{T_K}\right)(1+z)^{1/2}$$
$$\sim 200mK \quad \text{for} \quad z+1=30\,.$$

Signal is emission if $T_K > T_\gamma$ and absorption otherwise.

Critical curve (transition from emission to absorption):

$$(G\mu)_6^2 \simeq 0.1(v_s\gamma_s)^{-2}\,\frac{(z+1)^2}{z_i+1}$$

Relative brightness temperature:

$$\delta T_b(\nu) = [0.07\,\mathrm{K}] \frac{x_c}{1+x_c}\left(1 - \frac{T_\gamma}{T_K}\right)(1+z)^{1/2}$$
$$\sim 200mK \quad \text{for} \quad z+1 = 30\,.$$

Signal is emission if $T_K > T_\gamma$ and absorption otherwise.

Critical curve (transition from emission to absorption):

$$(G\mu)_6^2 \simeq 0.1(v_s\gamma_s)^{-2}\frac{(z+1)^2}{z_i+1}$$

Application to Cosmic String Wakes

Cosmic Strings

R. Brandenberger

Wake temperature T_K:

$$T_K \simeq [20\ \mathrm{K}](G\mu)_6^2(v_s\gamma_s)^2\frac{z_i+1}{z+1},$$

determined by considering thermalization at the shock which occurs after turnaround when $w = 1/2 w_{max}$ (see Eulerian hydro simulations by A. Sornborger et al, 1997).

Thickness in redshift space:

$$\begin{aligned}\frac{\delta\nu}{\nu} &= \frac{24\pi}{15}G\mu v_s\gamma_s(z_i+1)^{1/2}(z(t)+1)^{-1/2} \\ &\simeq 3\times 10^{-5}(G\mu)_6(v_s\gamma_s),\end{aligned}$$

using $z_i + 1 = 10^3$ and $z + 1 = 30$ in the second line.

Relative brightness temperature:

$$\delta T_b(\nu) = [0.07\,\mathrm{K}]\frac{x_c}{1+x_c}(1-\frac{T_\gamma}{T_K})(1+z)^{1/2}$$
$$\sim 200mK \quad \text{for} \quad z+1=30\,.$$

Signal is emission if $T_K > T_\gamma$ and absorption otherwise.

Critical curve (transition from emission to absorption):

$$(G\mu)_6^2 \simeq 0.1(v_s\gamma_s)^{-2}\frac{(z+1)^2}{z_i+1}$$

Geometry of the signal

Cosmic Strings

R. Brandenberger



Conclusions

- Cosmic strings $\rightarrow$ nonlinearities already at high redshifts.
- Signatures of cosmic strings more pronounced at high redshifts.
- Cosmic strings lead to perturbations which are non-Gaussian.
- Cosmic strings predict specific geometrical patterns in position space.
- 21 cm surveys provide an ideal arena to look for cosmic strings.
- Cosmic string wakes produce distinct wedges in redshift space with enhanced 21cm absorption or emission.

Conclusions

Cosmic Strings

R. Brandenberger

- Cosmic strings → nonlinearities already at high redshifts.
- Signatures of cosmic strings more pronounced at high redshifts.
- Cosmic strings lead to perturbations which are non-Gaussian.
- Cosmic strings predict specific geometrical patterns in position space.
- 21 cm surveys provide an ideal arena to look for cosmic strings.
- Cosmic string wakes produce distinct wedges in redshift space with enhanced 21cm absorption or emission.

Scalings of various temperatures

Cosmic Strings

R. Brandenberger



Top curve: $(G\mu)_6 = 1$, bottom curve: $(G\mu)_6 = 0.3$

The Effect

Cosmic Strings

R. Brandenberger

Introduction

Cosmic String Review

Kaiser-Stebbins Effect and Cosmic String Wakes

Signatures of Cosmic Strings in CMB Polarization

Signatures of Cosmic Strings in 21cm Maps

Conclusions

- $10^3 > z > 10$: baryonic matter dominated by neutral H.
- Neutral H has hydrogen hyperfine absorption/emission line.
- String wake is a gas cloud with special geometry which emits/absorbs 21cm radiation.
- Whether signal is emission/absorption depends on the temperature of the gas cloud.

Plan

Is B-mode Polarization the Holy Grail of Inflation? II

R.B., A. Nayeri, S. Patil and C. Vafa, hep-th/0604126.

Cosmic Strings

R. Brandenberger

- N.B. String Gas Cosmology produces a spectrum of gravitational waves with an amplitude larger than in many single field inflation models and with a small blue tilt.
- Inflationary cosmology must produce a red tilt.
- Observing a blue tilt of the gravitational wave spectrum would falsify inflationary cosmology.
- B-mode polarization may be the holy grail of early universe cosmology, but not of inflation.