Title: A Relativistic Relative of GS Superstrings on AdS5 x S5

Date: Aug 19, 2011 02:30 PM

URL: http://pirsa.org/11080035

Abstract: The motion of superstrings on symmetric space target spaces is classically equivalent, via the Pohlmeyer reduction, to a family of 2-d relativistic integrable field theories known as semisymmetric space sine-Gordon (SSSSG) theories. In this talk I will review recent progress in constructing the relativistic S-matrix corresponding to the quantum solution of the AdS5 x S5 SSSSG theory.

Pirsa: 11080035 Page 1/166

A relativistic relative of GS supertrings on $AdS_5 \times S^5$

The Semi-Symmetric Space sine-Gordon theory

J. Luis Miramontes

U. Santiago de Compostela/IGFAE, (Spain)

(*) in collaboration with Ben Hoare (London) and Timothy J. Hollowood (Swansea)

Based on arXiv:1107.0628, 1104.2429, ...

Outline

- The SSSG and SSSSG theories
- 2 The $AdS_5 \times S^5$ SSSSG theory
- 3 The relativistic S-matrix
- 4 Bound states: bootstrap
- Open questions

Pohlmeyer, Eichenherr, Forger, D'Auria, Regge, ...'76-81

.....



Symmetric space sine-Gordon (SSSG) theories

Tseytlin'03 Mikhailov'05

• Relativistic integrable theories in 1+1 dimensions that are classically equivalent, via the Pohlmeyer reduction, to the non-relativistic (gauged fixed) world-sheet theories of strings on symmetric space spacetimes like $\mathbb{R}_t \times S^n$, $\mathbb{R}_t \times \mathbb{C}P^n$, $AdS_n \times S^1_n$, AdS_n , ...

Hofman-Maldacena'06 Chen-N.Dorey-Okamura'06

• Admit soliton solutions that, for $\mathbb{R}_t \times S^n$ or $\mathbb{R}_t \times \mathbb{C}P^n$, are the images of the string giant magnons

Pirsa: 11080035 Page 4/166

Pohlmeyer, Eichenherr, Forger, D'Auria, Regge, ...'76-81



Symmetric space sine-Gordon (SSSG) theories

Tseytlin'03 Mikhailov'05

• Relativistic integrable theories in 1+1 dimensions that are classically equivalent, via the Pohlmeyer reduction, to the non-relativistic (gauged fixed) world-sheet theories of strings on symmetric space spacetimes like $\mathbb{R}_t \times S^n$, $\mathbb{R}_t \times \mathbb{C}P^n$, $AdS_n \times S^1_\vartheta$, AdS_n , ...

Hofman-Maldacena 106 Chen-N. Dorey-Okamura 106

• Admit soliton solutions that, for $\mathbb{R}_t \times S^n$ or $\mathbb{R}_t \times \mathbb{C}P^n$, are the images of the string giant magnons

Pirsa: 11080035 Page 5/166

 To describe the world-sheet theory of superstrings with all the fermionic degrees of freedom one has to generalize the SSSG theories to the case where the symmetric space is replaced by a semi-symmetric space F/G, with F a supergroup

> Serganova 33 Zarembo 310

- $\rightarrow \mathbb{Z}_4$ automorphism: $\sigma^4 = 1$, $\sigma(G) = G$
- → Lie superalgebra decomposition: $f = \underbrace{f_0}_{\text{even}} \oplus \underbrace{f_1}_{\text{odd}} \oplus \underbrace{f_2}_{\text{even}} \oplus \underbrace{f_3}_{\text{odd}}$



Semisymmetric space sine-Gordon (SSSSG) theories

Metsaev-Tseytlin'98

•
$$AdS_5 \times S^5 \longrightarrow PSU(2,2|4)/SO(4,1) \times SO(5)$$

Arutyunov-Frolov'08 Stefanski'08

•
$$AdS_4 \times \mathbb{C}P^3 \longrightarrow OSp(6|4)/SO(3,1) \times U(3)$$

 To describe the world-sheet theory of superstrings with all the fermionic degrees of freedom one has to generalize the SSSG theories to the case where the symmetric space is replaced by a semi-symmetric space F/G, with F a supergroup

> Serganova 83 Zarembo 10

- $\rightarrow \mathbb{Z}_4$ automorphism: $\sigma^4 = 1$, $\sigma(G) = G$
- \mapsto Lie superalgebra decomposition: $f = \underbrace{f_0}_{\text{even}} \oplus \underbrace{f_1}_{\text{odd}} \oplus \underbrace{f_2}_{\text{even}} \oplus \underbrace{f_3}_{\text{odd}}$



Semisymmetric space sine-Gordon (SSSSG) theories

• $AdS_5 \times S^5 \longrightarrow PSU(2,2|4)/SO(4,1) \times SO(5)$

Metsaev-Tseytlin'98

Arutyunov-Frolov'08 Stefanski'08

•
$$AdS_4 \times \mathbb{C}P^3 \longrightarrow OSp(6|4)/SO(3,1) \times U(3)$$

Grigoriev-Tseytlin'08 Mikhailov-SchaferNakemi'08



$$\mathcal{F}/G = \frac{PSU(2,2|4)}{Sp(2,2) \times Sp(4)}$$

is classically equivalent, via a fermionic generalization of Pohlmeyer reduction, to the non-relativistic (gauge fixed) Green-Schwarz superstring world-sheet theory on $AdS_5 \times S^5$

Grigoriev-Tseytlin'08 Mikhailov-SchaferNakemi'08

The relativistic SSSSG theory for

$$\mathcal{F}/G = \frac{PSU(2,2|4)}{Sp(2,2) \times Sp(4)}$$

is classically equivalent, via a fermionic generalization of Pohlmeyer reduction, to the non-relativistic (gauge fixed) Green-Schwarz superstring world-sheet theory on $AdS_5 \times S^5$

Pirsa: 11080035 Page 9/166

The relativistic SSSSG theory for

$$\mathcal{F}/G = \frac{PSU(2,2|4)}{Sp(2,2) \times Sp(4)}$$

is classically equivalent, via a fermionic generalization of Pohlmeyer reduction, to the non-relativistic (gauge fixed) Green-Schwarz superstring world-sheet theory on $AdS_5 \times S^5$

 In general, the equivalence between the (S)SSSG theories and (super)string world-sheet theories is expected to be purely classical

> Mikhailov'05 Schmidtt'11

They have different Hamiltonian structures

The relativistic SSSSG theory for

$$\mathcal{F}/G = \frac{PSU(2,2|4)}{Sp(2,2) \times Sp(4)}$$

is classically equivalent, via a fermionic generalization of Pohlmeyer reduction, to the non-relativistic (gauge fixed) Green-Schwarz superstring world-sheet theory on $AdS_5 \times S^5$

 In general, the equivalence between the (S)SSSG theories and (super)string world-sheet theories is expected to be purely classical

> Mikhailov 05 Schmidtt 111

They have different Hamiltonian structures

The relativistic SSSSG theory for

$$\mathcal{F}/G = \frac{PSU(2,2|4)}{Sp(2,2) \times Sp(4)}$$

is classically equivalent, via a fermionic generalization of Pohlmeyer reduction, to the non-relativistic (gauge fixed) Green-Schwarz superstring world-sheet theory on $AdS_5 \times S^5$

- In general, the equivalence between the (S)SSSG theories and (super)string world-sheet theories is expected to be purely classical
 - They have different Hamiltonian structures

Mikhailov'05 Schmidtt'11

Grigoriev-Tseytlin'08

★ ...but the equivalence has been conjectured to remain in the (conformally invariant) quantum theory

The relativistic SSSSG theory for

$$\mathcal{F}/G = \frac{PSU(2,2|4)}{Sp(2,2) \times Sp(4)}$$

is classically equivalent, via a fermionic generalization of Pohlmeyer reduction, to the non-relativistic (gauge fixed) Green-Schwarz superstring world-sheet theory on $AdS_5 \times S^5$

 In general, the equivalence between the (S)SSSG theories and (super)string world-sheet theories is expected to be purely classical

> Mikhailov'05 Schmidtt'11

They have different Hamiltonian structures

Grigoriev-Tseytlin'08

★ ...but the equivalence has been conjectured to remain in the (conformally invariant) quantum theory

The equivalence has already passed several tests: [— A. Tseytlin's talk]

Roiban-Tseytlin'09

• The $AdS_5 \times S^5$ SSSSG theory is UV-finite

Hoare-Iwashita-Tseytlin'09 Iwashita'10

Semiclassical partition function matches with string theory at one loop

Pirsa: 11080035 Page 14/166

The equivalence has already passed several tests: [— A. Tseytlin's talk]

Roiban-Tseytlin'09

• The $AdS_5 \times S^5$ SSSSG theory is UV-finite

Hoare-Iwashita-Tseytlin'09 Iwashita'10

Semiclassical partition function matches with string theory at one loop

Pirsa: 11080035 Page 15/166

The equivalence has already passed several tests: [— A. Tseytlin's talk]

Roiban-Tseytlin'09


• The $AdS_5 \times S^5$ SSSSG theory is UV-finite

Hoare-Iwashita-Tseytlin'09 Iwashita'10

Semiclassical partition function matches with string theory at one loop



Outstanding problem: Find the exact relativistic S-matrix of the $AdS_5 \times S^5$ SSSSG theory and clarify the relationship with the non-relativistic superstring S-matrix \longrightarrow interesting by itself!



Roiban-Tseytlin'09

• The $AdS_5 \times S^5$ SSSSG theory is UV-finite

Hoare-Iwashita-Tseytlin'09 Iwashita'10

Semiclassical partition function matches with string theory at one loop



Outstanding problem: Find the exact relativistic S-matrix of the $AdS_5 \times S^5$ SSSSG theory and clarify the relationship with the non-relativistic superstring S-matrix \longrightarrow interesting by itself!

Our approach:

- Focus not so much on the Lagrangian and perturbation theory but rather on the solitons: perturbative fields re-appear!
- Quantize the moduli space dynamics of the solitons yielding the semi-classical spectrum
- Conjecture the S-matrix by imposing all the axioms of S-matrix theory

 and solving the bootstrap (account for all poles on the physical Page 17/106)

The equivalence has already passed several tests: [-- A. Tseytlin's talk]

Roiban-Tseytlin'09

• The $AdS_5 \times S^5$ SSSSG theory is UV-finite

Hoare-Iwashita-Tseytlin'09 Iwashita'10

Semiclassical partition function matches with string theory at one loop



Outstanding problem: Find the exact relativistic S-matrix of the $AdS_5 \times S^5$ SSSSG theory and clarify the relationship with the non-relativistic superstring S-matrix \longrightarrow interesting by itself!

Our approach:

- Focus not so much on the Lagrangian and perturbation theory but rather on the solitons: perturbative fields re-appear!
- Quantize the moduli space dynamics of the solitons yielding the semi-classical spectrum
- Conjecture the S-matrix by imposing all the axioms of S-matrix theory

 Pirsa: 11080035 and solving the bootstrap (account for all poles on the physical Patrions)

The (relativistic) SSSSG Lagrangian

Grigoriev-Tseytlin'08 Mikhailov-SchaferNakemi'08

$$\mathcal{L} = \mathcal{L}_{\text{gWZW}}[G/H] - \frac{k}{\pi} \text{STr} \left(\Lambda \gamma^{-1} \Lambda \gamma \right)$$

$$+ \frac{k}{2\pi} \text{STr} \left(\psi_{+} [\Lambda, D_{-} \psi_{+}] - \psi_{-} [\Lambda, D_{+} \psi_{-}] - 2\psi_{+} \gamma^{-1} \psi_{-} \gamma \right)$$

- $\gamma \in G = e^{\mathfrak{f}_0}$ and $A_{\pm} \longrightarrow$ bosonic fields $\psi_+ \in \mathfrak{f}_1, \ \psi_- \in \mathfrak{f}_3 \longrightarrow$ fermionic fields
- The potential is fixed by $\Lambda \in \mathfrak{f}_2$ (constant)
- Gauge symmetry group $H = SU(2)^{\times 4} \subset G$ $[H, \Lambda] = 0$
- The coupling constant is the level of the WZW term $\equiv k$

The (relativistic) SSSSG Lagrangian

Grigoriev-Tseytlin'08 Mikhailov-SchaferNakemi'08

$$\mathcal{L} = \mathcal{L}_{\text{gWZW}}[G/H] - \frac{k}{\pi} \text{STr} \left(\Lambda \gamma^{-1} \Lambda \gamma \right)$$

$$+ \frac{k}{2\pi} \text{STr} \left(\psi_{+} [\Lambda, D_{-} \psi_{+}] - \psi_{-} [\Lambda, D_{+} \psi_{-}] - 2\psi_{+} \gamma^{-1} \psi_{-} \gamma \right)$$

- $\gamma \in G = e^{\mathfrak{f}_0}$ and $A_{\pm} \longrightarrow$ bosonic fields $\psi_+ \in \mathfrak{f}_1, \ \psi_- \in \mathfrak{f}_3 \longrightarrow$ fermionic fields
- The potential is fixed by $\Lambda \in \mathfrak{f}_2$ (constant)

Gauge symmetry group
$$H = SU(2)^{\times 4} \subset G$$
 $[H, \Lambda] = 0$

The coupling constant is the level of the WZW term $\equiv k$

The $AdS_5 \times S^5$ SSSSG theory

Equations of motion

• Zero-curvature conditions $\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$

$$\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$$

$$\mathcal{L}_{+}(z) = \partial_{+} + \gamma^{-1} \partial_{+} \gamma + \gamma^{-1} A_{+} \gamma + z \psi_{+} - z^{2} \Lambda,$$

$$\mathcal{L}_{-}(z) = \partial_{-} + A_{-} + z^{-1} \gamma^{-1} \psi_{-} \gamma - z^{-2} \gamma^{-1} \Lambda \gamma$$

 $z = \text{spectral parameter} \Rightarrow \text{Classical integrability}$

Page 21/166 Pirsa: 11080035

The $AdS_5 \times S^5$ SSSSG theory

Equations of motion

• Zero-curvature conditions $\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$

$$\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$$

$$\mathcal{L}_{+}(z) = \partial_{+} + \gamma^{-1}\partial_{+}\gamma + \gamma^{-1}A_{+}\gamma + z\psi_{+} - z^{2}\Lambda,$$

$$\mathcal{L}_{-}(z) = \partial_{-} + A_{-} + z^{-1}\gamma^{-1}\psi_{-}\gamma - z^{-2}\gamma^{-1}\Lambda\gamma$$

 $z = \text{spectral parameter} \Rightarrow \text{Classical integrability}$

Pirsa: 11080035 Page 22/166

• Zero-curvature conditions $\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$

$$\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$$

$$\mathcal{L}_{+}(z) = \partial_{+} + \gamma^{-1} \partial_{+} \gamma + \gamma^{-1} A_{+} \gamma + z \psi_{+} - z^{2} \Lambda,$$

$$\mathcal{L}_{-}(z) = \partial_{-} + A_{-} + z^{-1} \gamma^{-1} \psi_{-} \gamma - z^{-2} \gamma^{-1} \Lambda \gamma$$

$$z = \text{spectral parameter} \Rightarrow \text{Classical integrability}$$

Relativistic equations! Lorentz boost $x^{\pm} \to \lambda x^{\pm} \sim |z \to \lambda^{1/2} z|$

• Zero-curvature conditions $\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$

$$\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$$

$$\mathcal{L}_{+}(z) = \partial_{+} + \gamma^{-1}\partial_{+}\gamma + \gamma^{-1}A_{+}\gamma + z\psi_{+} - z^{2}\Lambda,$$

$$\mathcal{L}_{-}(z) = \partial_{-} + A_{-} + z^{-1}\gamma^{-1}\psi_{-}\gamma - z^{-2}\gamma^{-1}\Lambda\gamma$$

$$z = \text{spectral parameter} \Rightarrow \text{Classical integrability}$$

Relativistic equations! Lorentz boost $x^{\pm} \to \lambda x^{\pm} \sim |z \to \lambda^{1/2} z|$

• Zero-curvature conditions $\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$

$$\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$$

$$\mathcal{L}_{+}(z) = \partial_{+} + \gamma^{-1} \partial_{+} \gamma + \gamma^{-1} A_{+} \gamma + z \psi_{+} - z^{2} \Lambda,$$

$$\mathcal{L}_{-}(z) = \partial_{-} + A_{-} + z^{-1} \gamma^{-1} \psi_{-} \gamma - z^{-2} \gamma^{-1} \Lambda \gamma$$

$$z = \text{spectral parameter} \Rightarrow \text{Classical integrability}$$

- Relativistic equations! Lorentz boost $x^{\pm} \to \lambda x^{\pm} \sim |z \to \lambda^{1/2} z|$
- Integrability is controlled by the twisted affine loop superalgebra

$$\mathcal{L}(\mathfrak{psu}(2,2|4),\sigma) = \bigoplus_{n \in \mathbb{Z}} \bigoplus_{j=0}^{3} z^{4n+j} \otimes f_j = \bigoplus_{k \in \mathbb{Z}} \hat{\mathfrak{f}}_k, \quad [\hat{\mathfrak{f}}_k,\hat{\mathfrak{f}}_l] \subset \hat{\mathfrak{f}}_{k+l}$$

• Zero-curvature conditions $\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$

$$\left[\mathcal{L}_{\mu}(z),\mathcal{L}_{\nu}(z)\right]=0$$

$$\mathcal{L}_{+}(z) = \partial_{+} + \gamma^{-1} \partial_{+} \gamma + \gamma^{-1} A_{+} \gamma + z \psi_{+} - z^{2} \Lambda,$$

$$\mathcal{L}_{-}(z) = \partial_{-} + A_{-} + z^{-1} \gamma^{-1} \psi_{-} \gamma - z^{-2} \gamma^{-1} \Lambda \gamma$$

$$z = \text{spectral parameter} \Rightarrow \text{Classical integrability}$$

- Relativistic equations! Lorentz boost $x^{\pm} \to \lambda x^{\pm} \sim |z \to \lambda^{1/2} z|$
- Integrability is controlled by the twisted affine loop superalgebra

$$\mathcal{L}(\mathfrak{psu}(2,2|4),\sigma) = \bigoplus_{n \in \mathbb{Z}} \bigoplus_{j=0}^{3} z^{4n+j} \otimes f_j = \bigoplus_{k \in \mathbb{Z}} \hat{\mathfrak{f}}_k, \quad [\hat{\mathfrak{f}}_k,\hat{\mathfrak{f}}_l] \subset \hat{\mathfrak{f}}_{k+l}$$

The AdS₅ × S⁵ SSSSG theory

Hidden symmetries

Infinite (classical) symmetry algebra

$$\hat{\mathfrak{f}}^{\perp} = \operatorname{Ker}(\operatorname{ad}\Lambda) \cap \mathcal{L}(\mathfrak{psu}(2,2|4),\sigma)$$

Infinite (classical) symmetry algebra

$$\hat{\mathfrak{f}}^{\perp} = \operatorname{Ker}(\operatorname{ad}\Lambda) \cap \mathcal{L}(\mathfrak{psu}(2,2|4),\sigma)$$

Hollowood-JLM'11 Hoare-Tseytlin'11

The elements of grade ± 1 (or Lorentz spin $\pm 1/2$) generate SUSY transformations whose closure is the (exotic) $\mathcal{N}=(8|8)$ superalgebra

$$\mathfrak{s} = \big(\mathfrak{psu}(2|2) \oplus \mathfrak{psu}(2|2)\big) \ltimes \big(\mathbb{R} \oplus \mathbb{R}\big) = \mathfrak{s}_{-2} \oplus \mathfrak{s}_{-1} \oplus \mathfrak{s}_0 \oplus \mathfrak{s}_{+1} \oplus \mathfrak{s}_{+2}$$

- \mapsto $\mathfrak{s}_{\pm 2}$: central elements corresponding to the components of p_{μ}
- \mapsto $\mathfrak{s}_{\pm 1}$: generators of SUSY transformations
- \rightarrow 50: generators of global gauge transformations \equiv non-abelian R-symmetry group $SU(2)^{\times 4}$

Infinite (classical) symmetry algebra

$$\hat{\mathfrak{f}}^{\perp} = \operatorname{Ker}(\operatorname{ad}\Lambda) \cap \mathcal{L}(\mathfrak{psu}(2,2|4),\sigma)$$

Hollowood-JLM'11 Hoare-Tseytlin'11

The elements of grade ± 1 (or Lorentz spin $\pm 1/2$) generate SUSY transformations whose closure is the (exotic) $\mathcal{N}=(8|8)$ superalgebra

$$\mathfrak{s} = \big(\mathfrak{psu}(2|2) \oplus \mathfrak{psu}(2|2)\big) \ltimes \big(\mathbb{R} \oplus \mathbb{R}\big) = \mathfrak{s}_{-2} \oplus \mathfrak{s}_{-1} \oplus \mathfrak{s}_0 \oplus \mathfrak{s}_{+1} \oplus \mathfrak{s}_{+2}$$

- \mapsto 5±2: central elements corresponding to the components of p_{μ}
- → s_{±1}: generators of SUSY transformations
- \rightarrow 50: generators of global gauge transformations \equiv non-abelian R-symmetry group $SU(2)^{\times 4}$

Pirsa: 11080035 Page 29/166

Infinite (classical) symmetry algebra

$$\hat{\mathfrak{f}}^{\perp} = \operatorname{Ker}(\operatorname{ad}\Lambda) \cap \mathcal{L}(\mathfrak{psu}(2,2|4),\sigma)$$

Hoare-Tsevtlin'11

The elements of grade ± 1 (or Lorentz spin $\pm 1/2$) generate SUSY transformations whose closure is the (exotic) $\mathcal{N} = (8|8)$ superalgebra

$$\mathfrak{s} = \big(\mathfrak{psu}(2|2) \oplus \mathfrak{psu}(2|2)\big) \ltimes \big(\mathbb{R} \oplus \mathbb{R}\big) = \mathfrak{s}_{-2} \oplus \mathfrak{s}_{-1} \oplus \mathfrak{s}_0 \oplus \mathfrak{s}_{+1} \oplus \mathfrak{s}_{+2}$$

- \mapsto $\mathfrak{s}_{\pm 2}$: central elements corresponding to the components of p_{μ}
- \rightarrow 5+1: generators of SUSY transformations
- → 50: generators of global gauge transformations \equiv non-abelian R-symmetry group $SU(2)^{\times 4}$
- \mathfrak{s} is a finite subalgebra of $\mathcal{L}(\mathfrak{p}(\mathfrak{su}(2|2) \oplus \mathfrak{su}(2|2)), \sigma) \subset \hat{\mathfrak{f}}^{\perp}$

derivation zd/dz is the generator of Lorentz boosts

Infinite (classical) symmetry algebra

$$\hat{\mathfrak{f}}^{\perp} = \operatorname{Ker}(\operatorname{ad}\Lambda) \cap \mathcal{L}(\mathfrak{psu}(2,2|4),\sigma)$$

Hoare-Tseytlin'11

The elements of grade ± 1 (or Lorentz spin $\pm 1/2$) generate SUSY transformations whose closure is the (exotic) $\mathcal{N} = (8|8)$ superalgebra

$$\mathfrak{s} = \big(\mathfrak{psu}(2|2) \oplus \mathfrak{psu}(2|2)\big) \ltimes \big(\mathbb{R} \oplus \mathbb{R}\big) = \mathfrak{s}_{-2} \oplus \mathfrak{s}_{-1} \oplus \mathfrak{s}_0 \oplus \mathfrak{s}_{+1} \oplus \mathfrak{s}_{+2}$$

- \mapsto $\mathfrak{s}_{\pm 2}$: central elements corresponding to the components of p_{μ}
- \rightarrow 5+1: generators of SUSY transformations
- → 50: generators of global gauge transformations \equiv non-abelian R-symmetry group $SU(2)^{\times 4}$
- \mathfrak{s} is a finite subalgebra of $\mathcal{L}(\mathfrak{p}(\mathfrak{su}(2|2) \oplus \mathfrak{su}(2|2)), \sigma) \subset \hat{\mathfrak{f}}^{\perp}$ and the

derivation |zd/dz| is the generator of Lorentz boosts

The $AdS_5 \times S^5$ SSSSG theory

Goykhman-Ivanov'11 Hollowood-JLM'11

The SUSY transformations act in a mildly non-local way on the Lagrangian fields

The $AdS_5 \times S^5$ SSSSG theory

Goykhman-Ivanov'11 Hollowood-JLM'11

The SUSY transformations act in a mildly non-local way on the Lagrangian fields

Goykhman-Ivanov'11 Hollowood-JLM'11

- The SUSY transformations act in a mildly non-local way on the Lagrangian fields
 - \Rightarrow Conjecture: the symmetry algebra becomes q-deformed in the

quantum SSSSG theory

$$U_q(\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^2)$$

$$q = e^{i\pi/k}$$

Goykhman-Ivanov'11 Hollowood-JLM'11

- The SUSY transformations act in a mildly non-local way on the Lagrangian fields
 - \Rightarrow Conjecture: the symmetry algebra becomes q-deformed in the quantum SSSSG theory $U_q(\mathfrak{psu}(2|2)^2 \times \mathbb{R}^2)$ $q = e^{i\pi/k}$
- Different from the symmetry algebra of the superstring S-matrix

$$\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^3$$

Goykhman-Ivanov'11 Hollowood-JLM'11

- The SUSY transformations act in a mildly non-local way on the Lagrangian fields
 - \Rightarrow Conjecture: the symmetry algebra becomes q-deformed in the

quantum SSSSG theory

$$U_q(\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^2)$$

$$q = e^{i\pi/k}$$

- The SUSY transformations act in a mildly non-local way on the Lagrangian fields
 - \Rightarrow Conjecture: the symmetry algebra becomes q-deformed in the

quantum SSSSG theory

$$U_q(\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^2)$$

$$q = e^{i\pi/k}$$

- The SUSY transformations act in a mildly non-local way on the Lagrangian fields
 - \Rightarrow Conjecture: the symmetry algebra becomes q-deformed in the quantum SSSSG theory $U_q(\mathfrak{psu}(2|2)^2 \times \mathbb{R}^2)$ $q = e^{i\pi/k}$
- Different from the symmetry algebra of the superstring S-matrix

$$psu(2|2)^2 \times \mathbb{R}^3$$

- The SUSY transformations act in a mildly non-local way on the Lagrangian fields
 - \Rightarrow Conjecture: the symmetry algebra becomes q-deformed in the quantum SSSSG theory $U_q(\mathfrak{psu}(2|2)^2 \times \mathbb{R}^2)$ $q = e^{i\pi/k}$
- Different from the symmetry algebra of the superstring S-matrix

$$\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^3$$

- The SUSY transformations act in a mildly non-local way on the Lagrangian fields
 - \Rightarrow Conjecture: the symmetry algebra becomes q-deformed in the quantum SSSSG theory $U_q(\mathfrak{psu}(2|2)^2 \times \mathbb{R}^2)$ $q = e^{i\pi/k}$

$$psu(2|2)^2 \times \mathbb{R}^3$$

....although

$$\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^3 \subset \mathcal{L}(\mathfrak{p}(\mathfrak{su}(2|2) \oplus \mathfrak{su}(2|2)), \sigma) \oplus \mathbb{C} \mathfrak{C}$$

and both can be related to the same underlying loop algebra

- The SUSY transformations act in a mildly non-local way on the Lagrangian fields
 - \Rightarrow Conjecture: the symmetry algebra becomes q-deformed in the

$$U_q(\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^2)$$

$$q = e^{i\pi/k}$$

Different from the symmetry algebra of the superstring S-matrix

$$\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^3$$

....although

$$\mathfrak{psu}(2|2)^2\ltimes\mathbb{R}^3\subset\mathcal{L}\big(\mathfrak{p}(\mathfrak{su}(2|2)\oplus\mathfrak{su}(2|2)),\sigma\big)\oplus\mathbb{C}\,\mathfrak{C}$$

and both can be related to the same underlying loop algebra

Hallowood-JLM'11

Continuous spectrum of relativistic non-abelian Q-ball kinks with bosonic and fermionic degrees of freedom

labelled by
$$\varphi \in (0, \pi/2) \longrightarrow m_{\varphi} = \frac{2k}{\pi} \sin \varphi$$

Non-trivial moduli space
$$\mathfrak{M} = \frac{SU(2|2)}{U(2|1)} \times \frac{SU(2|2)}{U(2|1)}$$

Pirsa: 11080035 Page 42/166

Hollowood-JLM'11

Continuous spectrum of relativistic non-abelian Q-ball kinks with bosonic and fermionic degrees of freedom

labelled by
$$\varphi \in (0, \pi/2) \longrightarrow m_{\varphi} = \frac{2k}{\pi} \sin \varphi$$

Non-trivial moduli space
$$\mathfrak{M} = \frac{SU(2|2)}{U(2|1)} \times \frac{SU(2|2)}{U(2|1)}$$

Hollowood-JLM'11

Continuous spectrum of relativistic non-abelian Q-ball kinks with bosonic and fermionic degrees of freedom

labelled by
$$\varphi \in (0, \pi/2) \longrightarrow m_{\varphi} = \frac{2k}{\pi} \sin \varphi$$

Non-trivial moduli space

$$\mathfrak{M} = \frac{SU(2|2)}{U(2|1)} \times \frac{SU(2|2)}{U(2|1)}$$

Grassmann coordinates arise from the non-compact AdS₅ sector

 \longrightarrow turning off the Grassmann coordinates the solitons live in S^5

Hollowood-JLM'11

Continuous spectrum of relativistic non-abelian Q-ball kinks with bosonic and fermionic degrees of freedom

labelled by
$$\varphi \in (0, \pi/2) \longrightarrow m_{\varphi} = \frac{2k}{\pi} \sin \varphi$$

Non-trivial moduli space

$$\mathfrak{M} = \frac{SU(2|2)}{U(2|1)} \times \frac{SU(2|2)}{U(2|1)}$$

Grassmann coordinates arise from the non-compact AdS₅ sector

 \longrightarrow turning off the Grassmann coordinates the solitons live in S^5

Hollowood-ILM'11

Continuous spectrum of relativistic non-abelian Q-ball kinks with bosonic and fermionic degrees of freedom

labelled by
$$\varphi \in (0, \pi/2) \longrightarrow m_{\varphi} = \frac{2k}{\pi} \sin \varphi$$

Non-trivial moduli space
$$\mathfrak{M} = \frac{SU(2|2)}{U(2|1)} \times \frac{SU(2|2)}{U(2|1)}$$

- Grassmann coordinates arise from the non-compact AdS₅ sector
 - \rightarrow turning off the Grassmann coordinates the solitons live in S^5
- The SUSY group acts on M

Pirsa: 11080035 Page 46/166

Hollowood-JLM'11

Continuous spectrum of relativistic non-abelian Q-ball kinks with bosonic and fermionic degrees of freedom

labelled by
$$\varphi \in (0, \pi/2) \longrightarrow m_{\varphi} = \frac{2k}{\pi} \sin \varphi$$

Non-trivial moduli space

$$\mathfrak{M} = \frac{SU(2|2)}{U(2|1)} \times \frac{SU(2|2)}{U(2|1)}$$

- Grassmann coordinates arise from the non-compact AdS₅ sector
 - \longrightarrow turning off the Grassmann coordinates the solitons live in S^5

The SUSY group acts on M

The AdS₅ × S⁵ SSSSG theory

Quantization of moduli space dynamics

Semiclassical spectrum

Hilbert space of modules for the short (atypical) representations of the SUSY algebra of dimension $4a \times 4a$

Mass spectrum $m_a = \mu \sin \theta$

$$m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$$

$$a=1,\ldots,k$$

The AdS₅ × S⁵ SSSSG theory

Quantization of moduli space dynamics

Semiclassical spectrum

Hilbert space of modules for the short (atypical) representations of the SUSY algebra of dimension $4a \times 4a$

Mass spectrum $m_a = \mu \sin \left(\frac{1}{2} \right)$

$$m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$$

$$a=1,\ldots,k$$

Hallowood-IIM'11

Continuous spectrum of relativistic non-abelian Q-ball kinks with bosonic and fermionic degrees of freedom

labelled by
$$\varphi \in (0, \pi/2) \longrightarrow m_{\varphi} = \frac{2k}{\pi} \sin \varphi$$

Non-trivial moduli space
$$\mathfrak{M} = \frac{SU(2|2)}{U(2|1)} \times \frac{SU(2|2)}{U(2|1)}$$

- Grassmann coordinates arise from the non-compact AdS₅ sector
 - \longrightarrow turning off the Grassmann coordinates the solitons live in S^5
- The SUSY group acts on M

Hallawood-JLM'11

Continuous spectrum of relativistic non-abelian Q-ball kinks with bosonic and fermionic degrees of freedom

labelled by
$$\varphi \in (0, \pi/2) \longrightarrow m_{\varphi} = \frac{2k}{\pi} \sin \varphi$$

Non-trivial moduli space
$$\mathfrak{M} = \frac{SU(2|2)}{U(2|1)} \times \frac{SU(2|2)}{U(2|1)}$$

- Grassmann coordinates arise from the non-compact AdS₅ sector
 - \rightarrow turning off the Grassmann coordinates the solitons live in S^5
- The SUSY group acts on M

The AdS₅ × S⁵ SSSSG theory

Quantization of moduli space dynamics

Semiclassical spectrum

Hilbert space of modules for the short (atypical) representations of the SUSY algebra of dimension $4a \times 4a$

Mass spectrum
$$m_a = \mu \sin \left(\frac{1}{2} \right)$$

$$m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$$

$$a=1,\ldots,k$$

The AdS x S5 SSSSG theory

Quantization of moduli space dynamics

Semiclassical spectrum

Hilbert space of modules for the short (atypical) representations of the SUSY algebra of dimension $4a \times 4a$

Mass spectrum
$$m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$$

$$a=1,\ldots,k$$

Quantization of moduli space dynamics

Semiclassical spectrum

Hilbert space of modules for the short (atypical) representations of the SUSY algebra of dimension $4a \times 4a$

Mass spectrum
$$m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$$

$$a=1,\ldots,k$$

The semiclassical spectrum is discrete

a = 1 — perturbative states a > 1 — bound states

Quantization of moduli space dynamics

Semiclassical spectrum

Hilbert space of modules for the short (atypical) representations of the SUSY algebra of dimension $4a \times 4a$

Mass spectrum
$$m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$$

$$a=1,\ldots,k$$

The semiclassical spectrum is discrete

a = 1 — perturbative states a > 1 — bound states

The $AdS_5 \times S^5$ SSSSG theory

 \bigstar $m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$ can be understood as the BPS condition for the short representation of $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^2)$ of dimension $4a \times 4a$

with
$$q = e^{i\pi/k}$$

Quantization of moduli space dynamics

Semiclassical spectrum

Hilbert space of modules for the short (atypical) representations of the SUSY algebra of dimension $4a \times 4a$

Mass spectrum $m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$

$$a=1,\ldots,k$$

The semiclassical spectrum is discrete

 $a = 1 \longrightarrow \text{perturbative states}$

 $a > 1 \longrightarrow \text{bound states}$

Quantization of moduli space dynamics

Semiclassical spectrum

Hilbert space of modules for the short (atypical) representations of the SUSY algebra of dimension $4a \times 4a$

Mass spectrum $m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$

$$a=1,\ldots,k$$

The semiclassical spectrum is discrete

a = 1 \longrightarrow perturbative states a > 1 \longrightarrow bound states

The $AdS_5 \times S^5$ SSSSG theory

 \bigstar $m_a=\mu\sin\left(\frac{\pi a}{2k}\right)$ can be understood as the BPS condition for the short representation of $U_q(\mathfrak{psu}(2|2)\ltimes\mathbb{R}^2)$ of dimension $4a\times 4a$ with $q=e^{i\pi/k}$

The AdS₅ × S⁵ SSSSG theory

 $m_a = \mu \sin\left(\frac{\pi a}{2k}\right)$ can be understood as the BPS condition for the short representation of $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^2)$ of dimension $4a \times 4a$

with
$$q = e^{i\pi/k}$$

The $AdS_5 \times S^5$ SSSSG theory

 \bigstar $m_a=\mu\sin\left(\frac{\pi a}{2k}\right)$ can be understood as the BPS condition for the short representation of $U_q(\mathfrak{psu}(2|2)\ltimes\mathbb{R}^2)$ of dimension $4a\times 4a$ with $q=e^{i\pi/k}$

The semiclassical spectrum consists of a tower of massive states in short symmetric representations, just like the magnon bound states of the superstring world sheet theory.....

The AdS₅ × S⁵ SSSSG theory

 \bigstar $m_a=\mu\sin\left(\frac{\pi a}{2k}\right)$ can be understood as the BPS condition for the short representation of $U_q(\mathfrak{psu}(2|2)\ltimes\mathbb{R}^2)$ of dimension $4a\times 4a$ with $q=e^{i\pi/k}$

The semiclassical spectrum consists of a tower of massive states in short symmetric representations, just like the magnon bound states of the superstring world sheet theory.....

The AdS₅ × S⁵ SSSSG theory

 \bigstar $m_a=\mu\sin\left(\frac{\pi a}{2k}\right)$ can be understood as the BPS condition for the short representation of $U_q(\mathfrak{psu}(2|2)\ltimes\mathbb{R}^2)$ of dimension $4a\times 4a$ with $q=e^{i\pi/k}$

The semiclassical spectrum consists of a tower of massive states in short symmetric representations, just like the magnon bound states of the superstring world sheet theory......

but in the SSSG theory the tower is truncated by k, the level of the WZW term....

 \bigstar $m_a=\mu\sin\left(\frac{\pi a}{2k}\right)$ can be understood as the BPS condition for the short representation of $U_q(\mathfrak{psu}(2|2)\ltimes\mathbb{R}^2)$ of dimension $4a\times 4a$ with $q=e^{i\pi/k}$

The semiclassical spectrum consists of a tower of massive states in short symmetric representations, just like the magnon bound states of the superstring world sheet theory......

but in the SSSG theory the tower is truncated by k, the level of the WZW term....

Hollowood-JLM'11

Semiclassical quantization using the string world-sheet action gives rise to the usual infinite spectrum of magnon bound states in short symmetric representations!

 \bigstar $m_a=\mu\sin\left(\frac{\pi a}{2k}\right)$ can be understood as the BPS condition for the short representation of $U_q(\mathfrak{psu}(2|2)\ltimes\mathbb{R}^2)$ of dimension $4a\times 4a$ with $q=e^{i\pi/k}$

The semiclassical spectrum consists of a tower of massive states in short symmetric representations, just like the magnon bound states of the superstring world sheet theory.....

but in the SSSG theory the tower is truncated by k, the level of the WZW term....

Hollowood-JLM'11

Semiclassical quantization using the string world-sheet action gives rise to the usual infinite spectrum of magnon bound states in short symmetric representations!

• The SSSSG S-matrix with symmetry $U_q(\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^2)$ will be constructed as a graded tensor product of elementary blocks with symmetry $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^2)$

Hoare-Tseytlin'11

- Supported by perturbation theory
- The blocks are obtained as a limit of the fundamental R-matrix for $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^3)$ constructed by Beisert and Koroteev

• The SSSSG S-matrix with symmetry $U_q(\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^2)$ will be constructed as a graded tensor product of elementary blocks with symmetry $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^2)$

Hoare-Tseytlin'11

- Supported by perturbation theory
- The blocks are obtained as a limit of the fundamental R-matrix for $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^3)$ constructed by Beisert and Koroteev

• The SSSSG S-matrix with symmetry $U_q(\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^2)$ will be constructed as a graded tensor product of elementary blocks with symmetry $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^2)$

Hoare-Tseytlin'11

- Supported by perturbation theory
- The blocks are obtained as a limit of the fundamental R-matrix for $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^3)$ constructed by Beisert and Koroteev
- $\mathfrak{psu}(2|2) \ltimes \mathbb{R}^2$ is a "finite" affine algebra

$$\mathfrak{psu}(2|2) \ltimes \mathbb{R}^2 \subset \mathcal{L}(\mathfrak{su}(2|2), \sigma)$$

 \longrightarrow our relativistic S-matrix fits into the well known class of S-matrices associated to trigonometric solutions to the Yang-Baxter equation with affine quantum group $U_q(\hat{\mathfrak{g}})$ symmetry

• The SSSSG S-matrix with symmetry $U_q(\mathfrak{psu}(2|2)^2 \ltimes \mathbb{R}^2)$ will be constructed as a graded tensor product of elementary blocks with symmetry $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^2)$

- --- Supported by perturbation theory
- The blocks are obtained as a limit of the fundamental R-matrix for $U_q(\mathfrak{psu}(2|2) \ltimes \mathbb{R}^3)$ constructed by Beisert and Koroteev
- $\mathfrak{psu}(2|2)\ltimes\mathbb{R}^2$ is a "finite" affine algebra

$$\mathfrak{psu}(2|2) \ltimes \mathbb{R}^2 \subset \mathcal{L}(\mathfrak{su}(2|2), \sigma)$$

 \longrightarrow our relativistic S-matrix fits into the well known class of S-matrices associated to trigonometric solutions to the Yang-Baxter equation with affine quantum group $U_q(\hat{\mathfrak{g}})$ symmetry

Hoare-Tsevtlin'11

The extended sl(2|2) superalgebra

Generators

Even
$$\to \mathfrak{sl}(2)^2$$
: $\mathfrak{R}^a{}_b$, $\mathfrak{L}^\alpha{}_\beta$
Odd \to SUSY: $\mathfrak{Q}^\alpha{}_b$, $\mathfrak{S}^a{}_\beta$ Centres: \mathfrak{C} , \mathfrak{P} , \mathfrak{K}

• $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$ algebra

$$\begin{split} \left[\mathfrak{R}^{a}{}_{b},\mathfrak{R}^{c}{}_{d}\right] &= \delta^{c}_{b}\mathfrak{R}^{a}{}_{d} - \delta^{a}_{d}\mathfrak{R}^{c}{}_{b} \\ \left[\mathfrak{R}^{a}{}_{b},\mathfrak{Q}^{\gamma}{}_{d}\right] &= -\delta^{a}_{d}\mathfrak{Q}^{\gamma}{}_{b} + \frac{1}{2}\delta^{a}_{b}\mathfrak{Q}^{\gamma}{}_{d} \\ \left[\mathfrak{R}^{a}{}_{b},\mathfrak{Q}^{\gamma}{}_{d}\right] &= -\delta^{c}_{d}\mathfrak{Q}^{\gamma}{}_{b} + \frac{1}{2}\delta^{a}_{b}\mathfrak{Q}^{\gamma}{}_{d} \\ \left[\mathfrak{R}^{a}{}_{b},\mathfrak{G}^{c}{}_{\delta}\right] &= \delta^{c}_{b}\mathfrak{G}^{a}{}_{\delta} - \frac{1}{2}\delta^{a}_{b}\mathfrak{G}^{c}{}_{\delta} \\ \left[\mathfrak{L}^{\alpha}{}_{\beta},\mathfrak{Q}^{\gamma}{}_{d}\right] &= -\delta^{\alpha}_{\delta}\mathfrak{G}^{c}{}_{\beta} + \frac{1}{2}\delta^{\alpha}_{\beta}\mathfrak{Q}^{\gamma}{}_{d} \\ \left[\mathfrak{L}^{\alpha}{}_{\beta},\mathfrak{G}^{c}{}_{\delta}\right] &= -\delta^{\alpha}_{\delta}\mathfrak{G}^{c}{}_{\beta} + \frac{1}{2}\delta^{\alpha}_{\beta}\mathfrak{G}^{c}{}_{\delta} \end{split}$$

$$\begin{split} \{\mathfrak{Q}^{\alpha}{}_{b},\mathfrak{S}^{c}{}_{\delta}\} &= \delta^{c}_{b}\mathfrak{L}^{\alpha}{}_{\delta} + \delta^{\alpha}_{\delta}\mathfrak{R}^{c}{}_{b} + \delta^{c}_{b}\delta^{\alpha}_{\delta}\mathfrak{C} \\ \{\mathfrak{Q}^{\alpha}{}_{b},\mathfrak{Q}^{\gamma}{}_{d}\} &= \varepsilon^{\alpha\gamma}\varepsilon_{bd}\mathfrak{P} \qquad \{\mathfrak{S}^{a}{}_{\beta},\mathfrak{S}^{c}{}_{\delta}\} = \varepsilon^{ac}\varepsilon_{\beta\delta}\mathfrak{K} \end{split}$$

The extended sl(2|2) superalgebra

Generators

Even
$$\to \mathfrak{sl}(2)^2$$
: $\mathfrak{R}^a{}_b$, $\mathfrak{L}^\alpha{}_\beta$
Odd \to SUSY: $\mathfrak{Q}^\alpha{}_b$, $\mathfrak{S}^a{}_\beta$ Centres: \mathfrak{C} , \mathfrak{P} , \mathfrak{K}

• $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$ algebra

$$\begin{split} \left[\mathfrak{R}^{a}{}_{b},\mathfrak{R}^{c}{}_{d}\right] &= \delta^{c}_{b}\mathfrak{R}^{a}{}_{d} - \delta^{a}_{d}\mathfrak{R}^{c}{}_{b} \\ \left[\mathfrak{R}^{a}{}_{b},\mathfrak{Q}^{\gamma}{}_{d}\right] &= -\delta^{a}_{d}\mathfrak{Q}^{\gamma}{}_{b} + \frac{1}{2}\delta^{a}_{b}\mathfrak{Q}^{\gamma}{}_{d} \\ \left[\mathfrak{R}^{a}{}_{b},\mathfrak{Q}^{\gamma}{}_{d}\right] &= -\delta^{c}_{d}\mathfrak{Q}^{\gamma}{}_{b} + \frac{1}{2}\delta^{a}_{b}\mathfrak{Q}^{\gamma}{}_{d} \\ \left[\mathfrak{R}^{a}{}_{b},\mathfrak{G}^{c}{}_{\delta}\right] &= \delta^{c}_{b}\mathfrak{S}^{a}{}_{\delta} - \frac{1}{2}\delta^{a}_{b}\mathfrak{S}^{c}{}_{\delta} \\ \left[\mathfrak{L}^{\alpha}{}_{\beta},\mathfrak{Q}^{\gamma}{}_{d}\right] &= -\delta^{\alpha}_{\delta}\mathfrak{Q}^{c}{}_{\delta} + \frac{1}{2}\delta^{\alpha}_{\beta}\mathfrak{Q}^{\gamma}{}_{d} \\ \left[\mathfrak{L}^{\alpha}{}_{\beta},\mathfrak{S}^{c}{}_{\delta}\right] &= -\delta^{\alpha}_{\delta}\mathfrak{S}^{c}{}_{\beta} + \frac{1}{2}\delta^{\alpha}_{\beta}\mathfrak{S}^{c}{}_{\delta} \end{split}$$

$$\begin{split} \{\mathfrak{Q}^{\alpha}{}_{b},\mathfrak{S}^{c}{}_{\delta}\} &= \delta^{c}_{b}\mathfrak{L}^{\alpha}{}_{\delta} + \delta^{\alpha}_{\delta}\mathfrak{R}^{c}{}_{b} + \delta^{c}_{b}\delta^{\alpha}_{\delta}\mathfrak{C} \\ \{\mathfrak{Q}^{\alpha}{}_{b},\mathfrak{Q}^{\gamma}{}_{d}\} &= \varepsilon^{\alpha\gamma}\varepsilon_{bd}\mathfrak{P} \qquad \{\mathfrak{S}^{a}{}_{\beta},\mathfrak{S}^{c}{}_{\delta}\} = \varepsilon^{ac}\varepsilon_{\beta\delta}\mathfrak{K} \end{split}$$

Defining representation(s)

- psl(2|2) $\times \mathbb{R}^3$ has a family of 4-dimensional representations labelled by four parameters a, b, c, d constrained by ad bc = 1
 - Odd generators

$$\mathfrak{Q}^{1}{}_{1} = a \, \mathbf{e}_{31} + b \, \mathbf{e}_{24}, \qquad \mathfrak{Q}^{1}{}_{2} = a \, \mathbf{e}_{32} - b \, \mathbf{e}_{14},
\mathfrak{Q}^{2}{}_{1} = a \, \mathbf{e}_{41} - b \, \mathbf{e}_{23}, \qquad \mathfrak{Q}^{2}{}_{2} = a \, \mathbf{e}_{42} + b \, \mathbf{e}_{13},
\mathfrak{S}^{1}{}_{1} = d \, \mathbf{e}_{13} + c \, \mathbf{e}_{42}, \qquad \mathfrak{S}^{1}{}_{2} = d \, \mathbf{e}_{14} - c \, \mathbf{e}_{32},
\mathfrak{S}^{2}{}_{1} = d \, \mathbf{e}_{23} - c \, \mathbf{e}_{41}, \qquad \mathfrak{S}^{2}{}_{2} = d \, \mathbf{e}_{24} + c \, \mathbf{e}_{31},$$

• Centres $\mathfrak{C} = C \mathbf{1}, \ \mathfrak{P} = P \mathbf{1}, \ \mathfrak{K} = K \mathbf{1}$

$$C = \frac{1}{2}(ad + bc), \quad P = ab, \quad K = cd$$

 $\Rightarrow C^2 - PK = \frac{1}{4} \equiv \text{ shortening (BPS) condition}$

Defining representation(s)

- psl(2|2) $\times \mathbb{R}^3$ has a family of 4-dimensional representations labelled by four parameters a, b, c, d constrained by ad bc = 1
 - Odd generators

$$\mathfrak{Q}^{1}{}_{1} = a \, \mathbf{e}_{31} + b \, \mathbf{e}_{24}, \qquad \mathfrak{Q}^{1}{}_{2} = a \, \mathbf{e}_{32} - b \, \mathbf{e}_{14},
\mathfrak{Q}^{2}{}_{1} = a \, \mathbf{e}_{41} - b \, \mathbf{e}_{23}, \qquad \mathfrak{Q}^{2}{}_{2} = a \, \mathbf{e}_{42} + b \, \mathbf{e}_{13}
\mathfrak{S}^{1}{}_{1} = d \, \mathbf{e}_{13} + c \, \mathbf{e}_{42}, \qquad \mathfrak{S}^{1}{}_{2} = d \, \mathbf{e}_{14} - c \, \mathbf{e}_{32},
\mathfrak{S}^{2}{}_{1} = d \, \mathbf{e}_{23} - c \, \mathbf{e}_{41}, \qquad \mathfrak{S}^{2}{}_{2} = d \, \mathbf{e}_{24} + c \, \mathbf{e}_{31},$$

• Centres $\mathfrak{C} = C \mathbf{1}, \ \mathfrak{P} = P \mathbf{1}, \ \mathfrak{K} = K \mathbf{1}$

$$C = \frac{1}{2}(ad + bc), \quad P = ab, \quad K = cd$$

$$\Rightarrow$$
 $C^2 - PK = \frac{1}{4}$ \equiv shortening (BPS) condition

\mathbb{Z} -gradation and \mathbb{Z}_4 automorphism

• \mathbb{Z} -gradation of $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$

$$s(\mathfrak{R}^{a}{}_{b}) = s(\mathfrak{L}^{\alpha}{}_{\beta}) = 0,$$
 $s(\mathfrak{Q}^{\alpha}{}_{b}) = 1,$ $s(\mathfrak{S}^{a}{}_{\beta}) = -1,$ $s(\mathfrak{C}) = 0,$ $s(\mathfrak{P}) = 2,$ $s(\mathfrak{K}) = -2,$

• \mathbb{Z}_4 automorphism of $\mathfrak{sl}(2|2)$: $\mathfrak{L}^{\alpha}{}_{\beta}$, $\mathfrak{R}^{a}{}_{b}$, $\mathfrak{Q}^{\alpha}{}_{b}$, $\mathfrak{S}^{a}{}_{\beta}$, $\mathfrak{1}$

$$\sigma(u) = e^{i\pi s(u)/2}u, \quad \sigma(1) = -1$$

psl(2|2) $\times \mathbb{R}^3$ is a finite subalgebra of the centrally extended twisted affine algebra

$$\mathfrak{sl}(2|2)^{(\sigma)}=\mathcal{L}\big(\mathfrak{sl}(2|2),\sigma\big)\oplus\mathbb{C}\,\mathfrak{C}$$

$$\mathfrak{P} \equiv z^2 \otimes \mathbf{1}, \quad \mathfrak{K} \equiv z^{-2} \otimes \mathbf{1}$$

Defining representation(s)

- psl(2|2) $\times \mathbb{R}^3$ has a family of 4-dimensional representations labelled by four parameters a, b, c, d constrained by ad bc = 1
 - Odd generators

$$\mathfrak{Q}^{1}{}_{1} = a \, \mathbf{e}_{31} + b \, \mathbf{e}_{24}, \qquad \mathfrak{Q}^{1}{}_{2} = a \, \mathbf{e}_{32} - b \, \mathbf{e}_{14},
\mathfrak{Q}^{2}{}_{1} = a \, \mathbf{e}_{41} - b \, \mathbf{e}_{23}, \qquad \mathfrak{Q}^{2}{}_{2} = a \, \mathbf{e}_{42} + b \, \mathbf{e}_{13}
\mathfrak{S}^{1}{}_{1} = d \, \mathbf{e}_{13} + c \, \mathbf{e}_{42}, \qquad \mathfrak{S}^{1}{}_{2} = d \, \mathbf{e}_{14} - c \, \mathbf{e}_{32},
\mathfrak{S}^{2}{}_{1} = d \, \mathbf{e}_{23} - c \, \mathbf{e}_{41}, \qquad \mathfrak{S}^{2}{}_{2} = d \, \mathbf{e}_{24} + c \, \mathbf{e}_{31},$$

• Centres $\mathfrak{C} = C \mathbf{1}, \ \mathfrak{P} = P \mathbf{1}, \ \mathfrak{K} = K \mathbf{1}$

$$C = \frac{1}{2}(ad + bc), \quad P = ab, \quad K = cd$$

 $\Rightarrow C^2 - PK = \frac{1}{4} \equiv \text{ shortening (BPS) condition}$

\mathbb{Z} -gradation and \mathbb{Z}_4 automorphism

• \mathbb{Z} -gradation of $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$

$$s(\mathfrak{R}^{a}{}_{b}) = s(\mathfrak{L}^{\alpha}{}_{\beta}) = 0,$$
 $s(\mathfrak{Q}^{\alpha}{}_{b}) = 1,$ $s(\mathfrak{S}^{a}{}_{\beta}) = -1,$ $s(\mathfrak{C}) = 0,$ $s(\mathfrak{P}) = 2,$ $s(\mathfrak{K}) = -2,$

• \mathbb{Z}_4 automorphism of $\mathfrak{sl}(2|2)$: $\mathfrak{L}^{\alpha}{}_{\beta}$, $\mathfrak{R}^{a}{}_{b}$, $\mathfrak{Q}^{\alpha}{}_{b}$, $\mathfrak{S}^{a}{}_{\beta}$, $\mathfrak{1}$

$$\sigma(u) = e^{i\pi s(u)/2}u, \quad \sigma(1) = -1$$

psl(2|2) $\times \mathbb{R}^3$ is a finite subalgebra of the centrally extended twisted affine algebra

$$\mathfrak{sl}(2|2)^{(\sigma)}=\mathcal{L}\big(\mathfrak{sl}(2|2),\sigma\big)\oplus\mathbb{C}\,\mathfrak{C}$$

$$\mathfrak{P} \equiv z^2 \otimes \mathbf{1}, \quad \mathfrak{K} \equiv z^{-2} \otimes \mathbf{1}$$

The extended sl(2|2) superalgebra

Generators

Even
$$\to \mathfrak{sl}(2)^2$$
: $\mathfrak{R}^a{}_b$, $\mathfrak{L}^\alpha{}_\beta$
Odd \to SUSY: $\mathfrak{Q}^\alpha{}_b$, $\mathfrak{S}^a{}_\beta$ Centres: \mathfrak{C} , \mathfrak{P} , \mathfrak{K}

• $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$ algebra

$$\begin{split} \left[\mathfrak{R}^{a}{}_{b},\mathfrak{R}^{c}{}_{d}\right] &= \delta^{c}_{b}\mathfrak{R}^{a}{}_{d} - \delta^{a}_{d}\mathfrak{R}^{c}{}_{b} \\ \left[\mathfrak{R}^{a}{}_{b},\mathfrak{Q}^{\gamma}{}_{d}\right] &= -\delta^{a}_{d}\mathfrak{Q}^{\gamma}{}_{b} + \frac{1}{2}\delta^{a}_{b}\mathfrak{Q}^{\gamma}{}_{d} \\ \left[\mathfrak{R}^{a}{}_{b},\mathfrak{Q}^{\gamma}{}_{d}\right] &= -\delta^{c}_{d}\mathfrak{Q}^{\gamma}{}_{b} + \frac{1}{2}\delta^{a}_{b}\mathfrak{Q}^{\gamma}{}_{d} \\ \left[\mathfrak{R}^{a}{}_{b},\mathfrak{G}^{c}{}_{\delta}\right] &= \delta^{c}_{b}\mathfrak{S}^{a}{}_{\delta} - \frac{1}{2}\delta^{a}_{b}\mathfrak{S}^{c}{}_{\delta} \\ \left[\mathfrak{L}^{\alpha}{}_{\beta},\mathfrak{Q}^{\gamma}{}_{d}\right] &= -\delta^{\alpha}_{\delta}\mathfrak{Q}^{\alpha}{}_{d} - \frac{1}{2}\delta^{\alpha}_{\beta}\mathfrak{Q}^{\gamma}{}_{d} \\ \left[\mathfrak{L}^{\alpha}{}_{\beta},\mathfrak{S}^{c}{}_{\delta}\right] &= -\delta^{\alpha}_{\delta}\mathfrak{S}^{c}{}_{\beta} + \frac{1}{2}\delta^{\alpha}_{\beta}\mathfrak{S}^{c}{}_{\delta} \end{split}$$

$$\begin{split} \{\mathfrak{Q}^{\alpha}{}_{b},\mathfrak{S}^{c}{}_{\delta}\} &= \delta^{c}_{b}\mathfrak{L}^{\alpha}{}_{\delta} + \delta^{\alpha}_{\delta}\mathfrak{R}^{c}{}_{b} + \delta^{c}_{b}\delta^{\alpha}_{\delta}\,\mathfrak{C} \\ \{\mathfrak{Q}^{\alpha}{}_{b},\mathfrak{Q}^{\gamma}{}_{d}\} &= \varepsilon^{\alpha\gamma}\varepsilon_{bd}\,\mathfrak{P} \qquad \{\mathfrak{S}^{a}{}_{\beta},\mathfrak{S}^{c}{}_{\delta}\} = \varepsilon^{ac}\varepsilon_{\beta\delta}\,\mathfrak{K} \end{split}$$

\mathbb{Z} -gradation and \mathbb{Z}_4 automorphism

• \mathbb{Z} -gradation of $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$

$$s(\mathfrak{R}^{a}{}_{b}) = s(\mathfrak{L}^{\alpha}{}_{\beta}) = 0,$$
 $s(\mathfrak{Q}^{\alpha}{}_{b}) = 1,$ $s(\mathfrak{S}^{a}{}_{\beta}) = -1,$ $s(\mathfrak{C}) = 0,$ $s(\mathfrak{P}) = 2,$ $s(\mathfrak{K}) = -2,$

• \mathbb{Z}_4 automorphism of $\mathfrak{sl}(2|2)$: $\mathfrak{L}^{\alpha}{}_{\beta}$, $\mathfrak{R}^{a}{}_{b}$, $\mathfrak{Q}^{\alpha}{}_{b}$, $\mathfrak{S}^{a}{}_{\beta}$, 1

$$\sigma(u) = e^{i\pi s(u)/2}u, \quad \sigma(1) = -1$$

 $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$ is a finite subalgebra of the centrally extended twisted affine algebra

$$\mathfrak{sl}(2|2)^{(\sigma)} = \mathcal{L}(\mathfrak{sl}(2|2), \sigma) \oplus \mathbb{C}\mathfrak{C}$$
 $\mathfrak{P} \equiv z^2 \otimes 1, \quad \mathfrak{K} \equiv z^{-2} \otimes 1$

$$\mathfrak{P} \equiv z^2 \otimes \mathbf{1}, \quad \mathfrak{K} \equiv z^{-2} \otimes \mathbf{1}$$

\mathbb{Z} -gradation and \mathbb{Z}_4 automorphism

• \mathbb{Z} -gradation of $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$

$$s(\mathfrak{R}^{a}{}_{b}) = s(\mathfrak{L}^{\alpha}{}_{\beta}) = 0,$$
 $s(\mathfrak{Q}^{\alpha}{}_{b}) = 1,$ $s(\mathfrak{S}^{a}{}_{\beta}) = -1,$ $s(\mathfrak{C}) = 0,$ $s(\mathfrak{P}) = 2,$ $s(\mathfrak{R}) = -2,$

• \mathbb{Z}_4 automorphism of $\mathfrak{sl}(2|2)$: $\mathfrak{L}^{\alpha}{}_{\beta}$, $\mathfrak{R}^{a}{}_{b}$, $\mathfrak{Q}^{\alpha}{}_{b}$, $\mathfrak{S}^{a}{}_{\beta}$, 1

$$\sigma(u) = e^{i\pi s(u)/2}u, \quad \sigma(1) = -1$$

psl(2|2) \times \mathbb{R}^3 is a finite subalgebra of the centrally extended twisted affine algebra

$$\mathfrak{sl}(2|2)^{(\sigma)} = \mathcal{L}(\mathfrak{sl}(2|2), \sigma) \oplus \mathbb{C}\mathfrak{C}$$

$$\mathfrak{P} \equiv z^2 \otimes \mathbf{1}, \quad \mathfrak{K} \equiv z^{-2} \otimes \mathbf{1}$$

Quantum deformation $U_q(\mathfrak{psl}(2|2)\ltimes\mathbb{R}^3)$


Chevalley generators

$$[\mathfrak{E}_1,\mathfrak{F}_1] = [\mathfrak{H}_1]_q, \qquad \{\mathfrak{E}_2,\mathfrak{F}_2\} = -[\mathfrak{H}_2]_q, \qquad [\mathfrak{E}_3,\mathfrak{F}_3] = -[\mathfrak{H}_3]_q$$

$$q^{\mathfrak{H}_i}\mathfrak{E}_j = q^{A_{ij}}\mathfrak{E}_jq^{\mathfrak{H}_i}, \qquad q^{\mathfrak{H}_i}\mathfrak{F}_j = q^{-A_{ij}}\mathfrak{F}_jq^{\mathfrak{H}_i}$$

+ deformed Serre relations

• Cartan matrix: $A_{ij} = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & -2 \end{pmatrix}$ $[x]_q = \frac{q^x - q^{-x}}{q - q^{-1}}$



Quantum deformation $U_a(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3)$

Chevalley generators

$$[\mathfrak{E}_{1},\mathfrak{F}_{1}] = [\mathfrak{H}_{1}]_{q}, \qquad \{\mathfrak{E}_{2},\mathfrak{F}_{2}\} = -[\mathfrak{H}_{2}]_{q}, \qquad [\mathfrak{E}_{3},\mathfrak{F}_{3}] = -[\mathfrak{H}_{3}]_{q}$$

$$q^{\mathfrak{H}_{i}}\mathfrak{E}_{j} = q^{A_{ij}}\mathfrak{E}_{j}q^{\mathfrak{H}_{i}}, \qquad q^{\mathfrak{H}_{i}}\mathfrak{F}_{j} = q^{-A_{ij}}\mathfrak{F}_{j}q^{\mathfrak{H}_{i}}$$

+ deformed Serre relations

• Cartan matrix: $A_{ij} = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & -2 \end{pmatrix}$ $[x]_q = \frac{q^x - q^{-x}}{q - q^{-1}}$



$$q = e^{i\pi/k}$$

Chevalley generators in the fundamental representation(s)

$$\mathfrak{E}_1 = q^{1/2} \, \mathfrak{R}^2_1, \qquad \mathfrak{E}_2 = \mathfrak{Q}^2_2, \qquad \mathfrak{E}_3 = q^{-1/2} \, \mathfrak{L}^1_2$$

 $\mathfrak{F}_1 = q^{-1/2} \, \mathfrak{R}^1_2, \qquad \mathfrak{F}_2 = \mathfrak{S}^2_2, \qquad \mathfrak{F}_3 = q^{1/2} \, \mathfrak{L}^2_1$

Centres

$$P = ab$$
, $K = cd$, $ad = [C + 1/2]_q$, $bc = [C - 1/2]_q$

$$\Rightarrow$$
 $[C]_q^2 - PK = [1/2]_q^2 \equiv$ deformed shortening condition

Chevalley generators in the fundamental representation(s)

$$\mathfrak{E}_1 = q^{1/2} \, \mathfrak{R}^2_{\ 1}, \qquad \mathfrak{E}_2 = \mathfrak{Q}^2_{\ 2}, \qquad \mathfrak{E}_3 = q^{-1/2} \, \mathfrak{L}^1_{\ 2}$$
 $\mathfrak{F}_1 = q^{-1/2} \, \mathfrak{R}^1_{\ 2}, \qquad \mathfrak{F}_2 = \mathfrak{S}^2_{\ 2}, \qquad \mathfrak{F}_3 = q^{1/2} \, \mathfrak{L}^2_{\ 1}$

Centres

$$P = ab$$
, $K = cd$, $ad = [C + 1/2]_q$, $bc = [C - 1/2]_q$

$$\Rightarrow$$
 $[C]_q^2 - PK = [1/2]_q^2 \equiv \text{deformed shortening condition}$

Beisert-Koroteev'08

Beisert-Koroteev parameterization of {a, b, c, d}

$$a = \sqrt{g}\gamma, \qquad b = \frac{\sqrt{g}\alpha}{\gamma} \left(1 - q^{2C-1} \frac{x^+}{x^-}\right),$$

$$c = \frac{i\sqrt{g}\gamma}{\alpha} \frac{q^{-C+1/2}}{x^+}, \quad d = \frac{i\sqrt{g}}{\gamma} q^{C+1/2} \left(x^- - q^{-2C-1}x^+\right)$$

subject to

$$\frac{x^{+}}{q} + \frac{q}{x^{+}} - qx^{-} - \frac{1}{qx^{-}} + ig(q - q^{-1})\left(\frac{x^{+}}{qx^{-}} - \frac{qx^{-}}{x^{+}}\right) = \frac{i}{g}$$



$$(g,q) \equiv \text{coupling constants} \qquad (x^+,x^-) \equiv 1 \text{ variable}$$

$$(x^+, x^-) \equiv 1$$
 variable

 $(\alpha, \gamma) \equiv$ normalization factors

Beisert-Koroteev'08

Beisert-Koroteev parameterization of {a, b, c, d}

$$a = \sqrt{g}\gamma, \qquad b = \frac{\sqrt{g}\alpha}{\gamma} \left(1 - q^{2C-1} \frac{x^+}{x^-}\right),$$

$$c = \frac{i\sqrt{g}\gamma}{\alpha} \frac{q^{-C+1/2}}{x^+}, \quad d = \frac{i\sqrt{g}}{\gamma} q^{C+1/2} \left(x^- - q^{-2C-1}x^+\right)$$

subject to

$$\frac{x^{+}}{q} + \frac{q}{x^{+}} - qx^{-} - \frac{1}{qx^{-}} + ig(q - q^{-1})\left(\frac{x^{+}}{qx^{-}} - \frac{qx^{-}}{x^{+}}\right) = \frac{i}{g}$$



$$(g,q) \equiv \text{coupling constants} \quad (x^+,x^-) \equiv 1 \text{ variable}$$

$$(x^+, x^-) \equiv 1$$
 variable

$$(\alpha, \gamma) \equiv$$
 normalization factors

Magnon representation:

$$q \to 1$$
 $(\text{or } k \to \infty) \Rightarrow \frac{x^+}{x^-} = e^{ip}$

 $p \equiv \text{world-sheet momentum}$ $g \equiv \text{string tension}$

$$[C]_q^2 - PK = [1/2]_q^2 \longrightarrow \epsilon(p) = \sqrt{1 + 4g^2 \sin^2(p/2)}$$

Magnon representation:

$$q \to 1$$
 $(\text{or } k \to \infty) \Rightarrow \frac{x^+}{x^-} = e^{ip}$

 $p \equiv \text{world-sheet momentum}$ $g \equiv \text{string tension}$

$$[C]_q^2 - PK = [1/2]_q^2 \longrightarrow \epsilon(p) = \sqrt{1 + 4g^2 \sin^2(p/2)}$$

Magnon representation:

$$q \to 1$$
 $(\text{or } k \to \infty) \Rightarrow \frac{x^+}{x^-} = e^{ip}$

 $p \equiv \text{world-sheet momentum}$ $g \equiv \text{string tension}$

$$[C]_q^2 - PK = [1/2]_q^2 \longrightarrow \epsilon(p) = \sqrt{1 + 4g^2 \sin^2(p/2)}$$

Hoare-Hollowood-JLM'11

$$g \to \infty$$
 \Rightarrow $\frac{x^+}{x^-} = q$, $C = 0$

Pirsa: 11080035
$$\begin{bmatrix} C \end{bmatrix}_q^2 - PK = \begin{bmatrix} 1/2 \end{bmatrix}_q^2$$
 becomes a relativistic mass-shell condition

Magnon representation:

$$q \to 1$$
 $(\text{or } k \to \infty) \Rightarrow \frac{x^+}{x^-} = e^{ip}$

 $p \equiv \text{world-sheet momentum}$ $g \equiv \text{string tension}$

$$[C]_q^2 - PK = [1/2]_q^2 \longrightarrow \epsilon(p) = \sqrt{1 + 4g^2 \sin^2(p/2)}$$

Hoare-Hollowood-JLM'11

$$g \to \infty$$
 \Rightarrow $\frac{x^+}{x^-} = q$, $C = 0$

Pirsa: 11080035
$$\left[C\right]_q^2 - PK = \left[1/2\right]_q^2$$
 becomes a relativistic mass-shell condition

Magnon representation:

$$\boxed{q \to 1} \quad (\text{or } k \to \infty) \Rightarrow \qquad \frac{x^+}{x^-} = e^{ip}$$

 $p \equiv \text{world-sheet momentum}$ $g \equiv \text{string tension}$

$$[C]_q^2 - PK = [1/2]_q^2 \longrightarrow \epsilon(p) = \sqrt{1 + 4g^2 \sin^2(p/2)}$$

Hoare-Hollowood-JLM'11

$$g \to \infty$$
 \Rightarrow $\frac{x^+}{x^-} = q$, $C = 0$

Pirsa: 11080035
$$\begin{bmatrix} C \end{bmatrix}_q^2 - PK = \begin{bmatrix} 1/2 \end{bmatrix}_q^2$$
 becomes a relativistic mass-shell condition

• Magnon representation:

$$\boxed{q \to 1} \quad (\text{or } k \to \infty) \Rightarrow \boxed{\frac{x^+}{x^-}} = e^{ip}$$

 $p \equiv \text{world-sheet momentum}$ $g \equiv \text{string tension}$

$$[C]_q^2 - PK = [1/2]_q^2 \longrightarrow \epsilon(p) = \sqrt{1 + 4g^2 \sin^2(p/2)}$$

Hoare-Hollowood-JLM'11

$$g \to \infty$$
 \Rightarrow $\frac{x^+}{x^-} = q$, $C = 0$

Pirsa: 11080035
$$\left[C\right]_q^2 - PK = \left[1/2\right]_q^2$$
 becomes a relativistic mass-shell condition

•
$$P = i [1/2]_q e^{-\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_-$$

$$K = i [1/2]_q e^{+\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_+$$

$$-PK = [1/2]_q^2 \Rightarrow p_+ p_- = \mu^2 \sin^2(\pi/2k) = m_1^2$$

○ The rapidity appears in the way dictated by the Z-gradation!

•
$$P = i [1/2]_q e^{-\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_-$$

$$K = i [1/2]_q e^{+\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_+$$

$$-PK = [1/2]_q^2 \Rightarrow p_+ p_- = \mu^2 \sin^2(\pi/2k) = m_1^2$$

○ The rapidity appears in the way dictated by the Z-gradation!

•
$$P = i [1/2]_q e^{-\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_-$$

 $K = i [1/2]_q e^{+\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_+$

$$-PK = [1/2]_q^2 \Rightarrow p_+ p_- = \mu^2 \sin^2(\pi/2k) = m_1^2$$

 \odot The rapidity appears in the way dictated by the \mathbb{Z} -gradation!

$$\star U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3) \longrightarrow U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^2)$$

•
$$P = i [1/2]_q e^{-\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_-$$

$$K = i [1/2]_q e^{+\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_+$$

$$-PK = [1/2]_q^2 \Rightarrow p_+ p_- = \mu^2 \sin^2(\pi/2k) = m_1^2$$

 \odot The rapidity appears in the way dictated by the \mathbb{Z} -gradation!

$$\star U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3) \longrightarrow U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^2)$$

•
$$P = i [1/2]_q e^{-\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_-$$

 $K = i [1/2]_q e^{+\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_+$

$$-PK = [1/2]_q^2 \Rightarrow p_+ p_- = \mu^2 \sin^2(\pi/2k) = m_1^2$$

 \odot The rapidity appears in the way dictated by the \mathbb{Z} -gradation!

$$\star U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3) \longrightarrow U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^2)$$

Fundamental particle multiplet $V_1(\theta)$ of mass

$$m_1 = \mu \sin(\pi/2k)$$
 \longrightarrow lightest semiclassical solitons $m_1 \simeq \mu + O(1/k)$ \longrightarrow perturbative modes

•
$$P = i [1/2]_q e^{-\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_-$$

$$K = i [1/2]_q e^{+\theta} \equiv \frac{i}{\mu \sin(\pi/k)} p_+$$

$$-PK = [1/2]_q^2 \Rightarrow p_+ p_- = \mu^2 \sin^2(\pi/2k) = m_1^2$$

 \odot The rapidity appears in the way dictated by the \mathbb{Z} -gradation!

$$\star U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3) \longrightarrow U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^2)$$

Fundamental particle multiplet $V_1(\theta)$ of mass

$$m_1 = \mu \sin(\pi/2k)$$
 \longrightarrow lightest semiclassical solitons $m_1 \simeq \mu + O(1/k)$ \longrightarrow perturbative modes

Beisert-Koroteev'08

Involves a new abelian generator

$$\Delta(\mathfrak{E}_{j}) = \mathfrak{E}_{j} \otimes 1 + q^{-\mathfrak{H}_{j}} \mathfrak{U}^{s_{j}} \otimes \mathfrak{E}_{j}, \quad \Delta(\mathfrak{F}_{j}) = \mathfrak{F}_{j} \otimes q^{\mathfrak{H}_{j}} + \mathfrak{U}^{-s_{j}} \otimes \mathfrak{F}_{j}$$

$$\Delta(\mathfrak{C}) = \mathfrak{C} \otimes 1 + 1 \otimes \mathfrak{C}, \qquad \Delta(\mathfrak{P}) = \mathfrak{P} \otimes 1 + q^{2\mathfrak{C}} \mathfrak{U}^{2} \otimes \mathfrak{P}$$

$$\Delta(\mathfrak{K}) = \mathfrak{K} \otimes q^{-2\mathfrak{C}} + \mathfrak{U}^{-2} \otimes \mathfrak{K}, \qquad \Delta(\mathfrak{U}) = \mathfrak{U} \otimes \mathfrak{U}$$

 $\star \mathbb{Z}$ gradation $(s_1, s_2, s_3) = (0, 1, 0)$

•
$$\mathfrak{P} = g\alpha(1 - q^{2\mathfrak{C}}\mathfrak{U}^2)$$
, $\mathfrak{K} = g\alpha^{-1}(q^{-2\mathfrak{C}} - \mathfrak{U}^{-2})$, $\mathfrak{U}^2 = x^+/qx^- \cdot 1$

Beisert-Koroteev'08

Involves a new abelian generator

$$\Delta(\mathfrak{E}_{j}) = \mathfrak{E}_{j} \otimes 1 + q^{-\mathfrak{H}_{j}} \mathfrak{U}^{s_{j}} \otimes \mathfrak{E}_{j}, \quad \Delta(\mathfrak{F}_{j}) = \mathfrak{F}_{j} \otimes q^{\mathfrak{H}_{j}} + \mathfrak{U}^{-s_{j}} \otimes \mathfrak{F}_{j}$$

$$\Delta(\mathfrak{C}) = \mathfrak{C} \otimes 1 + 1 \otimes \mathfrak{C}, \qquad \Delta(\mathfrak{P}) = \mathfrak{P} \otimes 1 + q^{2\mathfrak{C}} \mathfrak{U}^{2} \otimes \mathfrak{P}$$

$$\Delta(\mathfrak{K}) = \mathfrak{K} \otimes q^{-2\mathfrak{C}} + \mathfrak{U}^{-2} \otimes \mathfrak{K}, \qquad \Delta(\mathfrak{U}) = \mathfrak{U} \otimes \mathfrak{U}$$

 $\star \mathbb{Z}$ gradation $(s_1, s_2, s_3) = (0, 1, 0)$

•
$$\mathfrak{P} = g\alpha(1 - q^{2\mathfrak{C}}\mathfrak{U}^2)$$
, $\mathfrak{K} = g\alpha^{-1}(q^{-2\mathfrak{C}} - \mathfrak{U}^{-2})$, $\mathfrak{U}^2 = x^+/qx^- \cdot 1$

Beisert-Koroteev'08

Involves a new abelian generator !!

$$\Delta(\mathfrak{E}_{j}) = \mathfrak{E}_{j} \otimes 1 + q^{-\mathfrak{H}_{j}} \mathfrak{U}^{s_{j}} \otimes \mathfrak{E}_{j}, \quad \Delta(\mathfrak{F}_{j}) = \mathfrak{F}_{j} \otimes q^{\mathfrak{H}_{j}} + \mathfrak{U}^{-s_{j}} \otimes \mathfrak{F}_{j}$$

$$\Delta(\mathfrak{C}) = \mathfrak{C} \otimes 1 + 1 \otimes \mathfrak{C}, \qquad \Delta(\mathfrak{P}) = \mathfrak{P} \otimes 1 + q^{2\mathfrak{C}} \mathfrak{U}^{2} \otimes \mathfrak{P}$$

$$\Delta(\mathfrak{K}) = \mathfrak{K} \otimes q^{-2\mathfrak{C}} + \mathfrak{U}^{-2} \otimes \mathfrak{K}, \qquad \Delta(\mathfrak{U}) = \mathfrak{U} \otimes \mathfrak{U}$$

- \star Z gradation $(s_1, s_2, s_3) = (0, 1, 0)$
- $\mathfrak{P} = g\alpha(1 q^{2\mathfrak{C}}\mathfrak{U}^2)$, $\mathfrak{K} = g\alpha^{-1}(q^{-2\mathfrak{C}} \mathfrak{U}^{-2})$, $\mathfrak{U}^2 = x^+/qx^- \cdot 1$
- In the soliton representation $\mathfrak{U}^2=1$
- $\star \Delta(\mathfrak{P}) = \mathfrak{P} \otimes 1 + 1 \otimes \mathfrak{P}, \qquad \Delta(\mathfrak{K}) = \mathfrak{K} \otimes 1 + 1 \otimes \mathfrak{K},$ needed to interpret \mathfrak{P} and \mathfrak{K} as p_+

Pirsa: 11080035

* U becomes the fermion number

Beisert-Koroteev'08

Involves a new abelian generator

$$\Delta(\mathfrak{E}_{j}) = \mathfrak{E}_{j} \otimes 1 + q^{-\mathfrak{H}_{j}} \mathfrak{U}^{s_{j}} \otimes \mathfrak{E}_{j}, \quad \Delta(\mathfrak{F}_{j}) = \mathfrak{F}_{j} \otimes q^{\mathfrak{H}_{j}} + \mathfrak{U}^{-s_{j}} \otimes \mathfrak{F}_{j}$$

$$\Delta(\mathfrak{C}) = \mathfrak{C} \otimes 1 + 1 \otimes \mathfrak{C}, \qquad \Delta(\mathfrak{P}) = \mathfrak{P} \otimes 1 + q^{2\mathfrak{C}} \mathfrak{U}^{2} \otimes \mathfrak{P}$$

$$\Delta(\mathfrak{K}) = \mathfrak{K} \otimes q^{-2\mathfrak{C}} + \mathfrak{U}^{-2} \otimes \mathfrak{K}, \qquad \Delta(\mathfrak{U}) = \mathfrak{U} \otimes \mathfrak{U}$$

- \star Z gradation $(s_1, s_2, s_3) = (0, 1, 0)$
- $\mathfrak{P} = g\alpha(1 q^{2\mathfrak{C}}\mathfrak{U}^2)$, $\mathfrak{K} = g\alpha^{-1}(q^{-2\mathfrak{C}} \mathfrak{U}^{-2})$, $\mathfrak{U}^2 = x^+/qx^- \cdot 1$
- In the soliton representation $\mathfrak{U}^2=1$
- $\star \Delta(\mathfrak{P}) = \mathfrak{P} \otimes 1 + 1 \otimes \mathfrak{P}, \qquad \Delta(\mathfrak{K}) = \mathfrak{K} \otimes 1 + 1 \otimes \mathfrak{K},$ needed to interpret $\mathfrak P$ and $\mathfrak R$ as p_{\pm}

* U becomes the fermion number

Fundamental (relativistic) S-matrix

Hoare-Tseytlin'11
Hoare-Hollowood-JLM'11

•
$$\widetilde{S}_{11}(\theta_{12}): V_1(\theta_1) \otimes V_1(\theta_2) \longrightarrow V_1(\theta_2) \otimes V_1(\theta_1)$$
 $\theta_{12} = \theta_1 - \theta_2$

$$\widetilde{S}_{11}(\theta) = Y(\theta)Y(i\pi - \theta)\check{R}(e^{\theta})$$

 $\check{R} = g \to \infty$ limit of the fundamental R-matrix for $U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3)$ of Beisert and Koroteev

Fundamental (relativistic) S-matrix

Hoare-Tseytlin'11
Hoare-Hollowood-JLM'11

•
$$\widetilde{S}_{11}(\theta_{12}): V_1(\theta_1) \otimes V_1(\theta_2) \longrightarrow V_1(\theta_2) \otimes V_1(\theta_1)$$
 $\theta_{12} = \theta_1 - \theta_2$

$$\widetilde{S}_{11}(\theta) = Y(\theta)Y(i\pi - \theta)\check{R}(e^{\theta})$$

 $\check{R} = g \to \infty$ limit of the fundamental R-matrix for $U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3)$ of Beisert and Koroteev

• \check{R} -matrix: basis $\{|\phi^a\rangle, |\psi^\alpha\rangle\}, x = e^{\theta}$

$$\begin{split} \tilde{R}(x) \left| \phi^{a} \phi^{a} \right\rangle &= A \left| \phi^{a} \phi^{a} \right\rangle \;, \qquad \tilde{R}(x) \left| \psi^{\alpha} \psi^{\alpha} \right\rangle = D \left| \psi^{\alpha} \psi^{\alpha} \right\rangle \;, \\ \tilde{R}(x) \left| \phi^{1} \phi^{2} \right\rangle &= \frac{q(A-B)}{q^{2}+1} \left| \phi^{2} \phi^{1} \right\rangle + \frac{q^{2}A+B}{q^{2}+1} \left| \phi^{1} \phi^{2} \right\rangle + \frac{C}{1+q^{2}} \left| \psi^{1} \psi^{2} \right\rangle - \frac{qC}{1+q^{2}} \left| \psi^{2} \psi^{1} \right\rangle \;, \\ \tilde{R}(x) \left| \phi^{2} \phi^{1} \right\rangle &= \frac{q(A-B)}{q^{2}+1} \left| \phi^{1} \phi^{2} \right\rangle + \frac{q^{2}B+A}{q^{2}+1} \left| \phi^{2} \phi^{1} \right\rangle - \frac{qC}{1+q^{2}} \left| \psi^{1} \psi^{2} \right\rangle + \frac{q^{2}C}{1+q^{2}} \left| \psi^{2} \psi^{1} \right\rangle \;, \\ \tilde{R}(x) \left| \psi^{1} \psi^{2} \right\rangle &= \frac{q(D-E)}{q^{2}+1} \left| \psi^{2} \psi^{1} \right\rangle + \frac{q^{2}D+E}{q^{2}+1} \left| \psi^{1} \psi^{2} \right\rangle + \frac{F}{1+q^{2}} \left| \phi^{1} \phi^{2} \right\rangle - \frac{qF}{1+q^{2}} \left| \phi^{2} \phi^{1} \right\rangle \;, \\ \tilde{R}(x) \left| \psi^{2} \psi^{1} \right\rangle &= \frac{q(D-E)}{q^{2}+1} \left| \psi^{1} \psi^{2} \right\rangle + \frac{q^{2}E+D}{q^{2}+1} \left| \psi^{2} \psi^{1} \right\rangle - \frac{qF}{1+q^{2}} \left| \phi^{1} \phi^{2} \right\rangle + \frac{q^{2}F}{1+q^{2}} \left| \phi^{2} \phi^{1} \right\rangle \;, \\ \tilde{R}(x) \left| \phi^{a} \psi^{\alpha} \right\rangle &= G \left| \psi^{\alpha} \phi^{a} \right\rangle + H \left| \phi^{a} \psi^{\alpha} \right\rangle \;, \qquad \tilde{R}(x) \left| \psi^{\alpha} \phi^{a} \right\rangle = K \left| \psi^{\alpha} \phi^{a} \right\rangle + L \left| \phi^{a} \psi^{\alpha} \right\rangle \;, \end{split}$$

$$A = \frac{(qx-1)(x+1)}{q^{1/2}x}, \qquad D = \frac{(q-x)(x+1)}{q^{1/2}x},$$

$$B = \frac{q^3 - (q^3 - 2q^2 + 2q - 1)x - x^2}{q^{3/2}x}, \qquad E = \frac{q^3x^2 - (q^3 - 2q^2 + 2q - 1)x - 1}{q^{3/2}x},$$

$$C = F = \frac{i(q-1)(q^2 + 1)(x-1)}{q^{3/2}x^{1/2}}, \qquad G = L = x - x^{-1},$$

$$H = K = \frac{(q-1)(x+1)}{q^{1/2}x^{1/2}}.$$

• \check{R} -matrix: basis $\{|\phi^a\rangle, |\psi^\alpha\rangle\}, x = e^{\theta}$

$$\begin{split} \tilde{R}(x) \left| \phi^{a} \phi^{a} \right\rangle &= A \left| \phi^{a} \phi^{a} \right\rangle \;, \qquad \tilde{R}(x) \left| \psi^{\alpha} \psi^{\alpha} \right\rangle = D \left| \psi^{\alpha} \psi^{\alpha} \right\rangle \;, \\ \tilde{R}(x) \left| \phi^{1} \phi^{2} \right\rangle &= \frac{q(A-B)}{q^{2}+1} \left| \phi^{2} \phi^{1} \right\rangle + \frac{q^{2}A+B}{q^{2}+1} \left| \phi^{1} \phi^{2} \right\rangle + \frac{C}{1+q^{2}} \left| \psi^{1} \psi^{2} \right\rangle - \frac{qC}{1+q^{2}} \left| \psi^{2} \psi^{1} \right\rangle \;, \\ \tilde{R}(x) \left| \phi^{2} \phi^{1} \right\rangle &= \frac{q(A-B)}{q^{2}+1} \left| \phi^{1} \phi^{2} \right\rangle + \frac{q^{2}B+A}{q^{2}+1} \left| \phi^{2} \phi^{1} \right\rangle - \frac{qC}{1+q^{2}} \left| \psi^{1} \psi^{2} \right\rangle + \frac{q^{2}C}{1+q^{2}} \left| \psi^{2} \psi^{1} \right\rangle \;, \\ \tilde{R}(x) \left| \psi^{1} \psi^{2} \right\rangle &= \frac{q(D-E)}{q^{2}+1} \left| \psi^{2} \psi^{1} \right\rangle + \frac{q^{2}D+E}{q^{2}+1} \left| \psi^{1} \psi^{2} \right\rangle + \frac{F}{1+q^{2}} \left| \phi^{1} \phi^{2} \right\rangle - \frac{qF}{1+q^{2}} \left| \phi^{2} \phi^{1} \right\rangle \;, \\ \tilde{R}(x) \left| \psi^{2} \psi^{1} \right\rangle &= \frac{q(D-E)}{q^{2}+1} \left| \psi^{1} \psi^{2} \right\rangle + \frac{q^{2}E+D}{q^{2}+1} \left| \psi^{2} \psi^{1} \right\rangle - \frac{qF}{1+q^{2}} \left| \phi^{1} \phi^{2} \right\rangle + \frac{q^{2}F}{1+q^{2}} \left| \phi^{2} \phi^{1} \right\rangle \;, \\ \tilde{R}(x) \left| \phi^{3} \psi^{\alpha} \right\rangle &= G \left| \psi^{\alpha} \phi^{3} \right\rangle + H \left| \phi^{3} \psi^{\alpha} \right\rangle \;, \qquad \tilde{R}(x) \left| \psi^{\alpha} \phi^{3} \right\rangle = K \left| \psi^{\alpha} \phi^{3} \right\rangle + L \left| \phi^{3} \psi^{\alpha} \right\rangle \;, \end{split}$$

$$A = \frac{(qx-1)(x+1)}{q^{1/2}x}, \qquad D = \frac{(q-x)(x+1)}{q^{1/2}x},$$

$$B = \frac{q^3 - (q^3 - 2q^2 + 2q - 1)x - x^2}{q^{3/2}x}, \qquad E = \frac{q^3x^2 - (q^3 - 2q^2 + 2q - 1)x - 1}{q^{3/2}x},$$

$$C = F = \frac{i(q-1)(q^2 + 1)(x-1)}{q^{3/2}x^{1/2}}, \qquad G = L = x - x^{-1},$$

$$H = K = \frac{(q-1)(x+1)}{q^{1/2}x^{1/2}}.$$

$$\star$$
 $\widetilde{S}_{11}(\theta)$ has $U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^2)$ symmetry

The form of $\widetilde{S}_{11}(\theta)$ ensures that it satisfies crossing symmetry

 $Y(\theta)$ is fixed by the unitarity condition $\widetilde{S}_{11}(\theta)\widetilde{S}_{11}(-\theta) = 1 \otimes 1$

$$\begin{split} Y(\theta)Y(i\pi-\theta) &= \frac{\mathcal{F}(\theta)}{2(q-q^{-1})} : \\ \mathcal{F}(\theta) &= \exp\left[-2\int_0^\infty \frac{dt}{t} \frac{\cosh^2(t(1-\frac{1}{k}))\sinh(t(1-\frac{\theta}{i\pi}))\sinh(\frac{t\theta}{i\pi})}{\sinh t\cosh^2 t}\right] \end{split}$$

$$\star$$
 $\widetilde{S}_{11}(\theta)$ has $U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^2)$ symmetry

The form of $\widetilde{S}_{11}(\theta)$ ensures that it satisfies crossing symmetry

 $Y(\theta)$ is fixed by the unitarity condition $\widetilde{S}_{11}(\theta)\widetilde{S}_{11}(-\theta) = 1 \otimes 1$

$$\begin{split} Y(\theta)Y(i\pi-\theta) &= \frac{\mathcal{F}(\theta)}{2(q-q^{-1})} \;; \\ \mathcal{F}(\theta) &= \exp\left[-2\int_0^\infty \frac{dt}{t} \frac{\cosh^2(t(1-\frac{1}{k}))\sinh(t(1-\frac{\theta}{i\pi}))\sinh(\frac{t\theta}{i\pi})}{\sinh t\cosh^2 t}\right] \end{split}$$

Bound states: the boostrap programme

The most difficult aspect of building a consistent QFT is to explain all the singularities of the S-matrix on the physical strip $0 < \text{Im } \theta < \pi$ in terms of bound states and anomalous thresholds

Pirsa: 11080035 Page 108/166

Bound states: the boostrap programme

- The most difficult aspect of building a consistent QFT is to explain all the singularities of the S-matrix on the physical strip $0 < \text{Im } \theta < \pi$ in terms of bound states and anomalous thresholds
- Bound states give rise to simple poles, and their positions have to mesh with the representation theory of the quantum group

If a bound state corresponding to a multiplet V_c is produced in the collision of V_a and V_b in the direct channel, then

- ② The representation $V_a(\theta_1) \otimes V_b(\theta_2)$ becomes reducible: $V_a \otimes V_b = V_c \oplus V_c^{\perp}$

Bound states: the boostrap programme

- The most difficult aspect of building a consistent QFT is to explain all the singularities of the S-matrix on the physical strip $0 < \text{Im } \theta < \pi$ in terms of bound states and anomalous thresholds
- Bound states give rise to simple poles, and their positions have to mesh with the representation theory of the quantum group

If a bound state corresponding to a multiplet V_c is produced in the collision of V_a and V_b in the direct channel, then

- $S_{ab}(\theta)$ has a simple pole at $\theta = iu_{ab}^c$
- ② The representation $V_a(\theta_1) \otimes V_b(\theta_2)$ becomes reducible: $V_a \otimes V_b = V_c \oplus V_c^{\perp}$

Res $S_{ab}(iu_{ab}^c)$ is a weighted sum of "projectors"

$$V_c = \bigoplus_j V_c^{(j)}, \quad V_c^{(j)} = \text{irreducible representations of } U_q(\mathfrak{sl}(2)^{\times 2})$$

$$S_{ab}(\theta) \sim \frac{i}{\theta - iu_{ab}^c} \sum_{i} \rho_j \, \mathbb{P}_{ab}^{c,j}, \qquad \mathbb{P}_{ab}^{c,j} = \mathbb{P}_{c,j}^{ba} \, \mathbb{P}_{ab}^{c,j}$$

Res $S_{ab}(iu_{ab}^c)$ is a weighted sum of "projectors"

$$V_c = \bigoplus_j V_c^{(j)}, \quad V_c^{(j)} = \text{irreducible representations of } U_q(\mathfrak{sl}(2)^{\times 2})$$

$$S_{ab}(\theta) \sim \frac{i}{\theta - iu_{ab}^c} \sum_{j} \rho_j \, \mathbb{P}_{ab}^{c,j}, \qquad \mathbb{P}_{ab}^{c,j} = \mathbb{P}_{c,j}^{ba} \, \mathbb{P}_{ab}^{c,j}$$

Res $S_{ab}(iu_{ab}^c)$ is a weighted sum of "projectors"

$$V_c = \bigoplus_j V_c^{(j)}, \quad V_c^{(j)} = \text{irreducible representations of } U_q(\mathfrak{sl}(2)^{\times 2})$$

$$S_{ab}(\theta) \sim \frac{i}{\theta - iu_{ab}^c} \sum_{j} \rho_j \, \mathbb{P}_{ab}^{c,j}, \qquad \mathbb{P}_{ab}^{c,j} = \mathcal{P}_{c,j}^{ba} \, \mathcal{P}_{ab}^{c,j}$$

★ The scattering amplitudes of the bound state c with state d can be constructed in term of those of d with a and b.

$$S_{dc}(\theta) = \left(\sum_{j} \sqrt{|\rho_{j}|} \, \mathcal{P}_{ab}^{c,j} \otimes 1\right) \left(1 \otimes S_{db}(\theta + i\bar{u}_{b\bar{c}}^{\bar{a}})\right) \\ \times \left(S_{da}(\theta - i\bar{u}_{a\bar{c}}^{\bar{b}}) \otimes 1\right) \left(1 \otimes \sum_{l} \frac{1}{\sqrt{|\rho_{l}|}} \, \mathcal{P}_{c,l}^{ab}\right)$$

Res $S_{ab}(iu_{ab}^c)$ is a weighted sum of "projectors"

$$V_c = \bigoplus_j V_c^{(j)}, \quad V_c^{(j)} = \text{irreducible representations of } U_q(\mathfrak{sl}(2)^{\times 2})$$

$$S_{ab}(\theta) \sim \frac{i}{\theta - iu_{ab}^c} \sum_{j} \rho_j \, \mathbb{P}_{ab}^{c,j}, \qquad \mathbb{P}_{ab}^{c,j} = \mathcal{P}_{c,j}^{ba} \, \mathcal{P}_{ab}^{c,j}$$

★ The scattering amplitudes of the bound state c with state d can be constructed in term of those of d with a and b.

$$S_{dc}(\theta) = \left(\sum_{j} \sqrt{|\rho_{j}|} \, \mathcal{P}_{ab}^{c,j} \otimes 1\right) \left(1 \otimes S_{db}(\theta + i\bar{u}_{b\bar{c}}^{\bar{a}})\right) \\ \times \left(S_{da}(\theta - i\bar{u}_{a\bar{c}}^{\bar{b}}) \otimes 1\right) \left(1 \otimes \sum_{l} \frac{1}{\sqrt{|\rho_{l}|}} \, \mathcal{P}_{c,l}^{ab}\right)$$

Representation theory

Zhang-Gould'05 Beisert'07 Beisert-Koroteev'08

• For q not a root of unity, the representations of $U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3)$ are simply deformations of the representations of $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$

Representation theory

Zhang-Gould'05 Beisert'07 Beisert-Koroteev'08

• For q not a root of unity, the representations of $U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3)$ are simply deformations of the representations of $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$

Representation theory

Zhang-Gould'05 Beisert'07 Beisert-Koroteev'08

- For q not a root of unity, the representations of $U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3)$ are simply deformations of the representations of $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$
- Long (typical) representations $\{m, n\} \mapsto \dim = 16(m+1)(n+1)$ Irreducible for generic values of C, P, K
 - Become reducible but indecomposable for specific values of $[C]_q^2 PK$ (shortening or BPS conditions)

$$V_{\{m,n\}} = V_{\text{sub-rep}} \oplus V^{\perp}, \qquad V_{factor} = V_{\{m,n\}}/V_{\text{sub-rep}}$$

 $V_{\text{sub-rep}}$, $V_{factor} \equiv \text{short (atypical)}$ representations

Representation theory

Zhang-Gould'05 Beisert'07 Beisert-Koroteev'08

- For q not a root of unity, the representations of $U_q(\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3)$ are simply deformations of the representations of $\mathfrak{psl}(2|2) \ltimes \mathbb{R}^3$
- Long (typical) representations $\{m, n\} \mapsto \dim = 16(m+1)(n+1)$ Irreducible for generic values of C, P, K
 - Become reducible but indecomposable for specific values of $[C]_q^2 PK$ (shortening or BPS conditions)

$$V_{\{m,n\}} = V_{\mathsf{sub-rep}} \oplus V^{\perp}, \qquad V_{\mathsf{factor}} = V_{\{m,n\}}/V_{\mathsf{sub-rep}}$$

 $V_{\text{sub-rep}}$, $V_{factor} \equiv \text{short (atypical)}$ representations

Short representation $(m, n) \mapsto \dim = 4(m+1)(n+1) + 4mn$ Exists for $[C]_q^2 - PK = [(m+n+1)/2]_q^2 \longrightarrow \text{Shortening condition}$

★ Fundamental representation = $\langle 0, 0 \rangle$ Semiclassical solitons live in $\langle m, 0 \rangle$

Short representation $(m, n) \mapsto \dim = 4(m+1)(n+1) + 4mn$ Exists for $[C]_q^2 - PK = [(m+n+1)/2]_q^2 \longrightarrow \text{Shortening condition}$

★ Fundamental representation = $\langle 0, 0 \rangle$ Semiclassical solitons live in $\langle m, 0 \rangle$

Short representation
$$(m, n) \mapsto \dim = 4(m+1)(n+1) + 4mn$$

Exists for $[C]_q^2 - PK = [(m+n+1)/2]_q^2 \longrightarrow \text{Shortening condition}$

- ★ Fundamental representation = $\langle 0, 0 \rangle$ Semiclassical solitons live in $\langle m, 0 \rangle$
 - Tensor product $\langle m, 0 \rangle \otimes \langle n, 0 \rangle = \sum_{k=0}^{\min(m,n)} \{m+n-2k, 0\}$
 - Multiplet splittings

$$\{m,0\} \longrightarrow \langle m+1,0\rangle \oplus \langle m,1\rangle \text{ for } [C]_q^2 - PK = [(m+2)/2]_q^2$$

 $\longrightarrow \langle m-1,0\rangle \oplus \langle m,0\rangle_3 \text{ for } [C]_q^2 - PK = [m/2]_q^2$

Short representation
$$(m, n) \mapsto \dim = 4(m+1)(n+1) + 4mn$$

Exists for $[C]_q^2 - PK = [(m+n+1)/2]_q^2 \longrightarrow \text{Shortening condition}$

- ★ Fundamental representation = $\langle 0, 0 \rangle$ Semiclassical solitons live in $\langle m, 0 \rangle$
 - Tensor product $\langle m, 0 \rangle \otimes \langle n, 0 \rangle = \sum_{k=0}^{\min(m,n)} \{m+n-2k, 0\}$
 - Multiplet splittings

$$\{m,0\} \longrightarrow \langle m+1,0\rangle \oplus \langle m,1\rangle \text{ for } [C]_q^2 - PK = [(m+2)/2]_q^2$$

 $\longrightarrow \langle m-1,0\rangle \oplus \langle m,0\rangle_3 \text{ for } [C]_q^2 - PK = [m/2]_q^2$

The shortening conditions indicate the location of the poles corresponding to the bound states

•
$$\widetilde{S}_{11}(\theta_{12}): \widetilde{V_{\langle 0,0\rangle}(\theta_1)} \otimes V_{\langle 0,0\rangle}(\theta_2) \longrightarrow V_{\langle 0,0\rangle}(\theta_2) \otimes V_{\langle 0,0\rangle}(\theta_1)$$

$$C_1 = C_2 = 0, \quad P = P_1 + P_2, \quad K = K_1 + K_2$$

• Shortening condition for $\{0,0\} \rightarrow \langle 1,0 \rangle \oplus \langle 0,1 \rangle$

$$-PK = [1]_q^2 \Rightarrow \theta_{12} = \pm \frac{i\pi}{k}$$

The shortening conditions indicate the location of the poles corresponding to the bound states

•
$$\widetilde{S}_{11}(\theta_{12}): V_{\langle 0,0\rangle}(\theta_1) \otimes V_{\langle 0,0\rangle}(\theta_2) \longrightarrow V_{\langle 0,0\rangle}(\theta_2) \otimes V_{\langle 0,0\rangle}(\theta_1)$$

$$C_1 = C_2 = 0, \quad P = P_1 + P_2, \quad K = K_1 + K_2$$

• Shortening condition for $\{0,0\} \rightarrow \langle 1,0 \rangle \oplus \langle 0,1 \rangle$

$$-PK = [1]_q^2 \Rightarrow \theta_{12} = \pm \frac{i\pi}{k}$$

The shortening conditions indicate the location of the poles corresponding to the bound states

•
$$\widetilde{S}_{11}(\theta_{12}): \widetilde{V_{\langle 0,0\rangle}(\theta_1)} \otimes V_{\langle 0,0\rangle}(\theta_2) \longrightarrow V_{\langle 0,0\rangle}(\theta_2) \otimes V_{\langle 0,0\rangle}(\theta_1)$$

• $C_1 = C_2 = 0$, $P = P_1 + P_2$, $K = K_1 + K_2$

• Shortening condition for $\{0,0\} \rightarrow \langle 1,0 \rangle \oplus \langle 0,1 \rangle$

$$-PK = [1]_q^2 \Rightarrow \theta_{12} = \pm \frac{i\pi}{k}$$

- Bound state at $\theta = i\pi/k$ in the (factor) representation $\langle 1, 0 \rangle$
- * Consistent with the quantum group representation theory

$$\widetilde{S}_{11}(i\pi/k):V_{\langle 0,1\rangle}\longrightarrow 0$$

The shortening conditions indicate the location of the poles corresponding to the bound states

•
$$\widetilde{S}_{11}(\theta_{12}): \widetilde{V_{\langle 0,0\rangle}(\theta_1)} \otimes V_{\langle 0,0\rangle}(\theta_2) \longrightarrow V_{\langle 0,0\rangle}(\theta_2) \otimes V_{\langle 0,0\rangle}(\theta_1)$$

$$C_1 = C_2 = 0, \quad P = P_1 + P_2, \quad K = K_1 + K_2$$

• Shortening condition for $\{0,0\} \rightarrow \langle 1,0 \rangle \oplus \langle 0,1 \rangle$

$$-PK = [1]_q^2 \Rightarrow \theta_{12} = \pm \frac{i\pi}{k}$$

- Bound state at $\theta = i\pi/k$ in the (factor) representation $\langle 1, 0 \rangle$
- ★ Consistent with the quantum group representation theory

$$\widetilde{S}_{11}(i\pi/k):V_{\langle 0,1\rangle}\longrightarrow 0$$

•
$$V_{\{0,0\}} = \underbrace{V_{(2,0)} \oplus V_{(1,1)}^{(+)} \oplus V_{(0,0)}^{(+)} \oplus V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle}$$

$$\widetilde{S}_{11}(\frac{i\pi}{k}) \propto \frac{q+1}{2q^{1/2}} \mathbb{P}_{(2,0)} + \mathbb{P}_{(1,1)}^{(+)} + \frac{q^4 - q^3 + 4q^2 - q + 1}{2q^{3/2}(q+1)} \mathbb{P}_{(0,0)}^{(+)}$$

•
$$V_{\{0,0\}} = \underbrace{V_{(2,0)} \oplus V_{(1,1)}^{(+)} \oplus V_{(0,0)}^{(+)} \oplus V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1\rangle} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1\rangle}$$

$$\widetilde{S}_{11}(\frac{i\pi}{k}) \propto \frac{q+1}{2q^{1/2}} \mathbb{P}_{(2,0)} + \mathbb{P}_{(1,1)}^{(+)} + \frac{q^4-q^3+4q^2-q+1}{2q^{3/2}(q+1)} \mathbb{P}_{(0,0)}^{(+)}$$

•
$$V_{\{0,0\}} = \underbrace{V_{(2,0)} \oplus V_{(1,1)}^{(+)} \oplus V_{(0,0)}^{(+)} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle}$$

$$\widetilde{S}_{11}(\frac{i\pi}{k}) \propto \frac{q+1}{2q^{1/2}} \mathbb{P}_{(2,0)} + \mathbb{P}_{(1,1)}^{(+)} + \frac{q^4-q^3+4q^2-q+1}{2q^{3/2}(q+1)} \mathbb{P}_{(0,0)}^{(+)}$$



$$\widetilde{S}_{12}(\theta) = \left(1 \otimes \widetilde{S}_{11}(\theta + \frac{i\pi}{2k})\right) \left(\widetilde{S}_{11}(\theta - \frac{i\pi}{2k}) \otimes 1\right)\Big|_{V_{(0,0)} \otimes V_{(1,0)}}$$

•
$$V_{\{0,0\}} = \underbrace{V_{(2,0)} \oplus V_{(1,1)}^{(+)} \oplus V_{(0,0)}^{(+)} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle}$$

$$\widetilde{S}_{11}(\frac{i\pi}{k}) \propto \frac{q+1}{2q^{1/2}} \mathbb{P}_{(2,0)} + \mathbb{P}_{(1,1)}^{(+)} + \frac{q^4-q^3+4q^2-q+1}{2q^{3/2}(q+1)} \mathbb{P}_{(0,0)}^{(+)}$$

*

$$\widetilde{S}_{12}(\theta) = \left(1 \otimes \widetilde{S}_{11}(\theta + \frac{i\pi}{2k})\right) \left(\widetilde{S}_{11}(\theta - \frac{i\pi}{2k}) \otimes 1\right)\Big|_{V_{(0,0)} \otimes V_{(1,0)}}$$

•
$$V_{\{0,0\}} = \underbrace{V_{(2,0)} \oplus V_{(1,1)}^{(+)} \oplus V_{(0,0)}^{(+)} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle}$$

$$\widetilde{S}_{11}(\frac{i\pi}{k}) \propto \frac{q+1}{2q^{1/2}} \mathbb{P}_{(2,0)} + \mathbb{P}_{(1,1)}^{(+)} + \frac{q^4-q^3+4q^2-q+1}{2q^{3/2}(q+1)} \mathbb{P}_{(0,0)}^{(+)}$$

$$\star \qquad \widetilde{S}_{12}(\theta) = \left(1 \otimes \widetilde{S}_{11}(\theta + \frac{i\pi}{2k})\right) \left(\widetilde{S}_{11}(\theta - \frac{i\pi}{2k}) \otimes 1\right) \Big|_{V_{(0,0)} \otimes V_{(1,0)}}$$

 $S_{12}(\theta)$ has four simple poles that can be explained in terms of the fusions $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle}\to V_{\langle 3,0\rangle}$ and $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle}\to V_{\langle 1,0\rangle}$ in the direct- and cross-channels

.

•
$$V_{\{0,0\}} = \underbrace{V_{(2,0)} \oplus V_{(1,1)}^{(+)} \oplus V_{(0,0)}^{(+)} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle}$$

$$\widetilde{S}_{11}(\frac{i\pi}{k}) \propto \frac{q+1}{2q^{1/2}} \mathbb{P}_{(2,0)} + \mathbb{P}_{(1,1)}^{(+)} + \frac{q^4-q^3+4q^2-q+1}{2q^{3/2}(q+1)} \mathbb{P}_{(0,0)}^{(+)}$$

$$\star \qquad \widetilde{S}_{12}(\theta) = \left(1 \otimes \widetilde{S}_{11}(\theta + \frac{i\pi}{2k})\right) \left(\widetilde{S}_{11}(\theta - \frac{i\pi}{2k}) \otimes 1\right) \Big|_{V_{(0,0)} \otimes V_{(1,0)}}$$

 $\widetilde{S}_{12}(\theta)$ has four simple poles that can be explained in terms of the fusions $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle}\to V_{\langle 3,0\rangle}$ and $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle}\to V_{\langle 1,0\rangle}$ in the direct- and cross-channels

Quantum spectrum of particles associated to the representations

$$V_a(\theta) = \langle a - 1, 0 \rangle$$
 with mass $m_a = \mu \sin(\pi a/2k)$

→ The semiclassical spectrum is exact!

Bound states of a and b correspond to simple poles at

$$\theta = i\pi(a+b)/2k$$
 and $i\pi - i\pi|a-b|/2k$

•
$$V_{\{0,0\}} = \underbrace{V_{(2,0)} \oplus V_{(1,1)}^{(+)} \oplus V_{(0,0)}^{(+)} \oplus V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle}$$

$$\widetilde{S}_{11}(\frac{i\pi}{k}) \propto \frac{q+1}{2q^{1/2}} \mathbb{P}_{(2,0)} + \mathbb{P}_{(1,1)}^{(+)} + \frac{q^4-q^3+4q^2-q+1}{2q^{3/2}(q+1)} \mathbb{P}_{(0,0)}^{(+)}$$

$$\star \qquad \widetilde{S}_{12}(\theta) = \left(1 \otimes \widetilde{S}_{11}(\theta + \frac{i\pi}{2k})\right) \left(\widetilde{S}_{11}(\theta - \frac{i\pi}{2k}) \otimes 1\right) \Big|_{V_{(0,0)} \otimes V_{(1,0)}}$$

 $\widetilde{S}_{12}(\theta)$ has four simple poles that can be explained in terms of the fusions $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle} \to V_{\langle 3,0\rangle}$ and $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle} \to V_{\langle 1,0\rangle}$ in the direct- and cross-channels

.

Quantum spectrum of particles associated to the representations

$$V_a(\theta) = \langle a - 1, 0 \rangle$$
 with mass $m_a = \mu \sin(\pi a/2k)$

→ The semiclassical spectrum is exact!

Bound states of a and b correspond to simple poles at

$$\theta = i\pi(a+b)/2k$$
 and $i\pi - i\pi|a-b|/2k$

•
$$V_{\{0,0\}} = \underbrace{V_{(2,0)} \oplus V_{(1,1)}^{(+)} \oplus V_{(0,0)}^{(+)} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle} \oplus \underbrace{V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle}$$

$$\widetilde{S}_{11}(\frac{i\pi}{k}) \propto \frac{q+1}{2q^{1/2}} \mathbb{P}_{(2,0)} + \mathbb{P}_{(1,1)}^{(+)} + \frac{q^4-q^3+4q^2-q+1}{2q^{3/2}(q+1)} \mathbb{P}_{(0,0)}^{(+)}$$

$$\star \qquad \widetilde{S}_{12}(\theta) = \left(1 \otimes \widetilde{S}_{11}(\theta + \frac{i\pi}{2k})\right) \left(\widetilde{S}_{11}(\theta - \frac{i\pi}{2k}) \otimes 1\right) \Big|_{V_{(0,0)} \otimes V_{(1,0)}}$$

 $\widetilde{S}_{12}(\theta)$ has four simple poles that can be explained in terms of the fusions $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle}\to V_{\langle 3,0\rangle}$ and $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle}\to V_{\langle 1,0\rangle}$ in the direct- and cross-channels

•
$$V_{\{0,0\}} = \underbrace{V_{(2,0)} \oplus V_{(1,1)}^{(+)} \oplus V_{(0,0)}^{(+)} \oplus V_{(0,0)}^{(-)} \oplus V_{(1,1)}^{(-)} \oplus V_{(0,2)}^{(-)}}_{\langle 0,1 \rangle}$$

$$\widetilde{S}_{11}(\frac{i\pi}{k}) \propto \frac{q+1}{2q^{1/2}} \mathbb{P}_{(2,0)} + \mathbb{P}_{(1,1)}^{(+)} + \frac{q^4-q^3+4q^2-q+1}{2q^{3/2}(q+1)} \mathbb{P}_{(0,0)}^{(+)}$$

$$\star \qquad \left| \widetilde{S}_{12}(\theta) = \left(1 \otimes \widetilde{S}_{11}(\theta + \frac{i\pi}{2k}) \right) \left(\widetilde{S}_{11}(\theta - \frac{i\pi}{2k}) \otimes 1 \right) \right|_{V_{\langle 0,0 \rangle} \otimes V_{\langle 1,0 \rangle}}$$

 $\widetilde{S}_{12}(\theta)$ has four simple poles that can be explained in terms of the fusions $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle}\to V_{\langle 3,0\rangle}$ and $V_{\langle 1,0\rangle}\otimes V_{\langle 2,0\rangle}\to V_{\langle 1,0\rangle}$ in the direct- and cross-channels

Quantum spectrum of particles associated to the representations

$$V_a(\theta) = \langle a - 1, 0 \rangle$$
 with mass $m_a = \mu \sin(\pi a/2k)$

→ The semiclassical spectrum is exact!

Bound states of a and b correspond to simple poles at

$$\theta = i\pi(a+b)/2k$$
 and $i\pi - i\pi|a-b|/2k$



Quantum spectrum of particles associated to the representations

$$V_a(\theta) = \langle a-1, 0 \rangle$$
 with mass $m_a = \mu \sin(\pi a/2k)$

→ The semiclassical spectrum is exact!

Bound states of a and b correspond to simple poles at

$$\theta = i\pi(a+b)/2k$$
 and $i\pi - i\pi|a-b|/2k$

Quantum spectrum of particles associated to the representations

$$V_a(\theta) = \langle a - 1, 0 \rangle$$
 with mass $m_a = \mu \sin(\pi a/2k)$

→ The semiclassical spectrum is exact!

Bound states of a and b correspond to simple poles at

$$\theta = i\pi(a+b)/2k$$
 and $i\pi - i\pi |a-b|/2k$

* This meshes with the quantum group representation theory

$$V_a(\theta_1) \otimes V_b(\theta_2) = \{a+b-2,0\} \oplus \{a+b-4,0\} \oplus \cdots \oplus \{|a-b|,0\}$$

- At $\theta_{12}=i\pi(a+b)/2k$ the representation $\{a+b-2,0\}$ becomes reducible with a factor representation V_{a+b} , and $\widetilde{S}_{ab}(\theta_{12})$ is only non-vanishing on V_{a+b}
- At $\theta_{12}=i\pi-i\pi|a-b|/2k$ the representation $\{|a-b|,0\}$ becomes reducible with a factor representation $V_{|a-b|}$, and $\widetilde{S}_{ab}(\theta_{12})$ is only non-vanishing on $V_{|a-b|}$

Quantum spectrum of particles associated to the representations

$$V_a(\theta) = \langle a - 1, 0 \rangle$$
 with mass $m_a = \mu \sin(\pi a/2k)$

→ The semiclassical spectrum is exact!

Bound states of a and b correspond to simple poles at

$$\theta = i\pi(a+b)/2k$$
 and $i\pi - i\pi |a-b|/2k$

* This meshes with the quantum group representation theory

$$V_a(\theta_1) \otimes V_b(\theta_2) = \{a+b-2,0\} \oplus \{a+b-4,0\} \oplus \cdots \oplus \{|a-b|,0\}$$

- At $\theta_{12}=i\pi(a+b)/2k$ the representation $\{a+b-2,0\}$ becomes reducible with a factor representation V_{a+b} , and $\widetilde{S}_{ab}(\theta_{12})$ is only non-vanishing on V_{a+b}
- At $\theta_{12}=i\pi-i\pi|a-b|/2k$ the representation $\{|a-b|,0\}$ becomes reducible with a factor representation $V_{|a-b|}$, and $\widetilde{S}_{ab}(\theta_{12})$ is only non-vanishing on $V_{|a-b|}$

The relativistic SSSSG S-matrix

- Particles in $\langle a-1,0\rangle\otimes_{\operatorname{gr}}\langle a-1,0\rangle$ representations of the quantum symmetry group U_q (psl(2|2) $^2\ltimes\mathbb{R}^2$)
- $S_{ab}(\theta) = X_{ab}(\theta) \widetilde{S}_{ab}(\theta) \otimes_{gr} \widetilde{S}_{ab}(\theta)$
- $X_{ab}(\theta)$ \longrightarrow supplies the simple poles on the physical strip

$$(x) = \frac{\sinh(\frac{\theta}{2} + \frac{i\pi x}{4k})}{\sinh(\frac{\theta}{2} - \frac{i\pi x}{4k})}, \quad \{x\} = (x - 1)(x + 1), \quad [x] = \{x\}\{2k - x\}$$

$$X_{ab}(\theta) = [a+b-1][a+b-3]\cdots[|a-b|+1]$$

Page 142/166

The relativistic SSSSG S-matrix

- Particles in $\langle a-1,0\rangle\otimes_{\operatorname{gr}}\langle a-1,0\rangle$ representations of the quantum symmetry group U_q (psl(2|2) $^2\ltimes\mathbb{R}^2$)
- $S_{ab}(\theta) = X_{ab}(\theta) \widetilde{S}_{ab}(\theta) \otimes_{gr} \widetilde{S}_{ab}(\theta)$
- $X_{ab}(\theta)$ supplies the simple poles on the physical strip

$$(x) = \frac{\sinh(\frac{\theta}{2} + \frac{i\pi x}{4k})}{\sinh(\frac{\theta}{2} - \frac{i\pi x}{4k})}, \quad \{x\} = (x - 1)(x + 1), \quad [x] = \{x\}\{2k - x\}$$

$$X_{ab}(\theta) = [a+b-1][a+b-3]\cdots[|a-b|+1]$$

Open questions

Open questions

- Complete the closure of the bootstrap
 - Interpretation of all the singularities on the physical strip: bound states, anomalous thresholds, etc.

- Complete the closure of the bootstrap
 - Interpretation of all the singularities on the physical strip: bound states, anomalous thresholds, etc.

- Complete the closure of the bootstrap
 - Interpretation of all the singularities on the physical strip: bound states, anomalous thresholds, etc.
 - **Truncation** of the spectrum: $m_a = \mu \sin(\pi a/2k)$, a = 1, ..., k
 - Requires a better understanding of the quantum group representations for $q=e^{i\pi/k}$ a root of unity

- Complete the closure of the bootstrap
 - Interpretation of all the singularities on the physical strip: bound states, anomalous thresholds, etc.
 - \star Truncation of the spectrum: $m_a = \mu \sin(\pi a/2k)$, a = 1, ..., k
 - Requires a better understanding of the quantum group representations for $q=e^{i\pi/k}$ a root of unity

- Complete the closure of the bootstrap
 - Interpretation of all the singularities on the physical strip: bound states, anomalous thresholds, etc.
 - \star Truncation of the spectrum: $m_a = \mu \sin(\pi a/2k)$, a = 1, ..., k
 - Requires a better understanding of the quantum group representations for $q=e^{i\pi/k}$ a root of unity
- Algebraic structure underlying the relativistic S-matrix

→ Beisert-Galleas-Matsumoto'11

- Complete the closure of the bootstrap
 - Interpretation of all the singularities on the physical strip: bound states, anomalous thresholds, etc.
 - \star Truncation of the spectrum: $m_a = \mu \sin(\pi a/2k)$, a = 1, ..., k
 - Requires a better understanding of the quantum group representations for $q=e^{i\pi/k}$ a root of unity
- Algebraic structure underlying the relativistic S-matrix

→ Beisert-Galleas-Matsumoto'11

- Complete the closure of the bootstrap
 - Interpretation of all the singularities on the physical strip: bound states, anomalous thresholds, etc.
 - \star Truncation of the spectrum: $m_a = \mu \sin(\pi a/2k)$, a = 1, ..., k
 - Requires a better understanding of the quantum group representations for $q=e^{i\pi/k}$ a root of unity
- Algebraic structure underlying the relativistic S-matrix

Connection with semiclassical calculations

---- Hoare-Iwashita-Roiban-Tseytlin'09-11

- Complete the closure of the bootstrap
 - Interpretation of all the singularities on the physical strip: bound states, anomalous thresholds, etc.
 - \star Truncation of the spectrum: $m_a = \mu \sin(\pi a/2k)$, a = 1, ..., k
 - Requires a better understanding of the quantum group representations for $q=e^{i\pi/k}$ a root of unity
- Algebraic structure underlying the relativistic S-matrix

Connection with semiclassical calculations

How can a relativistic S-matrix be equivalent to a non-relativistic one?

Pirsa: 11080035 Page 152/166

How can a relativistic S-matrix be equivalent to a non-relativistic one?

Non-relativistic interpolating S-matrix

$$S[g,q] \xrightarrow{q \to 1}$$
 (non-relativistic) string/magnon S-matrix $\xrightarrow{g \to \infty}$ (relativistic) SSSSG S-matrix

How can a relativistic S-matrix be equivalent to a non-relativistic one?

Non-relativistic interpolating S-matrix

$$S[g,q] \xrightarrow{q \to 1}$$
 (non-relativistic) string/magnon S-matrix $\xrightarrow{g \to \infty}$ (relativistic) SSSSG S-matrix

How can a relativistic S-matrix be equivalent to a non-relativistic one?

Non-relativistic interpolating S-matrix

$$S[g,q] \xrightarrow{q \to 1}$$
 (non-relativistic) string/magnon S-matrix $\xrightarrow{g \to \infty}$ (relativistic) SSSSG S-matrix

Quantum version of the intepolating classical Poisson brackets

How can a relativistic S-matrix be equivalent to a non-relativistic one?

Non-relativistic interpolating S-matrix

$$S[g,q] \xrightarrow{q \to 1}$$
 (non-relativistic) string/magnon S-matrix $\xrightarrow{g \to \infty}$ (relativistic) SSSSG S-matrix

Quantum version of the intepolating classical Poisson brackets

How can a relativistic S-matrix be equivalent to a non-relativistic one?

Non-relativistic interpolating S-matrix

$$S[g,q] \xrightarrow{q \to 1}$$
 (non-relativistic) string/magnon S-matrix $\xrightarrow{g \to \infty}$ (relativistic) SSSSG S-matrix

- Quantum version of the intepolating classical Poisson brackets
- Requires the construction of the interpolating dressing function

 [Hoare-Hollowood-JLM in progress]

How can a relativistic S-matrix be equivalent to a non-relativistic one?

Non-relativistic interpolating S-matrix

$$S[g,q] \xrightarrow{q \to 1}$$
 (non-relativistic) string/magnon S-matrix $\xrightarrow{g \to \infty}$ (relativistic) SSSSG S-matrix

- Quantum version of the intepolating classical Poisson brackets
- ★ Requires the construction of the interpolating dressing function

 [Hoare-Hollowood-JLM in progress]

Bound states: bootstrap

Quantum spectrum of particles associated to the representations

$$V_a(\theta) = \langle a-1, 0 \rangle$$
 with mass $m_a = \mu \sin(\pi a/2k)$

The semiclassical spectrum is exact!

Bound states of a and b correspond to simple poles at

$$\theta = i\pi(a+b)/2k$$
 and $i\pi - i\pi|a-b|/2k$

* This meshes with the quantum group representation theory

$$V_a(\theta_1) \otimes V_b(\theta_2) = \{a+b-2,0\} \oplus \{a+b-4,0\} \oplus \cdots \oplus \{|a-b|,0\}$$

- At $\theta_{12} = i\pi(a+b)/2k$ the representation $\{a+b-2,0\}$ becomes reducible with a factor representation V_{a+b} , and $\widetilde{S}_{ab}(\theta_{12})$ is only non-vanishing on V_{a+b}
- At $\theta_{12}=i\pi-i\pi|a-b|/2k$ the representation $\{|a-b|,0\}$ becomes reducible with a factor representation $V_{|a-b|}$, and $\widetilde{S}_{ab}(\theta_{12})$ is only non-vanishing on $V_{|a-b|}$

Bound states: bootstrap

Res $S_{ab}(iu_{ab}^c)$ is a weighted sum of "projectors"

$$V_c = \bigoplus_j V_c^{(j)}, \quad V_c^{(j)} = \text{irreducible representations of } U_q(\mathfrak{sl}(2)^{\times 2})$$

$$S_{ab}(\theta) \sim \frac{i}{\theta - iu_{ab}^c} \sum_{i} \rho_j \, \mathbb{P}_{ab}^{c,j}, \qquad \mathbb{P}_{ab}^{c,j} = \mathcal{P}_{c,j}^{ba} \, \mathcal{P}_{ab}^{c,j}$$

The magnon and the soliton representations

Magnon representation:

$$\boxed{q \to 1} \quad (\text{or } k \to \infty) \Rightarrow \qquad \frac{x^+}{x^-} = e^{ip}$$

 $p \equiv \text{world-sheet momentum}$ $g \equiv \text{string tension}$

$$[C]_q^2 - PK = [1/2]_q^2 \longrightarrow \epsilon(p) = \sqrt{1 + 4g^2 \sin^2(p/2)}$$

Hoare-Hollowood-JLM'11

Soliton representation:

$$g \to \infty$$
 \Rightarrow $\frac{x^+}{x^-} = q$, $C = 0$

Pirsa: 11080035
$$\begin{bmatrix} C \end{bmatrix}_q^2 - PK = \begin{bmatrix} 1/2 \end{bmatrix}_q^2$$
 becomes a relativistic mass-shell condition

The $AdS_5 \times S^5$ SSSSG theory

 \bigstar $m_a=\mu\sin\left(\frac{\pi a}{2k}\right)$ can be understood as the BPS condition for the short representation of $U_q(\mathfrak{psu}(2|2)\ltimes\mathbb{R}^2)$ of dimension $4a\times 4a$ with $q=e^{i\pi/k}$

The semiclassical spectrum consists of a tower of massive states in short symmetric representations, just like the magnon bound states of the superstring world sheet theory......

but in the SSSG theory the tower is truncated by k, the level of the WZW term....

Solitons

Hellowood-JLM'11

Continuous spectrum of relativistic non-abelian Q-ball kinks with bosonic and fermionic degrees of freedom

labelled by
$$\varphi \in (0, \pi/2) \longrightarrow m_{\varphi} = \frac{2k}{\pi} \sin \varphi$$

Non-trivial moduli space
$$\mathfrak{M} = \frac{SU(2|2)}{U(2|1)} \times \frac{SU(2|2)}{U(2|1)}$$

Hidden symmetries

Infinite (classical) symmetry algebra

$$\hat{\mathfrak{f}}^{\perp} = \operatorname{Ker}(\operatorname{ad}\Lambda) \cap \mathcal{L}(\mathfrak{psu}(2,2|4),\sigma)$$

Hellowood-JLM'11 Hoare-Tseytlin'11

The elements of grade ± 1 (or Lorentz spin $\pm 1/2$) generate SUSY transformations whose closure is the (exotic) $\mathcal{N} = (8|8)$ superalgebra

$$\mathfrak{s} = \big(\mathfrak{psu}(2|2) \oplus \mathfrak{psu}(2|2)\big) \ltimes \big(\mathbb{R} \oplus \mathbb{R}\big) = \mathfrak{s}_{-2} \oplus \mathfrak{s}_{-1} \oplus \mathfrak{s}_0 \oplus \mathfrak{s}_{+1} \oplus \mathfrak{s}_{+2}$$

- \mapsto \mathfrak{s}_{+2} : central elements corresponding to the components of p_{μ}
- \rightarrow 5+1: generators of SUSY transformations
- → 50: generators of global gauge transformations \equiv non-abelian R-symmetry group $SU(2)^{\times 4}$
- \mathfrak{s} is a finite subalgebra of $\mathcal{L}(\mathfrak{p}(\mathfrak{su}(2|2) \oplus \mathfrak{su}(2|2)), \sigma) \subset \hat{\mathfrak{f}}^{\perp}$

derivation zd/dz is the generator of Lorentz boosts

The (relativistic) SSSSG Lagrangian

Grigoriev-Tseytlin'08 Mikhailov-SchaferNakemi'08

$$\mathcal{L} = \mathcal{L}_{\text{gWZW}}[G/H] - \frac{k}{\pi} \text{STr} \left(\Lambda \gamma^{-1} \Lambda \gamma \right)$$

$$+ \frac{k}{2\pi} \text{STr} \left(\psi_{+} [\Lambda, D_{-} \psi_{+}] - \psi_{-} [\Lambda, D_{+} \psi_{-}] - 2\psi_{+} \gamma^{-1} \psi_{-} \gamma \right)$$

- $\gamma \in G = e^{\mathfrak{f}_0}$ and $A_{\pm} \longrightarrow$ bosonic fields $\psi_+ \in \mathfrak{f}_1, \ \psi_- \in \mathfrak{f}_3 \longrightarrow$ fermionic fields
- The potential is fixed by $\Lambda \in \mathfrak{f}_2$ (constant)

Gauge symmetry group
$$H = SU(2)^{\times 4} \subset G$$
 $[H, \Lambda] = 0$

The coupling constant is the level of the WZW term $\equiv k$

The (relativistic) SSSSG Lagrangian

Grigoriev-Tseytlin'08 Mikhailov-SchaferNakemi'08

$$\mathcal{L} = \mathcal{L}_{\text{gWZW}}[G/H] - \frac{k}{\pi} \text{STr} \left(\Lambda \gamma^{-1} \Lambda \gamma \right)$$

$$+ \frac{k}{2\pi} \text{STr} \left(\psi_{+} [\Lambda, D_{-} \psi_{+}] - \psi_{-} [\Lambda, D_{+} \psi_{-}] - 2\psi_{+} \gamma^{-1} \psi_{-} \gamma \right)$$

- $\gamma \in G = e^{\mathfrak{f}_0}$ and $A_{\pm} \longrightarrow$ bosonic fields $\psi_+ \in \mathfrak{f}_1, \ \psi_- \in \mathfrak{f}_3 \longrightarrow$ fermionic fields
- The potential is fixed by $\Lambda \in \mathfrak{f}_2$ (constant)

Gauge symmetry group
$$H = SU(2)^{\times 4} \subset G$$
 $[H, \Lambda] = 0$

The coupling constant is the level of the WZW term $\equiv k$