

Title: Statistical Mechanics - Lecture 8


Date: Oct 13, 2011 10:30 AM


URL: http://pirsa.org/11100035


Abstract:















































Pirsa: 11100035 Page 17/144









Pirsa: 11100035 Page 20/144




Pirsa: 11100035 Page 21/144






Pirsa: 11100035 Page 23/144



Pirsa: 11100035 Page 24/144



Pirsa: 11100035 Page 25/144

Liouville's Theorem and conservation of functions of the energy

$$\partial_t \rho(p,q,t) + \sum_{\alpha} [(\partial_{p_{\alpha}} H) \partial_{q_{\alpha}} - (\partial_{q_{\alpha}} H) \partial_{p_{\alpha}}] \rho(p,q,t) = 0$$

If $\rho(p,q,0)$ is any function of the Hamiltonian, e.g. $\rho(p,q,0)=Z^{-1}\exp[-\beta H(p,q)]$ then this same functional form will hold for all times, assuming that the Hamiltonian has no explicit time dependence. $\rho(p,q,0)=f(H(p,q))$ implies that $\rho(p,q,t)=f(H(p,q))$ for any f.


Further, if ρ is any function of a time-independent H and of any other conserved functions of ρ and q, with no explicit time-dependence, then ρ will be a solution of our equation. Thus, not only is the Maxwell- Boltzmann distribution function a solution describing the equilibrium time-dependence of a Hamiltonian system, there are many other solutions as well.

Classical mechanics is not enough to specify a unique equilibrium probability density in a classical system. Something else is needed in addition.


Give some examples of functions of H with and without explicit time-dependence.

14


Pirsa: 11100035 Page 26/144




Pirsa: 11100035 Page 27/144




Pirsa: 11100035 Page 28/144







Pirsa: 11100035 Page 30/144



Pirsa: 11100035 Page 31/144





Pirsa: 11100035 Page 33/144

Liouville's Theorem and conservation of functions of the energy

$$\partial_t \rho(p,q,t) + \sum_{\alpha} [(\partial_{p_{\alpha}} H) \partial_{q_{\alpha}} - (\partial_{q_{\alpha}} H) \partial_{p_{\alpha}}] \rho(p,q,t) = 0$$

If p(p,q,0) is any function of the Hamiltonian, e.g. $p(p,q,0) = Z^{-1} \exp[-\beta H(p,q)]$ then this same functional form will hold for all times, assuming that the Hamiltonian has no explicit time dependence. p(p,q,0)=f(H(p,q)) implies that p(p,q,t)=f(H(p,q)) for any f.


Further, if ρ is any function of a time-independent H and of any other conserved functions of ρ and q, with no explicit time-dependence, then ρ will be a solution of our equation. Thus, not only is the Maxwell- Boltzmann distribution function a solution describing the equilibrium time-dependence of a Hamiltonian system, there are many other solutions as well.

Classical mechanics is not enough to specify a unique equilibrium probability density in a classical system. Something else is needed in addition.


Give some examples of functions of H with and without explicit time-dependence.

14


Pirsa: 11100035 Page 34/144




Pirsa: 11100035 Page 35/144





Pirsa: 11100035 Page 36/144




Pirsa: 11100035 Page 37/144









Pirsa: 11100035 Page 40/144




Pirsa: 11100035 Page 41/144




Pirsa: 11100035 Page 42/144





Pirsa: 11100035 Page 43/144








Pirsa: 11100035 Page 45/144


































Pirsa: 11100035 Page 57/144




Pirsa: 11100035 Page 58/144




Pirsa: 11100035 Page 59/144





Pirsa: 11100035 Page 60/144




Pirsa: 11100035 Page 61/144









Pirsa: 11100035 Page 64/144




Pirsa: 11100035 Page 65/144




Pirsa: 11100035 Page 66/144




Pirsa: 11100035 Page 67/144




Pirsa: 11100035 Page 68/144




Pirsa: 11100035 Page 69/144







Pirsa: 11100035 Page 71/144




Pirsa: 11100035 Page 72/144




Pirsa: 11100035 Page 73/144




Pirsa: 11100035 Page 74/144




Pirsa: 11100035 Page 75/144




Pirsa: 11100035 Page 76/144




Pirsa: 11100035 Page 77/144




Pirsa: 11100035 Page 78/144




Pirsa: 11100035 Page 79/144




Pirsa: 11100035 Page 80/144




Pirsa: 11100035 Page 81/144




Pirsa: 11100035 Page 82/144




Pirsa: 11100035 Page 83/144




Pirsa: 11100035 Page 84/144




Pirsa: 11100035 Page 85/144




Pirsa: 11100035 Page 86/144





Pirsa: 11100035 Page 87/144







Pirsa: 11100035 Page 89/144













Pirsa: 11100035 Page 93/144









Pirsa: 11100035 Page 96/144




Pirsa: 11100035 Page 97/144




Pirsa: 11100035 Page 98/144



Pirsa: 11100035 Page 99/144



Pirsa: 11100035 Page 100/144



The average (<...>) in this equation is over values of $\eta(s)$ for s between t and t+ δt . All other averaging is included in $\rho(p,t)$.

Now expand in δt. (the result is particularly simple because δp does not depend upon p (or qi).

$$\delta t \, \partial_t \, \rho(p,t) \, = \, \langle \delta p \rangle \, \partial_p \, \rho(p,t) + \, \langle \delta p^2 \rangle \, (\partial_p)^2 \, \rho(p,t)/2$$

The average of δp is zero, and the rest gives

$$\partial_t \rho(p,t) = [\langle \delta p^2 \rangle/(2\delta t)] (\partial_p)^2 \rho(p,t)$$

The average has the value:

$$[\delta p]^2 = \int_t^{t+\delta t} \ du \int_t^{t+\delta t} \ ds < \eta(u)\eta(s) > = \int_t^{t+\delta t} \ du \int_t^{t+\delta t} \ ds \ \Gamma \delta(u-s) = \Gamma \delta t$$

(recall that $\leq \eta(t)|\eta(s)> = \Gamma\delta(t-s)$) so that we end up with the result


$$\partial_t \rho(p,t) = \dots + (\Gamma/2) (\partial_p)^2 \rho(p,t)$$

This describes a diffusion in momentum space.


(Notice that because it is diffusive, this result cannot be followed backward in time! Notice also that we had to go to second order in our calculation of the effect of the random kicks because the second order effect here is linear in δt and we were holding on to all linear terms in δt .)

Regists \$52 statement physics Legium Names plus V. Hopping in Manuscaut, Version (1) on (1) 11 from a net source


Pirsa: 11100035 Page 101/144




Pirsa: 11100035 Page 102/144




Pirsa: 11100035 Page 103/144





Pirsa: 11100035 Page 104/144





Pirsa: 11100035 Page 105/144

















Pirsa: 11100035 Page 111/144




Pirsa: 11100035 Page 112/144




Pirsa: 11100035 Page 113/144





Pirsa: 11100035 Page 114/144




Pirsa: 11100035 Page 115/144









Pirsa: 11100035 Page 118/144






Pirsa: 11100035 Page 120/144



Pirsa: 11100035 Page 121/144



Effect of Brownian motion: toward a unique probability distribution

We put together our two different pieces of the Brownian time derivative equation and get:

$$\partial_t \rho(p,x,t) = \dots + (\Gamma/2) (\partial_p)^2 \rho(p,x,t) + \partial_p [c (\partial_p \epsilon) \rho(p,x,t)]$$

The's refer to terms which might come from usual Hamiltonian mechanics. We shall put them aside for a moment. An equation, like the one here, obtained from averaging the stochastic mechanics of a Langevin equation, is called a Fokker-Planck equation.

We look for a time-independent zero of this part of the equation to tell us about the equilibrium behavior. Write $\rho(p,x,t)=\exp[-Q(p,x)+C]$ and find that zeroing this part gives


$$0 = \partial_{p} \left\{ \exp[-Q(p,x)][(\Gamma/2)(-\partial_{p}Q) + \partial_{p}\epsilon(p,r,t) c] \right\}$$
 v.12


which has a solution $Q = \epsilon(p,r,t)$ c. To get the right resusult pick $c=1/(k_BT)$ and pick the other constant of integration so that $\exp(C)=1/z$, which then gives us the right normalization, with $\rho = [\exp(-\epsilon(p,r,t)/(k_BT))]/z$

Einstein's calculation shows that, in order to get the Maxwell-Boltzmann equilibrium result, as a unique result, we have to go beyond Hamiltonian mechanics and include some stochastic behavior. This is a surprise.


21

Nexus 252 statement grevins Lecture Neptes part V. Hopping in Montenum Vet 1000 2.5 (H. 17.17.) 1. Next Kind Lecture






Pirsa: 11100035 Page 125/144



Pirsa: 11100035 Page 126/144



Pirsa: 11100035 Page 127/144

Effect of Brownian motion: toward a unique probability distribution

We put together our two different pieces of the Brownian time derivative equation and get:

$$\partial_t \rho(p,x,t) = \dots + (\Gamma/2) (\partial_p)^2 \rho(p,x,t) + \partial_p [c (\partial_p \epsilon) \rho(p,x,t)]$$

The ...'s refer to terms which might come from usual Hamiltonian mechanics. We shall put them aside for a moment. An equation, like the one here, obtained from averaging the stochastic mechanics of a Langevin equation, is called a Fokker-Planck equation.

We look for a time-independent zero of this part of the equation to tell us about the equilibrium behavior. Write $\rho(p,x,t)=\exp[-Q(p,x)+C]$ and find that zeroing this part gives

$$0 = \partial_{p} \left\{ \exp[-Q(p,x)][(\Gamma/2)(-\partial_{p}Q) + \partial_{p}\varepsilon(p,r,t) c] \right\}$$
 v.12


which has a solution $Q = \epsilon(p,r,t)$ c. To get the right resusult pick $c=1/(k_BT)$ and pick the other constant of integration so that $\exp(C)=1/z$, which then gives us the right normalization, with $\rho = [\exp(-\epsilon(p,r,t)/(k_BT))]/z$

Einstein's calculation shows that, in order to get the Maxwell-Boltzmann equilibrium result, as a unique result, we have to go beyond Hamiltonian mechanics and include some stochastic behavior. This is a surprise.


Streets 252 restators process because Nones part V. Happing in Management Version 5.5 to 5.5 to 5.5 to 5.5 to 6.5 to 6.5

21


Pirsa: 11100035 Page 128/144



Pirsa: 11100035 Page 129/144



Pirsa: 11100035 Page 130/144



Pirsa: 11100035 Page 131/144

Summary of Einstein's dynamics Our fullest time-dependent equation, including both Brownian motion and Hamiltonian mechanics is $\partial_{\tau} \, \rho(\textbf{p},\textbf{r},t) + (\nabla_{\textbf{p}} \, \epsilon(\textbf{p},\textbf{r},t)) \cdot \nabla_{\textbf{r}} \, \rho(\textbf{p},\textbf{r},t) - (\nabla_{\textbf{r}} \, \epsilon(\textbf{p},\textbf{r},t)) \cdot \nabla_{\textbf{p}} \, \rho(\textbf{p},\textbf{r},t)$ v.13a = $(\Gamma/2) (\nabla_p)^2 \rho(\mathbf{p}, \mathbf{r}, t) + \nabla_p \cdot [c (\nabla_p \varepsilon) \rho(\mathbf{p}, \mathbf{r}, t)]$ A. D. Fokker, Die mittlere Energie rotierender elektrischer Dipole im Strahlungsfeld, Ann. Phys., 348 (4, Folge 43), 810-820. (1914). M. Planck, Sitz ber, Preuß, Akad. (1917). Equation v.13a works equally well if we use as our basic variable the density of particles in phase space, f(p,r,t) = Np(p,r,t). We then have $\partial_{\tau} f(\mathbf{p}, \mathbf{r}, t) + \nabla_{\mathbf{p}} \, \epsilon(\mathbf{p}, \mathbf{r}, t) \cdot \nabla_{\mathbf{r}} f(\mathbf{p}, \mathbf{r}, t) - \nabla_{\mathbf{r}} \, \epsilon(\mathbf{p}, \mathbf{r}, t) \cdot \nabla_{\mathbf{p}} f(\mathbf{p}, \mathbf{r}, t)$ v.13b = $(\Gamma/2) (\nabla_p)^2 f(p,r,t) + \nabla_p \cdot [c(\nabla_p \epsilon) f(p,r,t)]$ This all looks very pretty. It is quite general, including Newtonian gravity, electrodynamics, and relativity in the possibilities of $\boldsymbol{\epsilon}$. 22

Pirsa: 11100035 Page 132/144

Summary of Einstein's dynamics Our fullest time-dependent equation, including both Brownian motion and Hamiltonian mechanics is $\partial_{\tau} \, \rho(\textbf{p},\textbf{r},t) + (\nabla_{\textbf{p}} \, \epsilon(\textbf{p},\textbf{r},t)) \cdot \nabla_{\textbf{r}} \, \rho(\textbf{p},\textbf{r},t) - (\nabla_{\textbf{r}} \, \epsilon(\textbf{p},\textbf{r},t)) \cdot \nabla_{\textbf{p}} \, \rho(\textbf{p},\textbf{r},t)$ v.13a = $(\Gamma/2) (\nabla_p)^2 \rho(\mathbf{p},\mathbf{r},t) + \nabla_p \cdot [c (\nabla_p \varepsilon) \rho(\mathbf{p},\mathbf{r},t)]$ A. D. Fokker, Die mittlere Energie rotierender elektrischer Dipole im Strahlungsfeld, Ann. Phys. 348 (4, Folge 43), 810–820. (1914). M. Planck, Sitz ber, Preuß, Akad. (1917). Equation v.13a works equally well if we use as our basic variable the density of particles in phase space, f(p,r,t) = Np(p,r,t). We then have $\partial_{\tau} f(\mathbf{p}, \mathbf{r}, t) + \nabla_{\mathbf{p}} \, \epsilon(\mathbf{p}, \mathbf{r}, t) \cdot \nabla_{\mathbf{r}} f(\mathbf{p}, \mathbf{r}, t) - \nabla_{\mathbf{r}} \, \epsilon(\mathbf{p}, \mathbf{r}, t) \cdot \nabla_{\mathbf{p}} f(\mathbf{p}, \mathbf{r}, t)$ v.13b $= (\Gamma/2) \ (\nabla_p)^2 \ f(\boldsymbol{p},\boldsymbol{r},t) + \nabla_p \cdot \left[c \ (\nabla_p \ \epsilon \) \ f(\boldsymbol{p},\boldsymbol{r},t) \right]$ This all looks very pretty. It is quite general, including Newtonian gravity, electrodynamics, and relativity in the possibilities of $\boldsymbol{\epsilon}$.


Pirsa: 11100035 Page 133/144

The Boltzmann Equation

Long before Einstein's work, Boltzmann had derived the Maxwell-Boltzmann distribution by using a more brute force approach than the one applied by Einstein. Like Einstein, Boltzmann looked at collisions of particles mostly described as moving independently. Therefore these particles could be well-characterized by the free particle Hamiltonian, $p^2/(2m) + U(r)$. In addition, however, Boltzmann imagined that these particles would, occasionally, come close to one another and collide, thereby substantially changing their momentum. So he started from an equation describing first the free particle motion, and then included the effect of collisions. The basic variable in this approach was $f(p_3r,t)$, which would then obey an equation of the form

$$\partial_t f(\mathbf{p}, \mathbf{r}, t) + (\mathbf{p}/m) \cdot \nabla_{\mathbf{r}} f(\mathbf{p}, \mathbf{r}, t) - \nabla_{\mathbf{r}} U(\mathbf{r}, t) \cdot \nabla_{\mathbf{p}} f(\mathbf{p}, \mathbf{r}, t) = \text{effects of collisions}$$


The collisional effects are best shown in a pair of pictures. The number of particles described by (p,r) will diminish because particles with momentum p scatter against particles with momentum q, producing particles with respective momentum p' and and q'. That scattering looks like




particles scatter out of p.

23


Pirsa: 11100035 Page 134/144




Pirsa: 11100035 Page 135/144





Pirsa: 11100035 Page 136/144





Pirsa: 11100035 Page 137/144













Pirsa: 11100035 Page 142/144





Pirsa: 11100035 Page 144/144