Title: Chern-Simons Theory, Vassiliev- Kontsevich Invariant, and Spinfoam Quantum Gravity

Date: Dec 01, 2011 10:30 AM

URL: http://pirsa.org/11120047

Abstract: We construct the q-deformed spinfoam vertex amplitude using Chern-Simons theory on the boundary 3-sphere of the 4-simplex. The rigorous definition involves the construction of Vassiliev-Kontsevich invariant for trivalent knot graph. Under the semiclassical asymptotics, the q-deformed spinfoam amplitude reproduce Regge gravity with cosmological constant at nondegenerate critical configurations.

Outline

- Motivation
- Chern-Simons Theory
- Vassiliev-Kontsevich Invariant
- q-deformed Spinfoam Vertex Amplitude
- Relate to Discrete Gravity with Λ
- Summary

Motivation

- Looking for a mathematical framework for LQG dynamics
- Study the relations between
  - Spinfoam QG in 4d
  - Topological QFT in 3d
  - Conformal Field Theory in 2d
- Construct a finite q-deformed spinfoam model using topological invariant
- Reproduce discrete GR with cosmological constant in the asymptotics.

  (Reproduce Λ-Regge calculus using flat simplices)

Previous works are done by: Turaev, Viro, Mizoguchi, Tada, Crane, Yetter Smolin, Major, Freidel, Krasnov, Noui, Roche Fairbairn, Mensburger, M.H

The vertex amplitude (Euclidean)

$$A_v^q = \int \Psi_\gamma[A^\pm]\; e^{\frac{2\pi i}{k^+} S_{cs}[A^+] + \frac{2\pi i}{k^-} S_{cs}[A^-]}\; DA^+ DA^-$$

(Motivated by L. Smolin '95)

$\Psi_\gamma[A^\pm]$: projective spin-network function on graph γ embedded in S^3 (Dupuis & Livine '10)

A_v^q is rigoronsly defined by Vassiliev - Kontsevich Invariant (Motivated by Fraidel & Krasnov '99)

$$k^\pm = \pm \frac{8}{(1 \pm \beta)^2}\,\omega \qquad \omega = \Lambda \ell_p^2 = \ell_p^2 / \ell_c^2$$

Asymptotic regime: $\ell_p^2 << \text{Area}_f << \ell_c^2$

$$A^q_{4\text{-simplex}} \sim \left(\frac{2\pi}{\lambda}\right)^{12} e^{i\lambda \sum_f \gamma j_f \Theta_f}\; e^{i\omega V_\sigma}\; [1 + o(\tfrac{1}{\lambda})]$$

The analysis can be generalized to simplicial manifold with many 4-simplices.

The vertex amplitude (Euclidean)

$$A_v^q = \int \Psi_\gamma[A^\pm]\; e^{\frac{2\pi i}{k^+} S_{cs}[A^+] + \frac{2\pi i}{k^-} S_{cs}[A^-]}\; DA^+ DA^-$$

(Motivated by L. Smolin '95)

$\Psi_\gamma[A^\pm]$: projective spin-network function on graph γ embedded in S^3 (Dupuis & Livine '10)

A_v^q is rigoronsly defined by Vassiliev - Kontsevich Invariant (Motivated by Fraidel & Krasnov '99)

$$k^\pm = \pm \frac{8}{(1 \pm \beta)^2}\,\omega \qquad \omega = \Lambda \ell_p^2 = \ell_p^2 / \ell_c^2$$

Asymptotic regime: $\ell_p^2 << \text{Area}_f << \ell_c^2$

$$A^q_{4\text{-simplex}} \sim \left(\frac{2\pi}{\lambda}\right)^{12} e^{i\lambda \sum_f \gamma j_f \Theta_f}\; e^{i\omega V_\sigma}\; [1 + o(\tfrac{1}{\lambda})]$$

The analysis can be generalized to simplicial manifold with many 4-simplices.

The vertex amplitude (Euclidean)

$$A_v^q = \int \Psi_\gamma[A^\pm]\; e^{\frac{2\pi i}{k^+} S_{cs}[A^+] + \frac{2\pi i}{k^-} S_{cs}[A^-]}\; DA^+ DA^-$$

(Motivated by L. Smolin '95)

$\Psi_\gamma[A^\pm]$: projective spin-network function on graph γ embedded in S^3 (Dupuis & Livine '10)

A_v^q is rigoronsly defined by Vassiliev - Kontsevich Invariant (Motivated by Fraidel & Krasnov '99)

$$k^\pm = \pm \frac{8}{(1 \pm \beta)^2}\,\omega \qquad \omega = \Lambda \ell_p^2 = \ell_p^2 / \ell_c^2$$

Asymptotic regime: $\ell_p^2 << \text{Area}_f << \ell_c^2$

$$A^q_{4\text{-simplex}} \sim \left(\frac{2\pi}{\lambda}\right)^{12} e^{i\lambda \sum_f \gamma j_f \Theta_f}\; e^{i\omega V_\sigma}\; [1 + o(\tfrac{1}{\lambda})]$$

The analysis can be generalized to simplicial manifold with many 4-simplices.

- The (q-deformed) spinfoam formalism of LQG defines a family of transition amplitudes, which are finite on all possible 2-complexes.

- The (q-deformed) spinfoam amplitude is a mathematical well-defined object, without infinity, and relate GR in the semiclassical limit.

Chern-Simons Theory

* Chern-Simons action on 3-manifold M (TQFT)

$$S_{CS}[A] = \frac{k}{4\pi}\int_M \mathrm{Tr}\left(A\wedge dA + \frac{2}{3}A\wedge A\wedge A\right)$$

A: G-connection, G: compact Lie group

Chern-Simons level/coupling: $k \in \mathbb{Z}$

* Chern-Simons expection value of Links

$$\langle \underset{R_1\ R_2}{\bigcirc\!\bigcirc} \rangle_{CS} = \int \underset{R_1\ R_2}{\bigcirc\!\bigcirc}\ e^{iS_{CS}[A]}\,\mathcal{D}A$$

framed Link L — Wilson loops

with regularization / normalization.

* Skein Relation for Jones Polynomial:

$$-q^{\frac{N}{2}}\langle \times \rangle_{CS} + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})\langle)(\rangle_{CS} + q^{\frac{N}{2}}\langle \times \rangle_{CS} = 0$$

trivial loop: $\langle \bigcirc \rangle_{CS} = \dfrac{q^{\frac{N}{2}} - q^{-\frac{N}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}$

$R = SU(N)$ fundamental rep. $\quad q = e^{\frac{2\pi i}{N+k}}$

(E. Witten '89)



Wilson Lines

EOM of Chern-Simons correlation functions

$$\frac{d}{dt}\left\langle U_1^{R_1}(t)\otimes\cdots\otimes U_n^{R_n}(t)\right\rangle_{CS}$$

$$=\frac{1}{k+N}\sum_{i<j=1}^{n}\frac{\dot z_i-\dot z_j}{z_i-z_j}\,\Omega_{ij}\left\langle U_1^{R_1}(t)\otimes\cdots\otimes U_n^{R_n}(t)\right\rangle_{CS}$$

Knizhnik - Zamolodchikov Equation

z_i: complex coordinate on complex plane Σ. $\dot z = \frac{d}{dt}z(t)$

* Infinitesimal Braiding

$$\Omega_{ij} = \sum_{k=1}^{\dim G} id\otimes\cdots\otimes id\otimes \overset{R_i}{T_k}\otimes\cdots\otimes \overset{R_j}{T_k}\otimes id\otimes\cdots\otimes id$$

$=$  chord diagram

* All the following constructions of invariants and spinfoam amplitude come from the solutions of Knizhnik - Zamolodchikov Equation.

* Knizhnik - Zamolodchikov Equation comes natually from 2d Conformal Field Theory



The operator algebra is a Kac-Moody algebra $\hat{g}_k$

$$[j_m^a, j_n^b] = i \sum_c f^{abc} j_{m+n}^c + k m \delta^{ab} \delta_{m+n,0}$$

$$\left([j_0^a, j_0^b] = i \sum_c f^{abc} j_0^c \right)$$

chiral current : $j^a(z) = \sum_{n \in \mathbb{Z}} j_n^a z^{-n-1}$

central charge : $c = \frac{k \dim G}{k+N}$, k : chern-Simons level

* All the following constructions of invariants and spinfoam amplitude come from the solutions of Knizhnik - Zamolodchikov Equation.

* Knizhnik - Zamolodchikov Equation comes natually from 2d Conformal Field Theory



The operator algebra is a Kac-Moody algebra $\hat{g}_k$

$$[j_m^a, j_n^b] = i \sum_c f^{abc} j_{m+n}^c + k m \delta^{ab} \delta_{m+n,0}$$

$$\left([j_0^a, j_0^b] = i \sum_c f^{abc} j_0^c \right)$$

chiral current : $j^a(z) = \sum_{n \in \mathbb{Z}} j_n^a z^{-n-1}$

central charge : $c = \frac{k \dim G}{k+N}$, k : chern-Simons level

Unitary rep. of the current algebra $\hat{g}_k$

finite # of $\hat{g}_k$-primary fields/states (ground states)

$\xrightarrow{j^a_n}$ descendants of each primary (finite # of $\hat{g}_k$-families)

($\hat{g}_k$-primary fields OPE : $j^a(z)\,\phi^r_R(w) = \frac{1}{z-w}\sum_s (T^a_R)^r_s\,\phi^s_R(w) + \text{regular}$)

$\widehat{su(2)}_k$: finite # of $\hat{g}_k$-families $\rightarrow$ finite # of spins

Constraint on the correlator of $\hat{g}_k$-primary fields

$$\left[\partial_{z_i} - \frac{1}{k+N}\sum_{j\neq i}\frac{\Omega_{ij}}{z_i - z_j}\right]\left\langle \phi_{R_1}(z_1)\cdots\phi_{R_n}(z_n)\right\rangle = 0$$

Knizhnik - Zamolodchikov Equation

(rep. indep.)

* The KZ eqn. $\frac{d}{dt}\langle\ \rangle = \frac{1}{k+N}\sum_{i<j=1}^{n}\frac{\dot{z}_i - \dot{z}_j}{z_i - z_j}$ $\langle\ \rangle$

is a parallel transportation equation with a flat KZ connection on $\mathbb{C}^n$

$$\omega = \frac{1}{k+N}\sum_{i<j}^{n}\frac{\overset{i}{\vert}\text{---}\overset{j}{\vert}}{z_i - z_j}\,(dz_i - dz_j)$$

The solutions of KZ eqn are holonomies of ω.

Unitary rep. of the current algebra $\hat{g}_k$

finite # of $\hat{g}_k$-primary fields/states (ground states)

$\xrightarrow{j^a_n}$ descendants of each primary (finite # of $\hat{g}_k$-families)

($\hat{g}_k$-primary fields OPE : $j^a(z)\,\phi^r_R(w) = \frac{1}{z-w}\sum_s (T^a_R)^r_s\,\phi^s_R(w) + \text{regular}$)

$\widehat{su(2)}_k$: finite # of $\hat{g}_k$-families $\rightarrow$ finite # of spins

Constraint on the correlator of $\hat{g}_k$-primary fields

$$\left[\partial_{z_i} - \frac{1}{k+N}\sum_{j\neq i}\frac{\Omega_{ij}}{z_i - z_j}\right]\left\langle \phi_{R_1}(z_1)\cdots\phi_{R_n}(z_n)\right\rangle = 0$$

Knizhnik - Zamolodchikov Equation

(rep. indep.)

* The KZ eqn. $\frac{d}{dt}\langle\ \rangle = \frac{1}{k+N}\sum_{i<j=1}^{n}\frac{\dot{z}_i - \dot{z}_j}{z_i - z_j}$ $\langle\ \rangle$

is a parallel transportation equation with a flat KZ connection on $\mathbb{C}^n$

$$\omega = \frac{1}{k+N}\sum_{i<j}^{n}\frac{\overset{i}{\vert}\text{---}\overset{j}{\vert}}{z_i - z_j}\,(dz_i - dz_j)$$

The solutions of KZ eqn are holonomies of ω.

Unitary rep. of the current algebra $\hat{g}_k$

finite # of $\hat{g}_k$-primary fields/states (ground states)

$\xrightarrow{j^a_n}$ descendants of each primary (finite # of $\hat{g}_k$-families)

($\hat{g}_k$-primary fields OPE : $j^a(z)\,\phi^r_R(w) = \frac{1}{z-w}\sum_s (T^a_R)^r_s\,\phi^s_R(w) + \text{regular}$)

$\widehat{su(2)}_k$: finite # of $\hat{g}_k$-families $\rightarrow$ finite # of spins

Constraint on the correlator of $\hat{g}_k$-primary fields

$$\left[\partial_{z_i} - \frac{1}{k+N}\sum_{j\neq i}\frac{\Omega_{ij}}{z_i - z_j}\right]\left\langle \phi_{R_1}(z_1)\cdots\phi_{R_n}(z_n)\right\rangle = 0$$

Knizhnik - Zamolodchikov Equation

(rep. indep.)

* The KZ eqn. $\frac{d}{dt}\langle\ \rangle = \frac{1}{k+N}\sum_{i<j=1}^{n}\frac{\dot{z}_i - \dot{z}_j}{z_i - z_j}$ $\langle\ \rangle$

is a parallel transportation equation with a flat KZ connection on $\mathbb{C}^n$

$$\omega = \frac{1}{k+N}\sum_{i<j}^{n}\frac{\overset{i}{\vert}\text{---}\overset{j}{\vert}}{z_i - z_j}\,(dz_i - dz_j)$$

The solutions of KZ eqn are holonomies of ω.

Unitary rep. of the current algebra $\hat{g}_k$

finite # of $\hat{g}_k$-primary fields/states (ground states)

$\xrightarrow{j^a_n}$ descendants of each primary (finite # of $\hat{g}_k$-families)

($\hat{g}_k$-primary fields OPE : $j^a(z)\,\phi^r_R(w) = \frac{1}{z-w}\sum_s (T^a_R)^r_s\,\phi^s_R(w) + \text{regular}$)

$\widehat{su(2)}_k$: finite # of $\hat{g}_k$-families $\rightarrow$ finite # of spins

Constraint on the correlator of $\hat{g}_k$-primary fields

$$\left[\partial_{z_i} - \frac{1}{k+N}\sum_{j\neq i}\frac{\Omega_{ij}}{z_i - z_j}\right]\left\langle \phi_{R_1}(z_1)\cdots\phi_{R_n}(z_n)\right\rangle = 0$$

Knizhnik - Zamolodchikov Equation

(rep. indep.)

* The KZ eqn. $\frac{d}{dt}\langle\ \rangle = \frac{1}{k+N}\sum_{i<j=1}^{n}\frac{\dot{z}_i - \dot{z}_j}{z_i - z_j}$ $\langle\ \rangle$

is a parallel transportation equation with a flat KZ connection on $\mathbb{C}^n$

$$\omega = \frac{1}{k+N}\sum_{i<j}^{n}\frac{\overset{i}{\vert}\text{---}\overset{j}{\vert}}{z_i - z_j}\,(dz_i - dz_j)$$

The solutions of KZ eqn are holonomies of ω.

* Kontsevich Integral (Kontsevich '93)

$$\langle \gamma \rangle = \sum_{n=0}^{\infty} \left(\frac{h}{2\pi i}\right)^n \int_{\substack{t_1 < \cdots < t_n \\ |z_i - z_i'| > \varepsilon}} \sum_{\substack{\text{Pairings} \\ P=(z_i z_i')}} (-1)^{\#P_\phi} D(\gamma, P) \bigwedge_{i=1}^{n} \frac{dz_i - dz_i'}{z_i - z_i'}$$

$\left(h = \frac{2\pi i}{k+N}\right)$

$D(\gamma, P)$ is a chord diagram based on γ



After regularization & normalization,

$\Rightarrow$ Perturbative Vassiliev - Kontsevich invariant

Vassiliev '90 Kontsevich '93 Bar-Natan '95

Vassiliev - Kontsevich Invariant (Combinatorial Definition)

Piunikhin '95, Le & Murakami '97, Murakami & Ohtsuki '97
Altschuler & Freidel '95, Dancso '08

Definition (Chord diagram supported on 3-valent graph)



* There are 3-valent and uni-valent vertices for chords
* The uni-valent chord vertices are on γ

The chord diagrams are perturbative Feynman diagram of Chern-Simons theory.

Definition

Given a 3-valent graph γ embedded in 3-manifold, $\mathcal{A}(\gamma)$ is the $\mathbb{C}$-linear space spanned by the chord diagrams supported by γ, subject to the relations:



(imagine $= f^{abc}$ $= (T^a)^i{}_j$; = intertwiner)

Definition

Given a 3-valent graph γ embedded in 3-manifold, $\mathcal{A}(\gamma)$ is the $\mathbb{C}$-linear space spanned by the chord diagrams supported by γ, subject to the relations:



(imagine  $= f^{abc}$ $= (T^a)^i{}_j$; = intertwiner)

Three building blocks for Vassiliev-Kontsevich invariant

- Braiding $R = \exp \frac{ih}{2}$ 

- Unknot $v =$  $+ \cdots$

- Associator $\Phi =$  $+ \cdots$

Power series in h

Definition (Le & Murakami '97, Murakami & Ohtsuki '97, Dancso '08)

The Vassiliev - Kontsevich invariant

$$\hat{Z} : \gamma \longmapsto \hat{Z}(\gamma) \in \mathcal{A}(\gamma)$$ ← linear space of chord diagram

constructed by the following rules

- Crossings $\hat{Z}(\times) = R \quad \hat{Z}(\times) = R^{-1}$
- Maximum/Minimum $\hat{Z}(\cap) = \nu^{-1} \quad \hat{Z}(\cup) = 1$
- Associator $\hat{Z}(\uparrow\uparrow\uparrow) = \Phi \quad \hat{Z}(\uparrow\uparrow\uparrow) = \Phi^{-1}$
- Vertices $\hat{Z}(\curlyvee) = a^{-1} \quad \hat{Z}(\curlyvee) = 1$

$a =$ 

Definition (Le & Murakami '97, Murakami & Ohtsuki '97, Dancso '08)

The Vassiliev - Kontsevich invariant

$$\hat{Z} : \gamma \longmapsto \hat{Z}(\gamma) \in \mathcal{A}(\gamma)$$ ← linear space of chord diagram

constructed by the following rules

- Crossings $\hat{Z}(\times) = R \quad \hat{Z}(\times) = R^{-1}$
- Maximum/Minimum $\hat{Z}(\cap) = \nu^{-1} \quad \hat{Z}(\cup) = 1$
- Associator $\hat{Z}(\uparrow\uparrow\uparrow) = \Phi \quad \hat{Z}(\uparrow\uparrow\uparrow) = \Phi^{-1}$
- Vertices $\hat{Z}(\curlyvee) = a^{-1} \quad \hat{Z}(\curlyvee) = 1$

$a =$ 

Theorem (Murakami & Ohtsuki '97, Dancso '08)

$\hat{Z}(\gamma)$ is an isotopy invariant of the embedding $\gamma \hookrightarrow S^3$

↓

Diff invariance

Theorem (Murakami & Ohtsuki '97, Dancso '08)

$\hat{Z}(\gamma)$ is an isotopy invariant of the embedding $\gamma \hookrightarrow S^3$

↓

Diff invariance

Definition (Le & Murakami '97, Murakami & Ohtsuki '97, Dancso '08)

The Vassiliev - Kontsevich invariant

$$\hat{Z} : \gamma \longmapsto \hat{Z}(\gamma) \in \mathcal{A}(\gamma)$$ ← linear space of chord diagram

constructed by the following rules

- Crossings $\hat{Z}(\times) = R \quad \hat{Z}(\times) = R^{-1}$
- Maximum/Minimum $\hat{Z}(\cap) = \nu^{-1} \quad \hat{Z}(\cup) = 1$
- Associator $\hat{Z}(\uparrow\uparrow\uparrow) = \Phi \quad \hat{Z}(\uparrow\uparrow\uparrow) = \Phi^{-1}$
- Vertices $\hat{Z}(\curlyvee) = a^{-1} \quad \hat{Z}(\curlyvee) = 1$

$a =$ 

Evaluate the chord diagram with Lie algebra g & rep.





X_a : Lie algebra generator $\quad [X_a, X_b] = f_{abc} X_c$

Quadratic Casimir : $t^{ab} X_a X_b$

3-valent vertices in γ : intertwiners.

$\hat{Z}(\gamma) = \sum_{n=0}^{\infty} \hat{Z}_n(\gamma)\, h^n$ is a power series of chord diagrams

$\hat{Z}(\gamma) \xrightarrow[\omega_g]{\text{Evaluation with } g \text{ order by order}} \omega_g \circ \hat{Z}(\gamma)$

Quantum Group invariant

Evaluate the chord diagram with Lie algebra g & rep.



X_a : Lie algebra generator $\quad [X_a, X_b] = f_{abc} X_c$

Quadratic Casimir : $t^{ab} X_a X_b$

3-valent vertices in γ : intertwiners.

$\hat{Z}(\gamma) = \sum_{n=0}^{\infty} \hat{Z}_n(\gamma)\, h^n$ is a power series of chord diagrams

$\hat{Z}(\gamma) \xrightarrow[\omega_g]{\text{Evaluation with } g \text{ order by order}} \omega_g \cdot \hat{Z}(\gamma)$

Quantum Group invariant

q-deformed Spinfoam Amplitude

Heuristically, the vertex amplitude (Euclidean)

$$A_v^q = \int \Psi_\gamma[A^\pm]\, e^{\frac{2\pi i}{h^+} S_{cs}[A^+] + \frac{2\pi i}{h^-} S_{cs}[A^-]}\, DA^+ DA^-$$

(A^+, A^-) : $SU(2) \times SU(2)$ - connection

S_{cs} : Chern-Simons action on S^3

$$h^\pm = \pm \frac{8}{(1\pm\beta)^2}\,\omega \qquad \omega = \Lambda \ell_p^2 = \ell_p^2 / \ell_c^2$$

$\Psi_\gamma[A^\pm]$: projective spin-network function on graph γ embedded in S^3

Given an $SU(2)$ spin-network function ψ_γ with $(j_e\ i_v)$

EPRL map : $j_e \longmapsto j_e^\pm = \frac{|1\pm\beta|}{2} j_e$ (spin(4) spins)

$$i_v \longmapsto I_v = i_v^{a_1\cdots a_n} C^{b_1^+ b_1^-}_{a_1} \cdots C^{b_n^+ b_n^-}_{a_n} P^{c_1^\pm \cdots c_n^\pm}_{b_1^\pm \cdots b_n^\pm}$$

($Spin(4)$ intertwiner)

$C_a^{b^+ b^-}(j, j^+, j^-)$ CG coefficients

Projector to the space of $Spin(4)$ intertwiners

$\longrightarrow$ Construct an $Spin(4)$ spin-network Ψ_γ from ψ_γ

Dupuis & Livine '10

The (arbitrary valent) vertex amplitude $A_v^q = A_v^q[\psi_\gamma]$ is a functional on $CYL \subset \mathcal{H}_{kin}$ via Chern-Simons and EPRL map.

Motivated by L. Smolin '95

Rigorous definition: $A_v^q = \omega_{Spin(4)} \circ \hat{Z}(\gamma)_{h^+, h^-}$

↑

Evaluation using Spin(4) rep & intertwiner which are images of EPRL map.

Relate to Gambini, Griego, Pullin '98

Generalize to Lorentzian theory:

$$A_v^q = \omega_{SL(2,\mathbb{C})} \circ \hat{Z}(\gamma)_h$$

A finite spinfoam amplitude on a 2-complex

$$A^q(K) = \sum_{j_f\, i_e} \prod_f \mu(j_f) \prod_v A_v^q(j_f, i_e)$$

The (arbitrary valent) vertex amplitude $A_v^q = A_v^q[\psi_\gamma]$ is a functional on $CYL \subset \mathcal{H}_{kin}$ via Chern-Simons and EPRL map.

Motivated by L. Smolin '95

Rigorous definition: $A_v^q = \omega_{Spin(4)} \circ \hat{Z}(\gamma)_{h^+, h^-}$

↑ Evaluation using Spin(4) rep & intertwiner which are images of EPRL map.

Relate to Gambini, Griego, Pullin '98

Generalize to Lorentzian theory:

$$A_v^q = \omega_{SL(2,\mathbb{C})} \circ \hat{Z}(\gamma)_h$$

A finite spinfoam amplitude on a 2-complex

$$A^q(K) = \sum_{j_f\, i_e} \prod_f \mu(j_f) \prod_v A_v^q(j_f, i_e)$$

Asymptotics (Euclidean, 4-simplex amplitude) M.H '11

Asymptotic regime: $\ell_p^2 \ll \text{Area}_f \ll \ell_c^2$

We keep triangle area fixed, but scale

$$\ell_p^2 \to \lambda^{-1} \ell_p^2$$
$$\ell_c^2 \to \lambda \ell_c^2 \qquad \lambda \to \infty$$

From $\text{Area} = j_f \ell_p^2$, $\omega = \ell_p^2 / \ell_c^2$

$$j_f \to \lambda j_f \qquad \omega \to \lambda^{-2} \omega$$

Asymptotics (Euclidean, 4-simplex amplitude) M.H '11

Asymptotic regime: $\ell_p^2 \ll \text{Area}_f \ll \ell_c^2$

We keep triangle area fixed, but scale

$$\ell_p^2 \to \lambda^{-1} \ell_p^2$$
$$\ell_c^2 \to \lambda \ell_c^2 \qquad \lambda \to \infty$$

From $\text{Area} = j_f \ell_p^2$, $\omega = \ell_p^2 / \ell_c^2$

$$j_f \to \lambda j_f \qquad \omega \to \lambda^{-2} \omega$$

Consider the expansion and the scaling $j \to \lambda j$, $\omega \to \omega \lambda^{-2}$

- Braiding $R = \exp \frac{ih}{2}$  $\sim 1 + o(1)$

- Unknot $\nu =$  $\sim 1 + o(\frac{1}{\lambda})$

- Associator $\Phi =$  $\sim 1 + o(\frac{1}{\lambda})$

with a certain nondeg. boundary condition of 4-simplex

$$A^{q}_{4\text{-simplex}} \sim \left(\frac{2\pi}{\lambda}\right)^{12} e^{i\lambda \sum_f \gamma j_f \Theta_f} e^{i\omega V_\sigma} \left[1 + o(\tfrac{1}{\lambda})\right]$$

contribution from R. ; ν, Φ

Consider the expansion and the scaling $j \to \lambda j$, $\omega \to \omega \lambda^{-2}$

- Braiding $R = \exp \frac{ih}{2}$  $\sim 1 + o(1)$

- Unknot $\nu =$  $\sim 1 + o(\frac{1}{\lambda})$

- Associator $\Phi =$  $\sim 1 + o(\frac{1}{\lambda})$

with a certain nondeg. boundary condition of 4-simplex

$$A^{q}_{4\text{-simplex}} \sim \left(\frac{2\pi}{\lambda}\right)^{12} e^{i\lambda \sum_f \gamma \hat{J}_f \Theta_f} e^{i\omega V_\sigma} \left[1 + o(\tfrac{1}{\lambda})\right]$$

contribution from R. ; ν, Φ

Some details:

$$A^{g}_{4\text{-simplex}} = \int \prod_{a=1}^{5} dg_a^{+} dg_a^{-} \; K^{\omega} \prod_{a<b} \langle j^{\pm}_{ab}, n_{ab} | (g_a^{\pm})^{-1} g_b^{\pm} | j^{\pm}_{ab}, n_{ba} \rangle$$

K^{ω}: expansion of chord diagrams (Feynman diagrams)

$\prod_{a<b} \langle \cdots \rangle$: usual Euclidean spinfoam integrand

From K^{ω}

n univalent vertices





$$\frac{\langle j_{ab}\, n_{ab} \,|\, g_a^{-1} X_{k_1} \cdots X_{k_n} g_b \,|\, j_{ab}\, n_{ba} \rangle}{\langle j_{ab}\, n_{ab} \,|\, g_a^{-1} g_b \,|\, j_{ab}\, n_{ba} \rangle} \qquad j_{ab} \to \lambda j_{ab}$$

$\lambda \to \infty$

$$= \lambda^{j_{ab}} (g_b\, n_{ba})^{k_1} \cdots \lambda^{j_{ab}} (g_b\, n_{ba})^{k_n} \left[1 + o\left(\tfrac{1}{\lambda}\right) \right] \sim \lambda^{n}$$

A chord diagram with n uni-valent & m tri-valent vertices of order $h^{\frac{m+n}{2}}$ ($h \propto \omega$), contributing $\lambda^{-(m+n)}$ from coupling constant

→ a diagram with m contribut λ^{-m}

→ leading contribution: $m = 0$ only comes from R-matrix



$$A^{q}_{4\text{-simplex}} \sim \left(\frac{2\pi}{\lambda}\right)^{12} e^{i\lambda \sum_f \gamma j_f \Theta_f} \; e^{i\omega V_\sigma} \left[1 + o\left(\frac{1}{\lambda}\right)\right]$$

Λ-Regge action with flat simplex

consistent with $\ell_p^2 \ll \ell_\sigma^2 \ll \ell_c^2$

↑

scale of 4-simplex

On a simplicial complex K

$$A^q(K) = \sum_{j_f\, i_e} \prod_f \mu(j_f) \prod_v A_v^q(j_f, i_e)$$

Asymptotics of spinfoam amplitude on a simplicial complex

Classify the critical configurations into 5 types



M.H & M.Zhang '11.

Recall the asymptotic formula:

$$f(\lambda) = \int dx\ a(x)\, e^{\lambda S(x)} \qquad \text{with } \mathrm{Re}\, S \le 0$$

As $\lambda \to \infty$

$$f(\lambda) = \sum_{x_c} a(x_c) \left(\frac{2\pi}{\lambda}\right)^{\frac{\gamma(x_c)}{2}} \frac{e^{i\,\mathrm{Ind}\, H'(x_c)}}{\sqrt{|\det{}_r H'(x_c)|}}\, e^{\lambda S(x_c)} \left[1 + o\left(\tfrac{1}{\lambda}\right)\right]$$

on different type of geometries: ↑ action at critical point

Nondeg, $V_4 > 0$ or $V_4 < 0$

$$e^{\lambda S(x_c)} = e^{-i\lambda\left[\varepsilon\, \mathrm{sgn} V_4 \sum_f \gamma\, j_f \textcircled{H}_f + \pi \sum_e n_e \sum_{f \subset t_e} j_f\right]}\, e^{i \sum_v \omega V_v}$$

Deg-A, $V_4 > 0$ or $V_4 < 0$ (asymptotics of BF type)

$$e^{\lambda S(x_c)} = e^{-i\lambda\left[\varepsilon\, \mathrm{sgn} V_4 \sum_f j_f \textcircled{H}_f + \pi \sum_e n_e \sum_{f \subset t_e} j_f\right]}$$

Deg-B

$$e^{\lambda S(x_c)} = e^{-i\lambda \sum_f j_f \Phi_f}$$ ← an angle from vector geometry.

Summary

- We formulate the spinfoam vertex amplitude in terms of Chern-Simons theory and Vassiliev-Kontsevich invariant.

- The constuction is essentially the solution of Knizhnik-Zamolodchikov Equation, with regularization & normalization, combined with EPRL map.

- The resulting spinfoam model is finite, and relate discrete GR with Λ in the regime $\ell_p^2 \ll \ell_\sigma^2 \ll \ell_c^2$.

Some details:

$$A^{g}_{4\text{-simplex}} = \int \prod_{a=1}^{5} dg_a^{+} dg_a^{-} \; K^{\omega} \prod_{a<b} \langle j^{\pm}_{ab}, n_{ab} | (g_a^{\pm})^{-1} g_b^{\pm} | j^{\pm}_{ab}, n_{ba} \rangle$$

K^{ω}: expansion of chord diagrams (Feynman diagrams)

$\prod_{a<b} \langle \cdots \rangle$: usual Euclidean spinfoam integrand

From K^{ω}

n univalent vertices





$$\frac{\langle j_{ab}\, n_{ab} | \, g_a^{-1} X_{k_1} \cdots X_{k_n} g_b \, | j_{ab}\, n_{ba} \rangle}{\langle j_{ab}\, n_{ab} | \, g_a^{-1} g_b \, | j_{ab}\, n_{ba} \rangle} \qquad j_{ab} \to \lambda j_{ab}$$

$$\lambda \to \infty \quad = \lambda j_{ab} (g_b n_{ba})^{k_1} \cdots \lambda j_{ab} (g_b n_{ba})^{k_n} \left[1 + o\left(\tfrac{1}{\lambda}\right) \right] \sim \lambda^{n}$$

A chord diagram with n uni-valent & m tri-valent vertices of order $h^{\frac{m+n}{2}}$ ($h \propto \omega$), contributing $\lambda^{-(m+n)}$ from coupling constant

→ a diagram with m contribut λ^{-m}

→ leading contribution: $m = 0$ only comes from R-matrix