

Title: Supersymmetric QCD Corrections to $b \bar{g} \hat{\tau}^+ b h$


Date: Dec 12, 2011 01:00 PM

URL: <http://pirsa.org/11120056>

Abstract: The dominant production mechanism for Standard Model (SM) Higgs boson is $g \bar{g} \hat{\tau}^+ h$. However, in certain beyond the SM scenarios, Higgs production



The Standard Model



- The only missing piece in the Standard Model - the Higgs boson.

The Higgs Boson

- How can we find the Higgs boson?

MISSING PARTICLE:

Name: *Higgs boson*

Age: *13.7 billion years*

Missing: *45 years*


Birthday: *Every few days at Fermilab*

Favorite trait: *Mass*

Favorite particle: *top quark*






Favorite Hangout: *Tevatron*

The Standard Model Higgs Boson



- Electroweak Precision measurements \Rightarrow a light Higgs boson
- LEP : $e^+ e^- \rightarrow Z h (h \rightarrow b\bar{b}/\tau^+ \tau^-)$
 - Bound : $m_h > 114$ GeV


SM Higgs boson Production at Hadron Colliders



- Main production modes in SM :
 - “gluon fusion”
 - W/Z fusion
 - W/Z associated production
 - associated production with t

Mirror Screens

SM Higgs boson Decay




- For small masses ($m_h < 2m_Z$) :
 - $h \rightarrow b\bar{b}/\tau^+\tau^-$
 - $h \rightarrow WW^*/ZZ^*$
 - $h \rightarrow \gamma\gamma$ (loop suppressed)
- For large masses
 - $h \rightarrow WW/ZZ$

The Standard Model Higgs Boson

What does LHC tell us about the Higgs?

- LHC : 5 fb^{-1} in 1 year at $\sqrt{s} = 7 \text{ TeV}$



Mirror Screens

- What about Higgs production beyond standard model?
- Is $bg \rightarrow bh$ even relevant compared to gluon fusion?

Prerit Jaiswal (YITP and BNL)

SQCD $bg \rightarrow bh$



Mirror Screens

Outline

- Motivation for $bg \rightarrow bh$ studies
- 4FNS vs 5FNS
- Large $\tan \beta$ radiative corrections
- SQCD Corrections and Δ_b Approximation
- How good is the Δ_b Approximation?

Preit Jaiswal (YITP and BNL)

SQCD $bg \rightarrow bh$



MSSM Higgs Sector

- Two Higgs doublets H_u and $H_d \Rightarrow$
 - 2 CP even (h^0, H^0), 1 CP odd (A^0)
 - 1 charged Higgs (H^\pm)
 - Goldstone bosons (G^0, G^\pm)
- VEVs : $\langle H_u \rangle = v_u$ and $\langle H_d \rangle = v_d$, $\tan \beta = v_u/v_d$
- Two independent parameters : m_A and $\tan \beta$ (at tree level)
 - examples : $m_{h/H}^2 = \frac{1}{2} \left[m_A^2 + m_Z^2 \mp \sqrt{(m_A^2 + m_Z^2)^2 - 4m_A^2 m_Z^2 \cos^2 2\beta} \right]$
 - α diagonalizes the Higgs mass matrix

$$\cos 2\alpha = -\cos 2\beta \left(\frac{m_A^2 - m_Z^2}{m_H^2 - m_h^2} \right)$$
- Decoupling limit : $m_A \gg m_Z$
 - $\beta - \alpha \rightarrow \frac{\pi}{2}$
 - Significance of Decoupling limit?



MSSM Higgs Couplings to Fermions


	Couplings w.r.t SM	Decoupling limit
$h^0 \bar{b}b$	$-\sin \alpha / \cos \beta$	1
$h^0 \bar{t}t$	$\cos \alpha / \sin \beta$	1
$H^0 \bar{b}b$	$\cos \alpha / \cos \beta$	$\tan \beta$
$H^0 \bar{t}t$	$\sin \alpha / \sin \beta$	$-1 / \tan \beta$
$A^0 \bar{b}b$	$\gamma_5 \tan \beta$	$\gamma_5 \tan \beta$
$A^0 \bar{t}t$	$\gamma_5 \cot \beta$	$\gamma_5 \cot \beta$

- Decoupling limit : h^0 couplings in MSSM same as that of SM at tree level (also true for couplings with gauge bosons)
 - Production cross-sections same as that in SM
 - Same channels as SM : gluon fusion, vector boson fusion, associated production
- For $m_A \gtrsim m_h$ and large $\tan \beta$, same conclusions hold (at tree level).

Mirror Screens

MOTIVATION

- Current status of MSSM parameter space




- LEP exclusion : small $\tan\beta$ (\Rightarrow small m_h)
- Tevatron exclusion : $h \rightarrow \tau^+\tau^- / b\bar{b}$

- Decoupling limit / moderately heavy A and $\tan\beta \gtrsim 7$ good parameter space to study.

- Assuming the allowed parameter space (Decoupling limit or moderately heavy A and $\tan \beta \gtrsim 7$), it seems the MSSM lightest CP even Higgs production should be identical to SM.
- Then why (and under what conditions) is production of Higgs in association with bottom quarks important?
- For large $\tan \beta$, the bottom Yukawa gets significantly modified.
 - At tree level, it is SM like
 - Radiative corrections can cause Yukawa to rescale by a factor of ~ 3 .
- For $m_A \lesssim 200$ GeV and $\tan \beta \gtrsim 7$, the dominant Higgs production mode is in association with bottom quark.


- Higgs production in association with bottom quarks



- $b\bar{b}h$ final state measurements :
 - Exclusive $b\bar{b}h$ production : Both b tagged (high p_T)
 - Semi-inclusive : At least one b tagged
 - Inclusive : No b tagged
- Inclusive modes \rightarrow Larger Cross-section but also large backgrounds.
 - \Rightarrow b tagging reduces background
 - \Rightarrow Semi-inclusive mode : a good compromise

4FNS (4-parton Flavour Number Scheme)


- Initial state quarks : Four lightest flavours
- *Advantage* :
 - Only way to calculate exclusive $b\bar{b}h$ production (both b tagged)
- *Drawback* :
 - In inclusive $b\bar{b}h$ production, large collinear logs
 - “Fixed order” perturbation theory (No resummation)



$$\Lambda_b = \ln \left(\frac{\mu^2}{m_b^2} \right)$$

Mirror Screens


4FNS VS 5FNS



Mirror Screens

4FNS VS 5FNS

- 4FNS and 5FNS do not match at LO but must match if calculated to all orders.




R. Harlander, M. Kramer, and M. Schumacher,

<https://twiki.cern.ch/twiki/pub/LHCPhysics/MSSMNeutral/santandermatching-hks.pdf>

Effective Lagrangian Approach

$$\mathcal{L}_{\text{eff}} = -\lambda_b \bar{b}_R \left[H_d + \frac{\Delta\lambda_b}{\lambda_b} H_u^* \right] Q_L$$

- Second term not allowed at tree level in MSSM.
- Effective theory : Integrate out heavy degrees of freedom (\tilde{g}, \tilde{b})
- Loop corrections enhanced since $v_u = v_d \tan\beta$




$$m_b \rightarrow \lambda_b v_d + \Delta\lambda_b v_u$$

$$= \lambda_b v_d \left(1 + \frac{\Delta\lambda_b}{\lambda_b} \tan\beta \right)$$

Effective Yukawa Coupling

- $m_b \rightarrow \lambda_b v_d (1 + \Delta_b)$
- At one-loop, $\Delta_b \propto \frac{\alpha_s}{\pi} \mu \tan\beta / M_{SUSY}$
- For $\mu \sim M_{SUSY}$, large $\tan\beta$
 $\Rightarrow \Delta_b \sim \mathcal{O}(1)$
- Perturbation in $\alpha_s \tan\beta$, must resum
- Resummation easy since no loop contribution to $\alpha_s^n \tan^n \beta$ for $n \geq 2$!!



- Final result : Bare b quark mass in effective Lagrangian approach

$$m_b = \frac{m_b}{(1+\Delta_b)},$$

- effective Yukawa coupling :

$$\lambda_b = \frac{m_b}{v_d(1+\Delta_b)} \left(1 - \frac{\Delta_b}{\tan\alpha \tan\beta} \right), \text{ where}$$

$$\Delta_b = \frac{2\alpha_s}{3\pi} M_{\tilde{g}} \mu \tan\beta \mathcal{F}(M_{\tilde{b}_1}, M_{\tilde{b}_2}, M_{\tilde{g}})$$


Mirror Screens

SQCD corrections


- Complete SQCD one-loop corrections to $bg \rightarrow bh$ computed in [S. Dawson and C. Jackson, Phys. Rev. D77, 015019 (2008)] , [S. Dawson, C. Jackson and P.J., Phys. Rev. D77, 015019 Phys.Rev. D83 (2011) 115007]
- SUSY QCD \Rightarrow Diagrams involving gluinos and sbottoms



Self-Energy diagrams



hbb Vertex diagrams



gbb Vertex diagrams


SQCD $bg \rightarrow bh$

10/14/2011

22 / 28

Mirror Screens

SQCD corrections



Box diagrams

..... and counterterms

- For cross-section calculations, consider two cases :

- Maximal mixing case :
 $|\tilde{m}_L^2 - \tilde{m}_R^2| \ll m_b |X_b| \Rightarrow \theta_{\tilde{b}} \sim \frac{\pi}{4}$
- Minimal mixing case :
 $|\tilde{m}_L^2 - \tilde{m}_R^2| \gg m_b |X_b| \Rightarrow \theta_{\tilde{b}} \sim 0$

- Squark mass matrix :


$$M_{\tilde{b}}^2 = \begin{pmatrix} \tilde{m}_L^2 & m_b X_b \\ m_b X_b & \tilde{m}_R^2 \end{pmatrix}$$

$$X_b = A_b - \mu \tan \beta$$

- Mixing angle :

$$\sin 2\theta_{\tilde{b}} = \frac{2m_b(A_b - \mu \tan \beta)}{M_{\tilde{b}_1}^2 - M_{\tilde{b}_2}^2}$$

7 TeV LHC, $bg \rightarrow bh$
 $\tan \beta = 40, \mu = M_A = 1 \text{ TeV}, M_S = 2M_{\text{gluino}} = m_L$



- Maximal mixing case :
 $m_R = m_L = M_S$
- Minimal mixing case :
 $m_R = \sqrt{2}M_L = \sqrt{2}m_S$

- How much of the SQCD corrections accounted by Δ_B approximation?

SQCD corrections : Δ_b Approximation

- Δ_b Approximation : Calculate SM cross-section but with rescaled Yukawa coupling

$$\frac{g_{bbh}^{MSSM}}{g_{bbh}^{SM}} = -\frac{\sin \alpha}{\cos \beta} \left(\frac{1}{1 + \Delta_b} \right) \left(1 - \frac{\Delta_b}{\tan \beta \tan \alpha} \right)$$

- Includes only resummed large $\tan \beta$ effects.

$M_{SUSY} = 1$ TeV, $M_0 = 1$ TeV, $A_b = A_t = 2$ TeV, and $\mu = M_2 = 200$ GeV

tan $\beta = 40$				
M_{h^0} [GeV]	100	110	120	130
α [rad]	-1.5063	-1.4716	-1.3798	-0.7150
$g_{bbh^0}^{MSSM} / g_{bbh}^{SM}$	33.913	33.823	33.387	22.390
$g_{ttb^0}^{MSSM} / g_{ttb}^{SM}$	0.0645	0.0991	0.1899	0.7553


S. Dawson, C.B. Jackson, L. Reina, D. Wackerth, Mod.Phys.Lett. A21 (2006) 89-110

SQCD corrections : Δ_b Approximation

- Δ_b Approximation in the context of $bg \rightarrow bh$:

$$\sigma_{MSSM} = \sigma_{SM} \left(\frac{g_{bbh}^{MSSM}}{g_{bbh}^{SM}} \right)^2 \equiv \sigma_{IBA}$$

7 TeV LHC, $bg \rightarrow bh$
 $\tan \beta=40, \mu=1000 \text{ GeV}, M_{\tilde{\Lambda}}=500 \text{ GeV}, m_{\tilde{\chi}}=700 \text{ GeV}$



- IBA works very well !!
- Other SQCD corrections $< 2 \%$

Mirror Screens


Δ_b Approximation : Why it works?

[S. Dawson, C. Jackson and PJ, Phys. Rev. D77, 015019 Phys.Rev. D83 (2011) 115007]

- Analytically compute the SQCD Corrections for two extreme cases : maximal and minimal mixing
- Maximal mixing : $|\tilde{m}_L^2 - \tilde{m}_R^2| \ll m_b |X_b| \Rightarrow \theta_{\tilde{b}} \sim \frac{\pi}{4}$
- Expand the amplitudes in power of $1/M_S$ assuming large $\tan \beta$ so that the terms like $m_b \tan \beta \sim \mathcal{O}(M_{EW})$.

$$|\overline{\mathcal{A}}|_{max}^2 = -\frac{2\pi\alpha_s(\mu_R)}{3} (g_{b\tilde{b}\tilde{h}}^{\Delta_b})^2 \left\{ \left(\frac{a^2 + M_h^4}{st} \right) \left[1 + 2 \left(\frac{\delta g_{b\tilde{b}\tilde{h}}}{g_{b\tilde{b}\tilde{h}}} \right)_{max}^{(2)} \right] + \frac{\alpha_s(\mu_R)}{2\pi} \frac{M_h^2}{M_S^2} \delta K_{max} \right\} + \mathcal{O} \left(\left[\frac{M_{EW}}{M_S} \right]^4 \cdot \alpha_s^3 \right).$$

- First term : IBA
- Other terms are $\mathcal{O}(M_{EW}^2/M_S^2)$



$$R = M_{\tilde{R}}/M_S$$

10/14/2011

27 / 28

- Higgs production in association with b quark an important mode for large parameter space in MSSM
- SUSY QCD corrections to $bg \rightarrow bh$ can be big for large $\tan \beta$.
- Bottom Yukawa coupling scales by a large factor when large $\tan \beta$ corrections resummed.
- SUSY QCD corrections are dominated by this rescaling : Δ_b approximation works.