

Title: The Other 99%: The Life and Times of the Black Hole Next Door

Date: Feb 15, 2012 02:00 PM

URL: <http://pirsa.org/12020127>

Abstract: Black holes lead a double life, simultaneously the dark, enigmatic, hairless consequence of general relativity and the powerful engines lurking at the hearts of galaxies with cosmological consequences.

$$L \sim 10^{48} \text{ erg/s}$$

- 10^{30} Canadas
- 10^{13} Suns
- 1 SNe/15 min
- $1 M_{\text{sun}} c^2/\text{month}$

PI Colloquium, Feb. 15, 2012

$$\frac{d}{d\sigma}(X \cdot X)$$

$$T_{00} = \dot{X}^2 + \frac{1}{2}(-\dot{X}^2 + X'^2) = \frac{1}{2}(\dot{X}^2 + X'^2)$$

check ||

$$T_{11}$$

$$T_{01} = T_{10} = X' \cdot \dot{X}$$

$$= 0$$

$$= 0$$

$$\pm = \frac{1}{2}(\tau \pm \sigma) \quad X_{(\sigma^{\pm}, \tau^{\pm})}'' = X_R''(\sigma^-) + X_L''(\sigma^+)$$

Closed string condition

$$X_{(\tau, \sigma)}'' = X_{(\tau)}''$$

$$(m\sigma)$$

$$\partial_- X'' = \partial_- X_R''$$

$$\partial_+ X'' = \partial_+ X_L''$$

The Other 99%: The Life and Times of the Black Hole Next Door

Avery E Broderick



Sheperd Doeleman (MIT Haystack)
Vincent Fish (MIT Haystack)
Alan Rogers (MIT Haystack)
Dimitrios Psaltis (Arizona)
Tim Johannsen (Arizona)

Avi Loeb (Harvard)
Ramesh Narayan (Harvard)
Jonathan Bittner (Harvard)
Mark Reid (Smithsonian)
Mareki Honma (NAOJ)

PI Colloquium, Feb. 15, 2012



Black Holes


$$R_{\mu\nu} = 0$$

$$g_{\mu\nu}(M, a, Q)$$

$$r_+, r_e$$

$$r_\gamma, r_{ISCO}$$







PI Colloquium, Feb. 15, 2012



View from above showing the shapes of



Black Holes + Stuff



PI Colloquium, Feb. 15, 2012


Black Holes as Engines



$$L \sim 10^{47} \text{ erg/s}$$

PI Colloquium, Feb. 15, 2012

Black Holes as Engines



$$L \sim 10^{48} \text{ erg/s}$$

- 10^{30} Canadas

PI Colloquium, Feb. 15, 2012

Black Holes as Engines




$$L \sim 10^{48} \text{ erg/s}$$

- 10^{30} Canadas
- 10^{13} Suns
- 1 SNe/15 min
- $1 M_{\text{sun}} c^2/\text{month}$


PI Colloquium, Feb. 15, 2012

The Story: Accretion Power




PI Colloquium, Feb. 15, 2012

The Story: Accretion Power



PI Colloquium, Feb. 15, 2012

The Story: Accretion Power




PI Colloquium, Feb. 15, 2012

The Story: Accretion Power

$$E_M \sim \frac{GM_E m}{R_E} \sim 10^{-9} mc^2$$

$$E_M \sim \frac{GMm}{R_{BH}} \sim \frac{GMm}{(GM/c^2)} \sim mc^2$$

$$\rightarrow L \sim \dot{M} c^2$$



PI Colloquium, Feb. 15, 2012

How Bright can BHs Get?

$$T \rightarrow c_s \sim T^{1/2}$$

1. Bondi Accretion:



PI Colloquium, Feb. 15, 2012

How Bright can BHs Get?

$$T \rightarrow c_s \sim T^{1/2}$$

1. Bondi Accretion:



PI Colloquium, Feb. 15, 2012

How Bright can BHs Get?

1. Bondi Accretion:

$$T \rightarrow c_s \sim T^{1/2}$$

$$v_{esc} < c_s$$

Unbound

$$v_{esc} > c_s$$

Bound

PI Colloquium, Feb. 15, 2012

How Bright can BHs Get?

1. Bondi Accretion:

$$\dot{M} \sim \rho \, 4\pi r_B^2 \, c_s \propto \frac{M^2}{T^{3/2}}$$

$$T \rightarrow c_s \sim T^{1/2}$$

$v_{esc} < c_s$
Unbound

$v_{esc} > c_s$
Bound


PI Colloquium, Feb. 15, 2012

How Bright can BHs Get?

1. Bondi Accretion:

$$\dot{M} \sim \rho \, 4\pi r_B^2 \, c_s \propto \frac{M^2}{T^{3/2}}$$

2. Eddington Limit:




PI Colloquium, Feb. 15, 2012

How Bright can BHs Get?

1. Bondi Accretion:

$$\dot{M} \sim \rho \, 4\pi r_B^2 \, c_s \propto \frac{M^2}{T^{3/2}}$$

2. Eddington Limit:




PI Colloquium, Feb. 15, 2012

How Bright can BHs Get?

1. Bondi Accretion:

$$\dot{M} \sim \rho \, 4\pi r_B^2 \, c_s \propto \frac{M^2}{T^{3/2}}$$

2. Eddington Limit:




$$f \sim \frac{L\sigma}{4\pi r^2 c} = f_g = \frac{GMm_p}{r^2}$$

How Bright can BHs Get?

1. Bondi Accretion:

$$\dot{M} \sim \rho \, 4\pi r_B^2 \, c_s \propto \frac{M^2}{T^{3/2}}$$

2. Eddington Limit:



$$f \sim \frac{L\sigma}{4\pi r^2 c} = f_g = \frac{GMm_p}{r^2}$$

$$\rightarrow L \sim \frac{4\pi GMm_p c}{\sigma} = 1.3 \times 10^{38} \left(\frac{M}{M_{sun}} \right) \text{ erg/s}$$

PI Colloquium, Feb. 15, 2012


How Bright can BHs Get?

1. Bondi Accretion:

$$\dot{M} \sim \rho \, 4\pi r_B^2 \, c_s \propto \frac{M^2}{T^{3/2}}$$

2. Eddington Limit:

$$\dot{M}_{Edd} \sim 2 \times 10^{-9} \left(\frac{M}{M_{sun}} \right) M_{sun}/yr$$



$$f \sim \frac{L\sigma}{4\pi r^2 c} = f_g = \frac{GMm_p}{r^2}$$

$$\rightarrow L \sim \frac{4\pi GMm_p c}{\sigma} = 1.3 \times 10^{38} \left(\frac{M}{M_{sun}} \right) \text{erg/s}$$

PI Colloquium, Feb. 15, 2012


How Bright can BHs Get?

1. Bondi Accretion:

$$\dot{M} \sim \rho \, 4\pi r_B^2 \, c_s \propto \frac{M^2}{T^{3/2}}$$


2. Eddington Limit:

$$\dot{M}_{Edd} \sim 2 \times 10^{-9} \left(\frac{M}{M_{sun}} \right) M_{sun}/yr$$



$$f \sim \frac{L\sigma}{4\pi r^2 c} = f_g = \frac{GM}{r^2}$$

$$\rightarrow L \sim \frac{4\pi GMm_p c}{\sigma} = 1.3 \times 10^{38} \left(\frac{M}{M_{sun}} \right) \text{erg/s}$$



PI Colloquium, Feb. 15, 2012

How Bright can BHs Get?

1. Bondi Accretion:



$$\dot{M} \sim \rho \, 4\pi r_B^2 \, c_s \propto \frac{M^2}{T^{3/2}}$$

2. Eddington Limit:

$$\dot{M}_{Edd} \sim 2 \times 10^{-9} \left(\frac{M}{M_{sun}} \right) M_{sun}/yr$$

$$f \sim \frac{L\sigma}{4\pi r^2 c} = f_g = \frac{GM}{r^2}$$

$$\rightarrow L \sim \frac{4\pi GM m_p c}{\sigma} = 1.3 \times 10^{38} \left(\frac{M}{M_{sun}} \right) \text{erg/s}$$



How Bright can BHs Get?

1. Bondi Accretion:

$$\dot{M} \sim \rho \, 4\pi r_B^2 \, c_s \propto \frac{M^2}{T^{3/2}}$$

2. Eddington Limit:

$$\dot{M}_{Edd} \sim 2 \times 10^{-9} \left(\frac{M}{M_{sun}} \right) M_{sun}/yr$$

3. Angular Momentum:

$$\frac{GM}{c^2} \sim 10^{11} - 10^{15} \text{ cm} \ll R_{gas} \sim 1 \text{ pc} \sim 3 \times 10^{18} \text{ cm}$$

$$l = \sqrt{GM r} \rightarrow \frac{l_i}{l_f} \sim 10^2 - 10^4!$$

PI Colloquium, Feb. 15, 2012

Astrophysical “Viscosity”


- Molecular viscosity (Spitzer viscosity) much too small to drive observed luminosities
- Parameterized ignorance: $\nu = \alpha h c_s \rightarrow v^r \simeq \alpha v_k$



PI Colloquium, Feb. 15, 2012

Astrophysical “Viscosity”


- Molecular viscosity (Spitzer viscosity) much too small to drive observed luminosities
- Parameterized ignorance: $\nu = \alpha h c_s \rightarrow v^r \simeq \alpha v_k$
- What about magnetic fields ...



PI Colloquium, Feb. 15, 2012

Astrophysical “Viscosity”

- Molecular viscosity (Spitzer viscosity) much too small to drive observed luminosities
- Parameterized ignorance: $\nu = \alpha h c_s \rightarrow v^r \simeq \alpha v_k$
- What about magnetic fields ...



PI Colloquium, Feb. 15, 2012

Astrophysical “Viscosity”


- Molecular viscosity (Spitzer viscosity) much too small to drive observed luminosities
- Parameterized ignorance: $\nu = \alpha h c_s \rightarrow v^r \simeq \alpha v_k$
- What about magnetic fields ...



PI Colloquium, Feb. 15, 2012


Astrophysical “Viscosity”

- Molecular viscosity (Spitzer viscosity) much too small to drive observed luminosities
- Parameterized ignorance: $\nu = \alpha h c_s \rightarrow v^r \simeq \alpha v_k$
- What about magnetic fields ...




PI Colloquium, Feb. 15, 2012

Disks in the Other Dimension!




PI Colloquium, Feb. 15, 2012

Disks in the Other Dimension!




PI Colloquium, Feb. 15, 2012

Disks in the Other Dimension!



PI Colloquium, Feb. 15, 2012

Disks in the Other Dimension!



PI Colloquium, Feb. 15, 2012

Efficient Radiators ($T \ll T_{vir}$)

Shakura-Sunyaev Disks (α -disks)

$$h \ll r \quad v^\phi \simeq v_k, v^r \simeq \alpha v_k \ll v_k$$


$$L \sim 0.1 \dot{M} c^2 \quad (\text{independent of } \alpha!)$$

Efficient Radiators ($T \ll T_{vir}$)

Shakura-Sunyaev Disks (α -disks)

$$h \ll r \quad v^\phi \simeq v_k, v^r \simeq \alpha v_k \ll v_k$$



$$L \sim 0.1 \dot{M} c^2 \quad (\text{independent of } \alpha!)$$

Efficient Radiators ($T \ll T_{vir}$)


Shakura-Sunyaev Disks (α -disks)

$$h \ll r$$

$$v^\phi \simeq v_k, v^r \simeq \alpha v_k \ll v_k$$




$$L \sim 0.1 \dot{M} c^2 \quad (\text{independent of } \alpha!)$$



PI Colloquium, Feb. 15, 2012

Inefficient Radiators ($T \sim T_{vir}$)


Radiatively Inefficient Accretion Flows (RIAFs)



PI Colloquium, Feb. 15, 2012


Inefficient Radiators ($T \sim T_{vir}$)

Radiatively Inefficient
Accretion Flows (RIAFs)



PI Colloquium, Feb. 15, 2012

RIAFs at low $\dot{M}$: Two-Temp flows




→ Energy in ions → Energy in leptons → Radiation → Thin Disk

PI Colloquium, Feb. 15, 2012

RIAFs at low $\dot{M}$: Two-Temp flows

$$t_{ee} \propto \frac{1}{n}, \quad t_{pp} \propto \frac{1}{n} \left(\frac{m_p}{m_e} \right)^{1/2}, \quad t_{ep} \propto \frac{1}{n} \left(\frac{m_p}{m_e} \right)$$




PI Colloquium, Feb. 15, 2012

RIAFs at low $\dot{M}$: Two-Temp flows

$$t_{ee} \propto \frac{1}{n}, \quad t_{pp} \propto \frac{1}{n} \left(\frac{m_p}{m_e} \right)^{1/2}, \quad t_{ep} \propto \frac{1}{n} \left(\frac{m_p}{m_e} \right)$$

$$\frac{t_{ep}}{t_{ff}} \propto \frac{1/n}{r/v_k} \propto \frac{\dot{M}_{Edd}}{\dot{M}} f\left(\frac{r}{M}\right)$$



→ Energy in ions → Energy in leptons → Radiation → Thin Disk

PI Colloquium, Feb. 15, 2012

RIAFs at low $\dot{M}$: Two-Temp flows

$$t_{ee} \propto \frac{1}{n}, \quad t_{pp} \propto \frac{1}{n} \left(\frac{m_p}{m_e} \right)^{1/2}, \quad t_{ep} \propto \frac{1}{n} \left(\frac{m_p}{m_e} \right)$$

$$\frac{t_{ep}}{t_{ff}} \propto \frac{1/n}{r/v_k} \propto \frac{\dot{M}_{Edd}}{\dot{M}} f\left(\frac{r}{M}\right)$$




$$\dot{M} < 10^{-2} \dot{M}_{Edd}$$

Today 90%-99% of all
BHs have $\dot{M} \ll \dot{M}_{Edd}$
→ Most BHs are RIAFS!

Energy in ions →  Energy in leptons → Radiation → Thin Disk



PI Colloquium, Feb. 15, 2012

Sagittarius A*: Neighborhood RIAF





PI Colloquium, Feb. 15, 2012

Sagittarius A*: Neighborhood RIAF




PI Colloquium, Feb. 15, 2012

Sagittarius A*: Neighborhood RIAF




PI Colloquium, Feb. 15, 2012

Sgr A*'s disappearing act!



PI Colloquium, Feb. 15, 2012


Mass and Just How Dim?




- $M = 4 \times 10^6 M_{\text{sun}}$

PI Colloquium, Feb. 15, 2012

Mass and Just How Dim?




- $M = 4 \times 10^6 M_{\text{sun}}$



PI Colloquium, Feb. 15, 2012


Mass and Just How Dim?



- $M = 4 \times 10^6 M_{\text{sun}}$

PI Colloquium, Feb. 15, 2012


Mass and Just How Dim?



- $M = 4 \times 10^6 M_{\text{sun}}$

PI Colloquium, Feb. 15, 2012


Mass and Just How Dim?



- $M = 4 \times 10^6 M_{\text{sun}}$

PI Colloquium, Feb. 15, 2012


Mass and Just How Dim?



- $M = 4 \times 10^6 M_{\text{sun}}$
- $L_{\text{Edd}} \simeq 10^{45} \text{ erg/s}$
- $L \sim 10^{-9} L_{\text{Edd}}!$

PI Colloquium, Feb. 15, 2012

Mass and Just How Dim?



- $M = 4 \times 10^6 M_{\text{sun}}$
- $L_{\text{Edd}} \simeq 10^{45} \text{ erg/s}$
- $L \sim 10^{-9} L_{\text{Edd}}!$
- $\dot{M} \sim 10^{-9} \dot{M}_{\text{Edd}}$
(if at 10% efficiency!)

PI Colloquium, Feb. 15, 2012

Is $\dot{M}$ really that low?

$$\dot{M}_{Bondi} \sim 10^{-5} M_{sun}/yr$$

$$\sim 10^{-4} \dot{M}_{Edd}$$

$$\gg 10^{-9} \dot{M}_{Edd}!$$

X-ray

$T \sim \text{keV}$



PI Colloquium, Feb. 15, 2012

Is $\dot{M}$ really that low?

$$\dot{M}_{\text{Bondi}} \sim 10^{-5} M_{\text{sun}}/\text{yr}$$

$$\sim 10^{-4} \dot{M}_{\text{Edd}}$$

$$\gg 10^{-9} \dot{M}_{\text{Edd}}!$$

X-ray

$T \sim \text{keV}$



PI Colloquium, Feb. 15, 2012

Is $\dot{M}$ really that low?

$$\dot{M}_{Bondi} \sim 10^{-5} M_{sun}/yr$$

$$\sim 10^{-4} \dot{M}_{Edd}$$

$$\gg 10^{-9} \dot{M}_{Edd}!$$

→ Sgr A* is a RIAF!
(like 99%!)

X-ray

$T \sim \text{keV}$



PI Colloquium, Feb. 15, 2012

Is $\dot{M}$ really that low?

$$\dot{M}_{Bondi} \sim 10^{-5} M_{sun}/yr$$

$$\sim 10^{-4} \dot{M}_{Edd}$$

$$\gg 10^{-9} \dot{M}_{Edd}!$$

→ Sgr A* is a RIAF!
(like 99%!)

Occupy Galactic Center?

X-ray


$T \sim \text{keV}$



PI Colloquium, Feb. 15, 2012

Loose Threads ...

- Polarization:




$$RM \equiv \frac{d\chi}{d\lambda^2} \sim 10^6 \text{ rad/m}^2 \propto nBR$$

$$\rightarrow \dot{M} \sim 10^{-8} M_{\text{sun}}/\text{yr}$$

PI Colloquium, Feb. 15, 2012

Loose Threads ...

- Polarization:




$$RM \equiv \frac{d\chi}{d\lambda^2} \sim 10^6 \text{ rad/m}^2 \propto nBR$$

$$\rightarrow \dot{M} \sim 10^{-8} M_{\text{sun}}/\text{yr}$$

PI Colloquium, Feb. 15, 2012

Loose Threads ...

- Polarization: $\rightarrow \dot{M} \sim 10^{-8} M_{\text{sun}}/\text{yr}$
- Accretion vs. Jet Power




Craig Walker

PI Colloquium, Feb. 15, 2012

Loose Threads ...

- Polarization: $\rightarrow \dot{M} \sim 10^{-8} M_{\text{sun}}/\text{yr}$
- Accretion vs. Jet Power
- Accretion power vs. something else?




- Power Density
- Inevitability

PI Colloquium, Feb. 15, 2012

Loose Threads ...

- Polarization: $\rightarrow \dot{M} \sim 10^{-8} M_{\text{sun}}/\text{yr}$
- Accretion vs. Jet Power
- Accretion power vs. something else?



"That some form of the meteoric theory is certainly the true and complete explanation of solar heat can scarcely be doubted, when the following reasons are considered: (1) No other natural explanation, except by chemical action, can be conceived. (2) The chemical theory is quite insufficient, because the most energetic chemical action we know, taking place between substances amounting to the whole sun's mass, would only generate about 3,000 years' heat. (3) There is no difficulty in accounting for 20,000,000 years' heat by the meteoric theory."

Lord Kelvin 1862

- Power Density
- Inevitability

PI Colloquium, Feb. 15, 2012

Loose Threads ...

- Polarization: $\rightarrow \dot{M} \sim 10^{-8} M_{\text{sun}}/\text{yr}$
- Accretion vs. Jet Power
- Accretion power vs. something else?
- Do RIAFs make sense?
- How does ang. mom. transfer really work?



PI Colloquium, Feb. 15, 2012

Loose Threads ...



- Polarization: $\rightarrow \dot{M} \sim 10^{-8} M_{\text{sun}}/\text{yr}$
- Accretion vs. Jet Power
- Accretion power vs. something else?
- Do RIAFs make sense?
- How does ang. mom. transfer really work?
- What are the consequences of strong gravity?

PI Colloquium, Feb. 15, 2012


Advantages of being next door!




Advantages of being next door!




Advantages of being next door!



Accretion Flows Near the Horizon



Accretion Flows Near the Horizon




Diffraction Limited Telescopes



PI Colloquium, Feb. 15, 2012

Diffraction Limited Telescopes




PI Colloquium, Feb. 15, 2012


Diffraction Limited Telescopes



PI Coll



Diffraction telescopes




Radio Interferometry

$$E_1 = E e^{i\omega t}$$

$$E_2 = E e^{i\omega t - i\phi}$$


$$\phi = \omega\tau \simeq 2\pi \frac{u\alpha}{\lambda}$$

$$\langle E_1 E_2^* \rangle = |E|^2 e^{2\pi i u \alpha / \lambda}$$



Smithsonian Astrophysical Observatory, 3/8/10

Radiometry



$$E_1 = E e^{i\omega t}$$

$$E_2 = E e^{i\omega t - i\phi}$$


$$\phi = \omega \tau \simeq 2\pi \frac{u\alpha}{\lambda}$$

$$\langle E_1 E_2^* \rangle = |E|^2 e^{2\pi i u \alpha / \lambda}$$



Smithsonian Astrophysical Observatory, 3/8/10

Radiometry



$$E_1 = E e^{i\omega t}$$

$$E_2 = E e^{i\omega t - i\phi}$$

$$\phi = \omega \tau \simeq 2\pi \frac{u\alpha}{\lambda}$$

$$\langle E_1 E_2^* \rangle = |E|^2 e^{2\pi i u \alpha / \lambda}$$

$$\begin{aligned} \rightarrow V_{12}(u) &= \langle E_1 E_2^* \rangle \\ &= \int I(\alpha) e^{2\pi i u \alpha / \lambda} d\alpha \end{aligned}$$


$$\Delta\theta \simeq \lambda/u$$



Smithsonian Astrophysical Observatory, 3/8/10

Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$




PI Colloquium, Feb. 15, 2012

Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$
- The bad:



$$V_{ij} = \int I(\alpha) e^{2\pi i \mathbf{u}_{ij} \cdot \alpha / \lambda} d^2\alpha$$



Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$
- The bad:

$$V_{ij} = \int I(\alpha) e^{2\pi i \mathbf{u}_{ij} \cdot \alpha / \lambda} d^2 \alpha$$





PI Colloquium, Feb. 15, 2012

Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$
- The bad:

$$V_{ij} = \int I(\alpha) e^{2\pi i \mathbf{u}_{ij} \cdot \alpha / \lambda} d^2\alpha$$





PI Colloquium, Feb. 15, 2012

Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$
- The bad:

$$V_{ij} = \int I(\alpha) e^{2\pi i \mathbf{u}_{ij} \cdot \alpha / \lambda} d^2 \alpha$$





PI Colloquium, Feb. 15, 2012

Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$
- The bad:

$$V_{ij} = \int I(\alpha) e^{2\pi i \mathbf{u}_{ij} \cdot \alpha / \lambda} d^2 \alpha$$



PI Colloquium, Feb. 15, 2012

Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$
- The bad: $V_{ij} = \int I(\alpha) e^{2\pi i \mathbf{u}_{ij} \cdot \alpha / \lambda} d^2\alpha$
- The ugly:
 - Sparse u-v coverage
 - Atmospheric phase errors




Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$
- The bad: $V_{ij} = \int I(\alpha) e^{2\pi i \mathbf{u}_{ij} \cdot \alpha / \lambda} d^2\alpha$
- The ugly:
 - Sparse u-v coverage
 - Atmospheric phase errors
- Why the VLBA will not work:
 - 100-300 μas





Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$
- The bad: $V_{ij} = \int I(\alpha) e^{2\pi i u \alpha} d\alpha$
- The ugly:
 - Sparse u-v coverage
 - Atmospheric phase errors
- Why the VLBA will not work:
 - 100-300 μas
 - Optically thick—thin transition



Very Long Baseline Interferometry



- The good: $\Delta\theta = \lambda/u$
- The bad: $V_{ij} = \int I(\alpha) e^{2\pi i u \alpha} d\alpha$
- The ugly:
 - Sparse u-v coverage
 - Atmospheric phase errors
- Why the VLBA will not work
 - 100-300 μas
 - Optically thick—thin transition



PI Colloquium, Feb. 15, 2012

Very Long Baseline Interferometry



- The good: $\Delta\theta = \lambda/u$
- The bad: $V_{ij} = \int I(\alpha) e^{2\pi i u \alpha} d\alpha$
- The ugly:
 - Sparse u-v coverage
 - Atmospheric phase errors
- Why the VLBA will
 - 100-300 μas
 - Optically thick—thin



Pr Colloquium, Feb. 13, 2012

Very Long Baseline Interferometry



- The good: $\Delta\theta = \lambda/u$
- The bad: $V_{ij} = \int I(\alpha) e^{2\pi i u \alpha} d\alpha$
- The ugly:
 - Sparse u-v coverage
 - Atmospheric phase errors
- Why the VLBA will not work:
 - 100-300 μas
 - Optically thick—thin transition
 - Interstellar electron scattering



PI Colloquium, Feb. 15, 2012


Very Long Baseline Interferometry

- The good: $\Delta\theta = \lambda/u$
- The bad: $V_{ij} = \int I(\alpha) e^{2\pi i u \alpha} d\alpha$
- The ugly:
 - Sparse u-v coverage
 - Atmospheric phase errors
- Why the VLBA will not work:
 - 100-300 μas
 - Optically thick—thin transition
 - Interstellar electron scattering
- But at 1 mm ...



PI Colloquium, Feb. 15, 2012


The Event Horizon Telescope



- Earth-sized mm/sub-mm VLBI array
- Existing and **Planned** telescopes.
- Accesses resolutions of $\sim 10 \mu\text{as}$,
→ Sufficient to resolve Sgr A*'s horizon!
- Why mm/sub-mm?
 - Optically thin transition ($< 3\text{mm}$)
 - Interstellar scattering ($< 3\text{mm}$)
 - Resolution from ground ($< 3\text{mm}$)
 - “Standard” VLBI techniques ($> 0.1\text{mm}$)
 - Good seeing at many sites ($> 0.5\text{mm}$)




Event Horizon Telescope




- Earth-sized mm/sub-mm VLBI array
- Existing and **Planned** telescopes.
- Accesses resolutions of $\sim 10 \mu\text{as}$,
→ Sufficient to resolve Sgr A*'s horizon!
- Why mm/sub-mm?
 - Optically thin transition ($< 3\text{mm}$)
 - Interstellar scattering ($< 3\text{mm}$)
 - Resolution from ground ($< 3\text{mm}$)
 - “Standard” VLBI techniques ($> 0.1\text{mm}$)
 - Good seeing at many sites ($> 0.5\text{mm}$)



First Data from the EHT



- Data already exists on Hawaii-CARMA-SMT
 - Doeleman et al., Nature, 455, 78 (2008)
 - Fish et al., ApJ, 727, 36 (2011)




PI Colloquium, Feb. 15, 2012

First Data from the EHT




- Data already exists on Hawaii-CARMA-SMT
 - Doeleman et al., Nature, 455, 78 (2008)
 - Fish et al., ApJ, 727, 36 (2011)



PI Colloquium, Feb. 15, 2012

Future Growth of the EHT




- Increasing Sensitivity:

- Phasing Arrays
 - Hawaii (Done!)
 - CARMA (In progress!)
 - ALMA (Approved!)
- Expanding Bandwidth to 4GHz
- Building New Receivers

PI Colloquium, Feb. 15, 2012

Future Growth of the EHT



- Increasing Sensitivity:


- Phasing Arrays
 - Hawaii (Done!)
 - CARMA (In progress!)
 - ALMA (Approved!)
- Expanding Bandwidth to 4GHz
- Building New Receivers

- Adding Sites:

- Planned
 - ALMA, APEX
 - LMT
 - Plateau de Bure
 - Pico Veleta
 - SPT
 - Haystack

PI Colloquium, Feb. 15, 2012

Future Growth of the EHT



- Increasing Sensitivity:


- Phasing Arrays
 - Hawaii (Done!)
 - CARMA (In progress!)
 - ALMA (Approved!)
- Expanding Bandwidth to 4GHz
- Building New Receivers

- Adding Sites:

- Planned
 - ALMA, APEX
 - LMT
 - Plateau de Bure
 - Pico Veleta
 - SPT
 - Haystack
- New (ALMA prototype dishes)
 - New Zealand
 - Kilimanjaro
 - South Africa
 - SPT

PI Colloquium, Feb. 15, 2012

Future Growth of the EHT



- Incr

—

- Adc


—

—

—


PI Colloquium, Feb. 15, 2012

EHT Data Today








PI Colloquium, Feb. 15, 2012

EHT Data Today




PI Colloquium, Feb. 15, 2012

EHT Data Today



PI Colloquium, Feb. 15, 2012

EHT Data Today





PI Colloquium, Feb. 15, 2012

Filling gaps in data with theory

Model Input:

- Phenomenological (gaussian, annuli, etc.)
- Physical (α -disk, RIAF, jet, etc.)



PI Colloquium, Feb. 15, 2012

Measuring Fits and Models

- Quality of a fit:
 - Do good (high likelihood) fits exist? χ^2
 - What model parameters are preferred? $P(a|V)$
 - Are these consistent among different epochs? $P(a|V)$
- Quality of a model:
 - Bayesian Information Criterion: $BIC = \chi^2 + k \log N$
 - Akaike Information Criterion: $AIC = \chi^2 + 2k + \frac{2k(k+1)}{N-k-1}$
 - Relative significance: $w_{ij} = e^{-(IC_i - IC_j)/2}$
 - Occam's Razor?
 - Lies, damn lies, and statistics!

ΔIC	Conclusion
0	Negative
0-1	Not worth more than a bare mention
1-3	Positive
3-5	Strong
5-10	Very Strong
>10	Conclusive

PI Colloquium, Feb. 15, 2012



Measuring Fits and Models

- Quality of a fit:
 - Do good (high likelihood) fits exist? χ^2
 - What model parameters are preferred? $P(a|V)$
 - Are these consistent among different epochs? $P(a|V)$
- Quality of a model:
 - Bayesian Information Criterion: $BIC = \chi^2 + k \log N$
 - Akaike Information Criterion: $AIC = \chi^2 + 2k + \frac{2k(k+1)}{N-k-1}$
 - Relative significance: $w_{ij} = e^{-(IC_i - IC_j)/2}$
 - Occam's Razor?
 - Lies, damn lies, and statistics!

ΔIC	Conclusion
0	Negative
0-1	Not worth more than a bare mention
1-3	Positive
3-5	Strong
5-10	Very Strong
>10	Conclusive

PI Colloquium, Feb. 15, 2012

Toy Model: Symmetric Gaussian



$$\sigma = 15.8 \pm 0.2 \text{ } \mu\text{s}$$

$$\chi^2/\text{DoF} = 1.16$$



$$\text{BIC} = 98.1$$

$$\text{AIC} = 87.7$$

AEB, Fish, Doeleman & Loeb, ApJ, 735, 110 (2011)

PI Colloquium, Feb. 15, 2012

Toy Model: Symmetric Gaussian



$$\sigma = 15.8 \pm 0.2 \mu\text{as}$$

$$\chi^2/\text{DoF} = 1.16$$



$$\text{BIC} = 98.1$$

$$\text{AIC} = 87.7$$

AEB, Fish, Doeleman & Loeb, ApJ, 735, 110 (2011)

PI Colloquium, Feb. 15, 2012

Toy Model: Asymmetric Gaussian



$$\sigma = \left[\frac{1}{2\sigma_m^2} + \frac{1}{2\sigma_M^2} \right]^{-1/2} = 20.5^{+0.3+0.5}_{-0.8-1.3} \mu\text{as}$$

$$A = \left[\frac{\sigma_M^2 - \sigma_m^2}{\sigma_M^2 + \sigma_m^2} \right]^{1/2} = 0.70^{+0.03+0.05}_{-0.1-0.18}$$

$$\xi = 19^{+3^\circ+6^\circ}_{-1^\circ-2^\circ}$$

$$\chi^2/\text{DoF} = 0.961$$

$$\text{BIC} = 91.3 \quad (-6.9)$$


$$\text{AIC} = 77.3 \quad (-7.4)$$

AEB, Fish, Doeleman & Loeb, ApJ, 735, 110 (2011)

PI Colloquium, Feb. 15, 2012

A Physical Model: RIAFs


- Thick, self-similar disk
- Keplerian beyond ISCO, ballistic inside
- Sub-equipartition magnetic field
- “Thermal” and power-law electron components
- Self-Absorbed Synchrotron
- Model defined by n_{th} , n_{nth} , T_e , and a



PI Colloquium, Feb. 15, 2012

A Physical Model: RIAFs



- Thick, self-similar disk
- Keplerian beyond ISCO, ballistic inside
- Sub-equipartition magnetic field
- “Thermal” and power-law electron components
- Self-Absorbed Synchrotron
- Model defined by n_{th} , n_{nth} , T_e , and a



PI Colloquium, Feb. 15, 2012

A Physical Model: RIAFs


- Thick, self-similar disk
- Keplerian beyond ISCO, ballistic inside
- Sub-equipartition magnetic field
- “Thermal” and power-law electron components
- Self-Absorbed Synchrotron
- Model defined by n_{th} , n_{nth} , T_e , and a
- Leverage existing spectra




PI Colloquium, Feb. 15, 2012

A Physical Model: RIAFs

Low Spin




High Spin





electron components

- Self-Absorbed Synchrotron
- Model defined by n_{th} , n_{nth} , T_e , and a
- Leverage existing spectra



PI Colloquium, Feb. 15, 2012

Radiatively inefficient accretion flow



$$\chi^2/\text{DoF}=0.830$$



$$\text{BIC}=82.9 \quad (-8.4, -15.3)$$

$$\text{AIC}=68.9 \quad (-8.4, -15.8)$$

AEB, Fish, Doeleman & Loeb, ApJ, 735, 110 (2011)

PI Colloquium, Feb. 15, 2012

Radiatively inefficient accretion flow



$$\chi^2/\text{DoF}=0.830$$


$$\text{BIC}=82.9 \quad (-8.4, -15.3)$$

$$\text{AIC}=68.9 \quad (-8.4, -15.8)$$

AEB, Fish, Doeleman & Loeb, ApJ, 735, 110 (2011)

PI Colloquium, Feb. 15, 2012

Radiatively inefficient accretion flow



AEB, Fish, Doeleman & Loeb, ApJ, 735, 110 (2011)


PI Colloquium, Feb. 15, 2012

The Spin of Sgr A*

$$a \sim 0^{+0.64+0.86}$$

$$\theta = 68^{+5+8}_{-20-28} \text{ deg.}$$

$$\xi = -52^{+17+33}_{-15-24} \text{ deg.}$$




The Spin of Sgr A*

$$a \sim 0^{+0.64+0.86}$$

$$\theta = 68^{+5+8}_{-20-28} \text{ deg.}$$

$$\xi = -52^{+17+33}_{-15-24} \text{ deg.}$$




Spin Magnitude

PI Colloquium, Feb. 15, 2012

The Importance of Phase

- Lack of phase info gives $\xi \rightarrow \xi + 180^\circ$ degeneracy



PI Colloquium, Feb. 15, 2012

The Importance of Phase

- Lack of phase info gives $\xi \rightarrow \xi + 180^\circ$ degeneracy
- Image structure strongly dependent upon phase

PI Colloquium, Feb. 15, 2012

The Importance of Phase

- Lack of phase info gives $\xi \rightarrow \xi + 180^\circ$ degeneracy
- Image structure strongly dependent upon phase



$|V|$



$\arg(V)$



PI Colloquium, Feb. 15, 2012

The Importance of Phase


- Lack of phase info gives $\xi \rightarrow \xi + 180^\circ$ degeneracy
- Image structure strongly dependent upon phase



$|V|$



$\arg(V)$



PI Colloquium, Feb. 15, 2012

The Importance of Phase


- Lack of phase info gives $\xi \rightarrow \xi + 180^\circ$ degeneracy
- Image structure strongly dependent upon phase



$|V|$




$\arg(V)$



PI Colloquium, Feb. 15, 2012

The Future with Closure Phases

$|V|$ Only




AEB, Fish, Doeleman & Loeb, ApJ, 738, 38 (2011)

PI Colloquium, Feb. 15, 2012

The Future with Closure Phases

$|V|$ & Anticipated Precision on all
triangles in an SMT-Hawaii-CARMA-APEX




AEB, Fish, Doeleman & Loeb, ApJ, 738, 38 (2011)

PI Colloquium, Feb. 15, 2012

The Future with Closure Phases

$|V|$ & Anticipated Precision on all triangles in an SMT-Hawaii-CARMA-APEX




AEB, Fish, Doeleman & Loeb, ApJ, 738, 38 (2011)

PI Colloquium, Feb. 15, 2012

The Future with Closure Phases

|V| & Anticipated Precision on all triangles
in an SMT-Hawaii-CARMA-ALMA10 array




AEB, Fish, Doeleman & Loeb, ApJ, 738, 38 (2011)

PI Colloquium, Feb. 15, 2012

The Future with Closure Phases

|V| & Anticipated Precision on all triangles
in an SMT-Hawaii-CARMA-ALMA10-LMT




AEB, Fish, Doeleman & Loeb, ApJ, 738, 38 (2011)

PI Colloquium, Feb. 15, 2012

The Future with Closure Phases

|V| & Anticipated Precision on all triangles
in an SMT-Hawaii-CARMA-ALMA10-LMT




AEB, Fish, Doeleman & Loeb, ApJ, 738, 38 (2011)

PI Colloquium, Feb. 15, 2012

The Future with Closure Phases


|V| & Anticipated Precision on all triangles
in an SMT-Hawaii-CARMA-ALMA10-LMT



AEB, Fish, Doeleman & Loeb, ApJ, 738, 38 (2011)


PI Colloquium, Feb. 15, 2012

The Future: Is Sgr A* a RIAF?



→ How inefficient is ion-electron coupling really?
(Do RIAFs make sense?)


→ Angular momentum transport near horizon.





Bittner, AEB & Loeb (2012)

PI Colloquium, Feb. 15, 2012

The Future: Does the MRI work?



Dexter, Agol & Fragile (2009)





Shcherbakov, Penna & McKinney (2011)

- Only fit smooth models thus far (though see Dexter, Agol & Fragile (2009),)
- Power from turbulence on small scales (large u)
- Measure turbulent spectrum:

$$\varepsilon_k \propto k^{-\gamma} \text{ and } j_\nu \propto \nu^{-\alpha} \rightarrow V(u) \propto u^{-\eta}$$

PI Colloquium, Feb. 15, 2012

The Future: Does the MRI work?




- Only fit smooth models that Agol & Fragile (2009),)
- Power from turbulence on small scales (large u)
- Measure turbulent spectrum:

$$\varepsilon_k \propto k^{-\gamma} \text{ and } j_\nu \propto \nu^{-\alpha} \rightarrow V(u) \propto u^{-\eta}$$

PI Colloquium, Feb. 15, 2012

The Future: Is Sgr A* Static?




- Probed timescales from 10s – 2 yrs already
- Thick disks don't align
- Should precess due to:
 - Lense-Thirring
 - Local massive perturbers
 - Deviations from GR?

- Timescale

$$\sim \frac{L}{\tau} \sim a^{-1} \left(\frac{R}{10^3 GM/c^2} \right)^2 \text{ yrs}$$

PI Colloquium, Feb. 15, 2012

The Future: Is Sgr A* Static?




- Probed timescales from 10s – 2 yrs already
- Thick disks don't align
- Should precess due to:
 - Lense-Thirring
 - Local massive perturbers
 - Deviations from GR?

- Timescale

$$\sim \frac{L}{\tau} \sim a^{-1} \left(\frac{R}{10^3 GM/c^2} \right)^2 \text{ yrs}$$

PI Colloquium, Feb. 15, 2012

The Future: Is Sgr A* Static?




- Probed timescales from 10s – 2 yrs already
- Thick disks don't align
- Should precess due to:
 - Lense-Thirring
 - Local massive perturbers
 - Deviations from GR?
- Timescale

$$\sim \frac{L}{\tau} \sim a^{-1} \left(\frac{R}{10^3 GM/c^2} \right)^2 \text{ yrs}$$



PI Colloquium, Feb. 15, 2012

The future: Supply & Demand




PI Colloquium, Feb. 15, 2012

The future: Supply & Demand




PI Colloquium, Feb. 15, 2012

The future: Supply & Demand



PI Colloquium, Feb. 15, 2012


The future: Supply & Demand



→ Increase in $\dot{M}$ by roughly 2 or so.

PI Colloquium, Feb. 15, 2012


Testing GR with quasi-Kerr



- $g_{\mu\nu} = g_{\mu\nu}^{Kerr} + \varepsilon h_{\mu\nu}$
 $\rightarrow M, a, q = -(a^2 + \varepsilon)$
- Wrong quadrupole (measured by ε)
- θ and ξ surprisingly insensitive to ε
 $\rightarrow$ robust direction
- Already best fit model clearly has nonzero ε !

Johannsen, AEB, Psaltis & Loeb (2011) Colloquium, Feb. 15, 2012

Testing GR with quasi-Kerr




- $g_{\mu\nu} = g_{\mu\nu}^{Kerr} + \varepsilon h_{\mu\nu}$
 $\rightarrow M, a, q = -(a^2 + \varepsilon)$
- Wrong quadrupole (measured by ε)
- θ and ξ surprisingly insensitive to ε
 $\rightarrow$ robust direction
- Already best fit model clearly has nonzero ε !

Johannsen, AEB, Psaltis & Loeb (2011) Colloquium, Feb. 15, 2012

Non-Imaging Analysis: The Near Future!

Testing GR with quasi-Kerr




- Bandwidth: 1GHz
- Observatories
 - SMT
 - JCMT, SMA (LO)
 - CARMA (2 ind. dishes)
- $|V|$ only

Johannsen, AEB, Psaltis & Loeb (2011) PI Colloquium, Feb. 15, 2012

Non-Imaging Analysis: The Near Future!

Testing GR with quasi-Kerr




- Bandwidth: 4GHz
- Observatories
 - SMT
 - Phased Hawaii
 - Phased CARMA (8)
 - Phased ALMA (10)
- $|V|$ only

Johannsen, AEB, Psaltis & Loeb (2011) PI Colloquium, Feb. 15, 2012

Non-Imaging Analysis: The Near Future!

Testing GR with quasi-Kerr



- Bandwidth: 4GHz
- Observatories
 - SMT
 - Phased Hawaii
 - Phased CARMA (8)
 - Phased ALMA (10)
- $|V|$
- *+ Closure Phases*

Johannsen, AEB, Psaltis & Loeb (2011) PI Colloquium, Feb. 15, 2012

Conclusions (so far...)

- Most black holes are “boring”, i.e., $\dot{M} \ll \dot{M}_{Edd}$
→ RIAFs are common.
- Lot's of reasons to believe Sgr A* is a RIAF
(i.e., Copernicus strikes again!).
- Low $\dot{M}$ + proximity enables unique horizon resolving study of Sgr A* with the Event Horizon Telescope.
- Already finding that RIAFs work extremely well.
- In the near future we will begin to test:
 - Internal consistency of RIAFs specifically
 - Critical components of accretion models generally
 - Structure of the spacetime around Sgr A*

PI Colloquium, Feb. 15, 2012

Conclusions (so far...)

- Most black holes are “boring”, i.e., $\dot{M} \ll \dot{M}_{Edd}$
→ RIAFs are common.
- Lot's of reasons to believe Sgr A* is a RIAF
(i.e., Copernicus strikes again!).
- Low $\dot{M}$ + proximity enables unique horizon resolving study of Sgr A* with the **EventHorizonTelescope**
- Already finding that RIAFs work extremely well.
- In the near future we will begin to test:
 - Internal consistency of RIAFs specifically
 - Critical components of accretion models generally
 - Structure of the spacetime around Sgr A*

PI Colloquium, Feb. 15, 2012

Conclusions (so far...)

- Most black holes are “boring”, i.e., $\dot{M} \ll \dot{M}_{Edd}$
→ RIAFs are common.
- Lot's of reasons to believe Sgr A* is a RIAF
(i.e., Copernicus strikes again!).
- Low $\dot{M}$ + proximity enables unique horizon resolving study of Sgr A* with the **EventHorizonTelescope**
- Already finding that RIAFs work extremely well.
- In the near future we will begin to test:
 - Internal consistency of RIAFs specifically
 - Critical components of accretion models generally
 - Structure of the spacetime around Sgr A*

PI Colloquium, Feb. 15, 2012