

Title: Explorations in Numerical Relativity - Lecture 1

Date: Apr 02, 2012 11:30 AM

URL: <http://pirsa.org/12040038>

Abstract:

Applications Places System **Ubuntu_11.04 - VMware Pla...** File Virtual Machine Help 11:35 AM ubuntu-user

Explorations in Numerical Relativity | PSI Student Portal - Mozilla Firefox

File Edit View History Bookmarks Tools Help

Explorations in Numerical Rel...

perimeterinstitute.ca https://perimeterinstitute.ca/psi_portal/wiki/explorations-num Google

Explorations in Numerical Relativity

Home > Groups > PSI 2012 Group

View What links here Edit Revisions

Lecturer: [[llehner@perimeterinstitute.ca|Luis Lehner] and [[fpretori@princeton.edu|Frans Pretorius]

Luis Lehner is a Professor at the University of Guelph and Perimeter Institute Associate.

Frans Pretorius is a Professor at the University of Princeton. He is a DRC of Perimeter Institute.

Both have made pioneering contributions to using advanced numerical methods to extract definite predictions for the behavior of black holes and other strongly gravitating systems from Einstein's equations.

Important Instructions to Install NR Software

Please follow the instructions in softconfig.pdf attached

to install missing/required software in your linux distribution (open vmware, ubuntu for this)

[Software install description](#) (txt version of software installation description)

[Font file for font required when installing software](#) (unzip it, and rename it Helvetica.ttf)

Note: After you are done, in a new terminal if you try to run: xvs or DV (i.e. type xvs and press enter), you should see a menu coming up.

Don't worry we'll learn to use them in the course.

<http://www.perimeterscholars.org/> count configuration so that the software can be accessed easily, follow the

Wiki page Create

Latest wiki posts

PI Explorations in Numerical Relativity
Posted: 20 Mar 2012 - 13:31 by [PerimeterIT](#)
0 comment(s)

 [Explorations in Quantum Information](#)
new
Posted: 9 Mar 2012 - 12:12 by [scroke](#)
0 comment(s)

1 of 8 »

Latest wiki comments

 [why there is no lecture note](#)
Posted: 3 Feb 2012 - 14:48 by [dkubiznak](#)

Explorations in ... [Update Manager] ubuntu-user@pi-... Downloads Lec1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:37 AM ubuntu-user

PSI: Explorations in Gravitational Physics: Numerical Relativity (Lehner, Noble & Choptuik) - Mozilla Firefox

File Edit View History Bookmarks Tools Help

PSI: Explorations in Gravitatio... +

http://bh0.physics.ubc.ca/2011-pi-nr/www/ Google

Perimeter Scholars International Explorations in Gravitational Physics Numerical Relativity

L. Lehner and S. Noble (as well as MW Choptuik & F Pretorius)

Please report all errors/typos. etc to Luis Lehner at [llehner "at" perimeterinstitute.ca](mailto:llehner@perimeterinstitute.ca)

Last updated April 3rd, 2011

Quick Links: [[Project 1](#) : [Project 2](#) : [Project 3](#)]

Lecture Notes

Week 1: Solution of Classical Field Equations Using Finite Difference Techniques (L. Lehner, MW Choptuik)

1. Solving the wave equation using finite difference techniques [[PDF](#)]
2. The 3+1 approach to the Einstein equations [[PDF](#)]
3. Time dependent spherically symmetric (SS) spacetimes and the SS Einstein-massless-Klein-Gordon system [[PDF](#)]
4. Project 1 handout [[PDF](#)]

Week 2: Much Ado About Hydrodynamics and Related Problems (S. Noble, L. Lehner)

PSI: Explorations... [Update Manager] ubuntu-user@pi-... Downloads Lec1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:37 AM ubuntu-user

PSI: Explorations in Gravitational Physics: Numerical Relativity (Lehner, Noble & Choptuik) - Mozilla Firefox

File Edit View History Bookmarks Tools Help

PSI: Explorations in Gravitatio...

http://bh0.physics.ubc.ca/2011-pi-nr/www/ Google

Please report ali errors/typos. etc to Luis Lehner at [llehner "at" perimeterinstitute.ca](mailto:llehner@perimeterinstitute.ca)

Last updated April 3rd, 2011

Quick Links: [[Project 1](#) : [Project 2](#) : [Project 3](#)]

Lecture Notes

Week 1: Solution of Classical Field Equations Using Finite Difference Techniques (L.Lehner, MW Choptuik)

1. Solving the wave equation using finite difference techniques [[PDF](#)]
2. The 3+1 approach to the Einstein equations [[PDF](#)]
3. Time dependent spherically symmetric (SS) spacetimes and the SS Einstein-massless-Klein-Gordon system [[PDF](#)]
4. Project 1 handout [[PDF](#)]

Week 2: Much Ado About Hydrodynamics and Related Problems (S. Noble, L Lehner)

1. Overview and Burger's equation (includes Project 2) [[PDF](#)]
2. General Relativistic Hydrodynamics and beyond [[PDF](#)]

Week 3: Miscellaneous topics (L Lehner, F Pretorius)

1. Lecture 1: Overview [[PDF](#)]
2. Lecture 4: AMR [[PDF](#)]

[Background/Reference Material and Related Software](#)

PSI: Explorations... [Update Manager] ubuntu-user@pi-... Downloads Lec1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:40 AM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 1 (2 of 83) Fit Page Width

Thumbnails



0



1



2



3

Course Outline

- **Solution of Classical Field Equations Using Finite Difference Techniques (Luis,Matt)**
 1. Solving the wave equation using finite difference techniques
 2. $3 + 1$ approach to the Einstein equations
 3. Dynamical spherically symmetric spacetimes
 4. Spherically symmetric Einstein-Klein-Gordon Evolution
 5. Introduction to Black Hole Critical Phenomena
- **General Relativistic Hydrodynamics Using Godunov/HRSC Schemes (Scott,Luis)**
 1. Mathematical structure; Linearly degenerate vs truly nonlinear eqns
 2. Burgers eqn; Godunov Methods & the Riemann problem
 3. $3 + 1$ Approach to GRHydrodynamics
 4. Stationary solutions, TOV stars & perturbations
 5. Magnetohydrodynamics & miscellaneous topics
- **Topics in Numerical Relativity (Luis,Frans)**
 1. Gravitational waves overview (nature in GR & sources)
 2. Newman Penrose formalism, Teukolsky equation

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:40 AM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 1 (2 of 83) Fit Page Width

Thumbnails



0



1



2



3

Course Outline

- **Solution of Classical Field Equations Using Finite Difference Techniques (Luis,Matt)**
 1. Solving the wave equation using finite difference techniques
 2. $3 + 1$ approach to the Einstein equations
 3. Dynamical spherically symmetric spacetimes
 4. Spherically symmetric Einstein-Klein-Gordon Evolution
 5. Introduction to Black Hole Critical Phenomena
- **General Relativistic Hydrodynamics Using Godunov/HRSC Schemes (Scott,Luis)**
 1. Mathematical structure; Linearly degenerate vs truly nonlinear eqns
 2. Burgers eqn; Godunov Methods & the Riemann problem
 3. $3 + 1$ Approach to GRHydrodynamics
 4. Stationary solutions, TOV stars & perturbations
 5. Magnetohydrodynamics & miscellaneous topics
- **Topics in Numerical Relativity (Luis,Frans)**
 1. Gravitational waves overview (nature in GR & sources)
 2. Newman Penrose formalism, Teukolsky equation

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:41 AM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 3 (4 of 83) Fit Page Width

Thumbnails

0

1

2

3

Week 1

Solution of Classical Field Equations Using Finite Difference Techniques

3

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:41 AM ubuntu-user

Lec1-1.pdf
File Edit View Go Help

Previous Next 4 (5 of 83) Fit Page Width

Week 1: References

- Mitchell, A. R., and D. F. Griffiths, **The Finite Difference Method in Partial Differential Equations**, New York: Wiley (1980)
- Richtmeyer, R. D., and Morton, K. W., **Difference Methods for Initial-Value Problems**, New York: Interscience (1967)
- H.-O. Kreiss and J. Oliger, **Methods for the Approximate Solution of Time Dependent Problems**, GARP Publications Series No. 10, (1973)
- Gustatsson, B., H. Kreiss and J. Oliger, **Time-Dependent Problems and Difference Methods**, New York: Wiley (1995)

Thumbnails

0

1

2

3

4

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:42 AM ubuntu-user


Lec1-1.pdf
File Edit View Go Help


Previous Next 6 (7 of 83) Fit Page Width


Preliminaries


- Classical field equations $\equiv$ time dependent partial differential equations (PDEs)
- Can divide time-dependent PDEs into two broad classes:
 1. **Initial-value Problems (Cauchy Problems)**, spatial domain has no boundaries (either infinite or “closed”—e.g. “periodic boundary conditions”)
 2. **Initial-Boundary-Value Problems**, spatial domain *finite*, need to specify boundary conditions
- **Note:** Even if *physical* problem is really of type 1, finite computational resources $\rightarrow$ finite spatial domain $\rightarrow$ approximate as type 2; will hereafter loosely refer to either type as an IVP.
- **Working Definition: Initial Value Problem**
 - State of physical system arbitrarily (usually) specified at some initial time $t = t_0$.
 - Solution exists for $t \geq t_0$; uniquely determined by equations of motion (EOM) and boundary conditions (BCs).

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf









Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:46 AM ubuntu-user

Lec1-1.pdf
File Edit View Go Help

Previous Next 7 (8 of 83) Fit Page Width

Thumbnails ▾

3

4

5

6

7

Preliminaries

- Approximate solution of initial value problems using *any* numerical method, including finite differencing, will always involve three key steps
 1. Complete mathematical specification of system of PDEs, including boundary conditions and initial conditions
 2. Discretization of the system: replacement of continuous domain by discrete domain, and approximation of differential equations by algebraic equations for discrete unknowns
 3. Solution of discrete algebraic equations
- Will assume that the set of PDEs has a unique solution for given initial conditions and boundary conditions, and that the solution does not “blow up” in time, unless such blow up is expected from the physics
- Whenever this last condition holds for an initial value problem, we say that the problem is well posed
- Note that this is a non-trivial issue in general relativity, since there are in practice *many* distinct forms the PDEs can take for a given physical scenario (in principle infinitely many) and not all will be well-posed in general

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:47 AM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 7 (8 of 83) Fit Page Width

Preliminaries

- Approximate solution of initial value problems using *any* numerical method, including finite differencing, will always involve three key steps
 1. Complete mathematical specification of system of PDEs, including boundary conditions and initial conditions
 2. Discretization of the system: replacement of continuous domain by discrete domain, and approximation of differential equations by algebraic equations for discrete unknowns
 3. Solution of discrete algebraic equations
- Will assume that the set of PDEs has a unique solution for given initial conditions and boundary conditions, and that the solution does not “blow up” in time, unless such blow up is expected from the physics
- Whenever this last condition holds for an initial value problem, we say that the problem is well posed
- Note that this is a non-trivial issue in general relativity, since there are in practice *many* distinct forms the PDEs can take for a given physical scenario (in principle infinitely many), and not all will be well-posed in general

Thumbnails

3

4

5

6

7

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

The image shows a man in a red jacket and grey trousers standing next to a large projection screen. He is holding a long, thin pointer stick and pointing at the screen. The screen displays a presentation slide titled "Preliminaries" from a PDF viewer. The slide contains a bulleted list of points about solving initial value problems for PDEs. The PDF viewer interface includes a menu bar (Applications, Places, System), a toolbar (Previous, Next, 7, 8 of 83, Fit PageWidth), and a sidebar with thumbnails. The system clock at the top right of the window shows 11:48 AM.

Applications Places System Ubuntu 11.04 - VMware Player File Virtual Machine Help 11:48 AM ubuntu-user

Lect1.pdf

File Edit View Go Help

Previous Next 7 (8 of 83) Fit PageWidth


Preliminaries

- Approximate solution of initial value problems using any numerical method, including finite differencing, will always involve three key steps
 1. Complete mathematical specification of system of PDEs, including boundary conditions and initial conditions
 2. Discretization of the system: replacement of continuous domain by discrete domain, and approximation of differential equations by algebraic equations for discrete unknowns
 3. Solution of discrete algebraic equations
- Will assume that the set of PDEs has a unique solution for given initial conditions and boundary conditions, and that the solution does not "blow up" in time, unless such blow up is expected from the physics
- Whenever this last condition holds for an initial value problem, we say that the problem is well posed
- Note that this is a non-trivial issue in general relativity, since there are in practice many distinct forms the PDEs can take for a given physical scenario (in principle infinitely many), and not all will be well-posed in general

PS: Explor... [Update Ma... ubuntu-user... Downloads Lect1.pdf Lect1-1.pdf

Well posedness


- 1- A sdn exists
- 2- Unique soln
- 3- Soln depends "continuously" on the
ID/BD



~ I Conf.


Well posedness

- 1- A soln exists
- 2- Unique soln
- 3- Soln is "continuously" on the ID/BD




Well posedness

- 1- A soln exists
- 2- Unique soln
- 3- Soln depends "continuously" on the
ID / BD



Well posedness

- 1- A soln exists
 - 2- Unique soln
 - 3- Soln depends "continuously" on the
ID/BD
-



$$u_{,t} = u_{,x}$$

$$u_{,t} = u_{,x} \rightarrow f_t = f \cdot i k$$

$$u(t, x) = f(t).$$

$$u_{,t} = u_{,x} \rightarrow$$


$$u(t,x) = f(t) \cdot e^{ikx}$$

$$f_t = f \cdot ik$$


$$f(t) = e^{ikt}$$

$$u(t,x) = e^{ik(t+x)}$$

$$\begin{aligned}
 u_{,t} &= u_{,x} \quad \rightarrow \\
 u(t,x) &= f(t) \cdot e^{ikx} \quad \rightarrow \quad f_{,t} = f \cdot ik \\
 &\quad \quad \quad ik(t+x) \quad \quad \quad f(t) = e^{ikt} \\
 u(t,x) &= e^{ik(t+x)} \\
 |u(t,x)| &= 1
 \end{aligned}$$




$$u_{i,t} = \delta u_{i,t+1}$$



$$u_{i,t} = \delta u_{i,t+1}$$

$$f_{i,t} = \delta (ik)^2 f_{i,t+1}$$

$$f_{i,t}$$




$$u_{,t} = \delta u_{,xxx}$$

$$f_{,t} = \delta (ik)^2 f$$

$$f_{,t} = -\delta k^2 f$$

$$p(t) = e^{-\delta k^2 t}$$

W




$$u_{,t} = \delta u_{,xxx}$$

$$f_{,t} = \delta (ik)^2 f$$

$$f_{,t} = -\delta k^2 \rightarrow f(t) = e^{-\delta k^2 t}$$

$$u(t) = e^{ikx}$$




$$u_{,t} = \delta u_{,xxx}$$

$$f_{,t} = \delta (ik)^2 f$$

$$f_{,t} = -\delta k^2 f$$

$$u(t) = e^{-\delta k^2 t}$$

$$|u(t)| = e^{-\delta k^2 t}$$

$$\square g = g_{,xxx}$$


I Conf.


$$u_{,t} = \delta u_{,xxx}$$

$$f_{,t} = \delta (ik)^2 f$$

$$f_{,t} = -\delta k^2 f \rightarrow f(t) = e^{-\delta k^2 t}$$

$$u(t) = e^{-\delta k^2 t} e^{ikx}$$

$$|u(t)| = e^{-\delta k^2 t}$$

$$\square g = g_{,xxx}$$


I Conf.



$$u_{,t} = \delta u_{,xx}$$

$$f_{,t} = \delta (ik)^2 f$$

$$f_{,t} = -\delta k^2 f \rightarrow f(t) = e^{-\delta k^2 t}$$

$$u(t) = e^{-\delta k^2 t} e^{ikx}$$

$$|u(t)| = e^{-\delta k^2 t}$$

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 11:59 AM ubuntu-user

Lec1-1.pdf
File Edit View Go Help

Previous Next 7 (8 of 83) Fit Page Width

Preliminaries

- Approximate solution of initial value problems using *any* numerical method, including finite differencing, will always involve three key steps
 1. Complete mathematical specification of system of PDEs, including boundary conditions and initial conditions
 2. Discretization of the system: replacement of continuous domain by discrete domain, and approximation of differential equations by algebraic equations for discrete unknowns
 3. Solution of discrete algebraic equations
- Will assume that the set of PDEs has a unique solution for given initial conditions and boundary conditions, and that the solution does not “blow up” in time, unless such blow up is expected from the physics
- Whenever this last condition holds for an initial value problem, we say that the problem is well posed
- Note that this is a non-trivial issue in general relativity, since there are in practice *many* distinct forms the PDEs can take for a given physical scenario (in principle infinitely many), and not all will be well-posed in general

Thumbnails

3

4

5

6

7

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:01 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 8 (9 of 83) Fit Page Width

Thumbnails

- 4
- 5
- 6
- 7
- 8

- There are several general approaches to the numerical solution of time dependent PDEs, including
 1. Finite differences
 2. Finite volume
 3. Finite elements
 4. Spectral
- Finite difference (FD) methods are particularly appropriate when the solution is expected to be smooth (infinitely differentiable) given that the initial data is smooth
- This is the case for many classical field theories including those for a scalar (linear/nonlinear Klein Gordon), vector (electromagnetism [Maxwell]), rank-2 symmetric tensor (general relativity [Einstein])
- In cases where solutions do *not* remain smooth, even if the initial data is—as happens in compressible hydrodynamics, for example, where shocks can form—the finite volume approach is the method of choice (next week)

8

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:01 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 9 (10 of 83) Fit Page Width

Thumbnails

5

6

7

8

9

Why Finite Differencing?

- Accessibility: Requires a minimum of mathematical background: if you're mathematically mature enough to understand the nature of the PDEs you need to solve, you're mathematically mature enough to understand finite differencing
- Flexibility: Technique can be used for essentially any system of PDEs that has smooth solutions, irrespective of
 - Number of dependent variables (unknown functions)
 - Number of independent variables (a.k.a. "dimensionality" of the system: nomenclature "1-D" means dependence on one spatial dimension plus time, "2-D", "3-D" similarly mean dependence on two/three dimensions, plus time, respectively)
 - Nonlinearity
 - Form of equations: technique does not require that the system of equations has any particular/special form (contrast with finite volume methods where one generally wants to cast the equations in so-called conservation-law form)

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:02 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 10 (11 of 83) Fit Page Width

Why Finite Differencing?

- Error analysis:
 - Mathematically rigorous: Quite difficult
 - Practical/empirical: Extremely straightforward—basic principle is to compute multiple solutions using same initial data and problem parameters, but differing fundamental discretization scales. Comparison of solutions provides direct estimate of error in solutions
- Adaptivity: Can combine basic method with changes in
 - Local scale of discretization
 - Order of approximation

in order to maximize increase in solution accuracy as a function of computational work invested (e.g. adaptive mesh refinement, week 3)
- Parallelization: Due to "locality of influence" in finite difference schemes, it is relatively easy to write FD codes than run efficiently on large distributed memory computer clusters having 1000s or cores (these days 10,000s or even 100,000s!)

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:04 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 13 (14 of 83) Fit Page Width

Thumbnails

9

10

11

12

13

The 1-D Wave Equation

- Consider the following initial value (Cauchy) problem for the scalar function $\phi(t, x)$

$$\phi_{tt} = c^2 \phi_{xx}, \quad -\infty \leq x \leq \infty, \quad t \geq 0 \quad (1)$$

$$\phi(0, x) = \phi_0(x) \quad (2)$$

$$\phi_t(0, x) = \Pi_0(x) \quad (3)$$

where c is a positive constant, we have adopted the subscript notation for partial differentiation, e.g. $\phi_{tt} \equiv \partial^2 \phi / \partial t^2$, and we wish to determine $\phi(t, x)$ in the solution domain from the initial conditions (2-3) and the governing equation (1)
- Note the following:
 - Since the spatial domain is unbounded, there are *no* boundary conditions
 - Since the equation is second order in time, two functions-worth of initial data must be specified: the initial scalar field profile, $\phi_0(x)$, and the initial time derivative, $\Pi_0(x)$
 - This system is well posed, and if the initial conditions $\phi_0(x)$ and $\Pi_0(x)$ are smooth—which we will hereafter assume—so is the complete solution $\phi(t, x)$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:04 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 13 (14 of 83) Fit Page Width

The 1-D Wave Equation

- Consider the following initial value (Cauchy) problem for the scalar function $\phi(t, x)$

$$\phi_{tt} = c^2 \phi_{xx}, \quad -\infty \leq x \leq \infty, \quad t \geq 0 \quad (1)$$

$$\phi(0, x) = \phi_0(x) \quad (2)$$

$$\phi_t(0, x) = \Pi_0(x) \quad (3)$$

where c is a positive constant, we have adopted the subscript notation for partial differentiation, e.g. $\phi_{tt} \equiv \partial^2 \phi / \partial t^2$, and we wish to determine $\phi(t, x)$ in the solution domain from the initial conditions (2-3) and the governing equation (1)
- Note the following:
 - Since the spatial domain is unbounded, there are *no* boundary conditions
 - Since the equation is second order in time, two functions-worth of initial data must be specified: the initial scalar field profile, $\phi_0(x)$, and the initial time derivative, $\Pi_0(x)$
 - This system is well posed, and if the initial conditions $\phi_0(x)$ and $\Pi_0(x)$ are smooth—which we will hereafter assume—so is the complete solution $\phi(t, x)$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

$$\begin{aligned}
 u_{,t} &= u_{,x} \quad \rightarrow \\
 u(t,x) &= f(t) \cdot e^{ikx} & f_{,t} &= f \cdot ik \\
 & \quad \quad \quad ik(t+x) & f(t) &= e^{ikt} \\
 u(t,x) &= e^{ik(t+x)} \\
 |u(t,x)| &= 1
 \end{aligned}$$

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:05 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 13 (14 of 83) Fit Page Width

Thumbnails

9

10

11

12

13

- Consider the following initial value (Cauchy) problem for the scalar function $\phi(t, x)$

$$\phi_{tt} = c^2 \phi_{xx}, \quad -\infty \leq x \leq \infty, \quad t \geq 0 \quad (1)$$

$$\phi(0, x) = \phi_0(x) \quad (2)$$

$$\phi_t(0, x) = \Pi_0(x) \quad (3)$$

where c is a positive constant, we have adopted the subscript notation for partial differentiation, e.g. $\phi_{tt} \equiv \partial^2 \phi / \partial t^2$, and we wish to determine $\phi(t, x)$ in the solution domain from the initial conditions (2-3) and the governing equation (1)

- Note the following:
 - Since the spatial domain is unbounded, there are *no* boundary conditions
 - Since the equation is second order in time, two functions-worth of initial data must be specified: the initial scalar field profile, $\phi_0(x)$, and the initial time derivative, $\Pi_0(x)$
 - This system is well posed, and if the initial conditions $\phi_0(x)$ and $\Pi_0(x)$ are smooth—which we will hereafter assume—so is the complete solution $\phi(t, x)$

13

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:06 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 14 (15 of 83) Fit Page Width

The 1-D Wave Equation

- Eqn. (1) is a *hyperbolic* PDE, and as such, its solutions generically describe the propagation of disturbances at some finite speed(s), which in this case is c
- Without loss of generality, we can assume that we have adopted units in which this speed satisfies $c = 1$. Our problem then becomes

$$\phi_{tt} = \phi_{xx}, \quad -\infty \leq x \leq \infty, \quad t \geq 0 \quad (4)$$

$$\phi(0, x) = \phi_0(x) \quad (5)$$

$$\phi_t(0, x) = \Pi_0(x) \quad (6)$$

- In the study of the solutions of hyperbolic PDEs, using either closed form (preferred to "analytic") or numerical approaches, the concept of characteristic is crucial
- Loosely, in a spacetime diagram, characteristics are the lines/surfaces along which information/signals propagate(s).

Thumbnails

10

11

12

13

14

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:06 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 15 (16 of 83) Fit Page Width

Thumbnails

11

12

13

14

15

The 1-D Wave Equation

..... : "left-directed" characteristics, $x + t = \text{constant}$, $l(x + t)$
 : "right-directed" characteristics, $x - t = \text{constant}$, $r(x - t)$

- General solution of (4) is a superposition of an arbitrary *left-moving* profile ($v = -c = -1$), and an arbitrary *right-moving* profile ($v = +c = +1$); i.e.

$$\phi(t, x) = \ell(x + t) + r(x - t) \quad (7)$$

where

ℓ : constant along "left-directed" characteristics

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:07 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 15 (16 of 83) Fit Page Width

Thumbnails

11

12

13

14

15

The 1-D Wave Equation

..... : "left-directed" characteristics, $x + t = \text{constant}$, $l(x + t)$

..... : "right-directed" characteristics, $x - t = \text{constant}$, $r(x - t)$

- General solution of (4) is a superposition of an arbitrary *left-moving* profile ($v = -c = -1$), and an arbitrary *right-moving* profile ($v = +c = +1$); i.e.

$$\phi(t, x) = \ell(x + t) + r(x - t) \quad (7)$$

where

ℓ : constant along "left-directed" characteristics

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:07 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 15 (16 of 83) Fit Page Width

Thumbnails

11

12

13

14

15

The 1-D Wave Equation

..... : "left-directed" characteristics, $x + t = \text{constant}$, $l(x + t)$

..... : "right-directed" characteristics, $x - t = \text{constant}$, $r(x - t)$

- General solution of (4) is a superposition of an arbitrary *left-moving* profile ($v = -c = -1$), and an arbitrary *right-moving* profile ($v = +c = +1$); i.e.

$$\phi(t, x) = \ell(x + t) + r(x - t) \quad (7)$$

where

ℓ : constant along "left-directed" characteristics

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:08 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 16 (17 of 83) Fit Page Width

The 1-D Wave Equation

- Observation provides alternative way of specifying initial values—often convenient in practice
- Rather than specifying $u(x, 0)$ and $u_t(x, 0)$ directly, specify *initial* left-moving and right-moving parts of the solution, $\ell(x)$ and $r(x)$
- Specifically, set

$$\phi(x, 0) = \ell(x) + r(x) \quad (8)$$

$$\phi_t(x, 0) = \ell'(x) - r'(x) \equiv \frac{d\ell}{dx}(x) - \frac{dr}{dx}(x) \quad (9)$$
- For illustrative purposes will frequently take profile functions $\phi_0(x)$, $\ell(x)$, $r(x)$ to be “gaussians”, e.g.

$$\phi_0(x) = A \exp \left[-((x - x_0) / \delta)^2 \right] \quad (10)$$

where A , x_0 and δ are viewed as adjustable parameters that control the overall size/height of the profile (A), its centre point (x_0) and its effective width (δ)

Thumbnails

12

13

14

15

16

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:08 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 16 (17 of 83) Fit Page Width

Thumbnails

- 12
- 13
- 14
- 15
- 16

- Rather than specifying $u(x, 0)$ and $u_t(x, 0)$ directly, specify *initial* left-moving and right-moving parts of the solution, $\ell(x)$ and $r(x)$
- Specifically, set

$$\phi(x, 0) = \ell(x) + r(x) \quad (8)$$

$$\phi_t(x, 0) = \ell'(x) - r'(x) \equiv \frac{d\ell}{dx}(x) - \frac{dr}{dx}(x) \quad (9)$$
- For illustrative purposes will frequently take profile functions $\phi_0(x)$, $\ell(x)$, $r(x)$ to be "gaussians", e.g.

$$\phi_0(x) = A \exp \left[-((x - x_0) / \delta)^2 \right] \quad (10)$$

where A , x_0 and δ are viewed as adjustable parameters that control the overall size/height of the profile (A), its centre point (x_0) and its effective width (δ)

16

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:08 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 17 (18 of 83) Fit Page Width

Thumbnails

13

14

15

16

17

- For illustrative purposes will frequently take profile functions $\phi_0(x)$, $\ell(x)$, $r(x)$ to be "gaussians", e.g.


$$\phi_0(x) = A \exp \left[-((x - x_0) / \delta)^2 \right] \quad (10)$$


where A , x_0 and δ are viewed as adjustable parameters that control the overall size/height of the profile (A), its centre point (x_0) and its effective width (δ)

16

2. Discretization


PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf





$$u(t, x) \longrightarrow u_J^n = u(n \Delta t, J \Delta x)$$

$$u(t, x) \longrightarrow u_J^n = u(n \Delta t, J \Delta x)$$



Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:11 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 19 (20 of 83) Fit Page Width

Thumbnails

15


16

17

18

19

Deriving Finite Difference Formulae



- One-dimensional, uniform finite difference mesh.
- Note that the spacing, $\Delta x = h$, between adjacent mesh points is *constant*.
- Will tacitly assume that the origin, x_0 , of coordinate system is $x_0 = 0$.

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:11 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 19 (20 of 83) Fit Page Width

Thumbnails

15


16

17

18

19

Deriving Finite Difference Formulae



- One-dimensional, uniform finite difference mesh.
- Note that the spacing, $\Delta x = h$, between adjacent mesh points is *constant*.
- Will tacitly assume that the origin, x_0 , of coordinate system is $x_0 = 0$.

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:11 PM ubuntu-user


Lec1-1.pdf

File Edit View Go Help

Previous Next 19 (20 of 83) Fit Page Width

18


Deriving Finite Difference Formulae



- One-dimensional, uniform finite difference mesh.
- Note that the spacing, $\Delta x = h$, between adjacent mesh points is *constant*.
- Will tacitly assume that the origin, x_0 , of coordinate system is $x_0 = 0$.

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

$$u(t, x) \longrightarrow u^n_J = u(n \Delta t, J \Delta x)$$



Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:13 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 20 (21 of 83) Fit Page Width

Thumbnails

16

17

18

19

20

Deriving Finite Difference Formulae

- Given the three values $u(x_j - h)$, $u(x_j)$ and $u(x_j + h)$, denoted u_{j-1} , u_j , and u_{j+1} respectively, can compute an $O(h^2)$ approximation to $u_x(x_j) \equiv (u_x)_j$ as follows
- Taylor expanding, have

$$\begin{aligned}
 u_{j-1} &= u_j - h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j - \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5) \\
 u_j &= u_j \\
 u_{j+1} &= u_j + h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j + \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5)
 \end{aligned}$$
- Now seek a linear combination of u_{j-1} , u_j , and u_{j+1} which yields $(u_x)_j$ to $O(h^2)$ accuracy, i.e. we seek c_- , c_0 and c_+ such that

$$c_- u_{j-1} + c_0 u_j + c_+ u_{j+1} = (u_x)_j + O(h^2)$$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:13 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 20 (21 of 83) Fit Page Width

Thumbnails

16

17

18

19

20

Deriving Finite Difference Formulae

- Given the three values $u(x_j - h)$, $u(x_j)$ and $u(x_j + h)$, denoted u_{j-1} , u_j , and u_{j+1} respectively, can compute an $O(h^2)$ approximation to $u_x(x_j) \equiv (u_x)_j$ as follows
- Taylor expanding, have

$$\begin{aligned}
 u_{j-1} &= u_j - h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j - \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5) \\
 u_j &= u_j \\
 u_{j+1} &= u_j + h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j + \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5)
 \end{aligned}$$
- Now seek a linear combination of u_{j-1} , u_j , and u_{j+1} which yields $(u_x)_j$ to $O(h^2)$ accuracy, i.e. we seek c_- , c_0 and c_+ such that

$$c_- u_{j-1} + c_0 u_j + c_+ u_{j+1} = (u_x)_j + O(h^2)$$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:14 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 20 (21 of 83) Fit Page Width

Thumbnails

Deriving Finite Difference Formulae

- Given the three values $u(x_j - h)$, $u(x_j)$ and $u(x_j + h)$, denoted u_{j-1} , u_j , and u_{j+1} respectively, can compute an $O(h^2)$ approximation to $u_x(x_j) \equiv (u_x)_j$ as follows
- Taylor expanding, have

$$\begin{aligned}
 u_{j-1} &= u_j - h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j - \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5) \\
 u_j &= u_j \\
 u_{j+1} &= u_j + h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j + \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5)
 \end{aligned}$$
- Now seek a linear combination of u_{j-1} , u_j , and u_{j+1} which yields $(u_x)_j$ to $O(h^2)$ accuracy, i.e. we seek c_- , c_0 and c_+ such that

$$c_- u_{j-1} + c_0 u_j + c_+ u_{j+1} = (u_x)_j + O(h^2)$$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:14 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 20 (21 of 83) Fit Page Width

Thumbnails

16

17

18

19

20

Deriving Finite Difference Formulae

- Given the three values $u(x_j - h)$, $u(x_j)$ and $u(x_j + h)$, denoted u_{j-1} , u_j , and u_{j+1} respectively, can compute an $O(h^2)$ approximation to $u_x(x_j) \equiv (u_x)_j$ as follows
- Taylor expanding, have

$$\begin{aligned}
 u_{j-1} &= u_j - h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j - \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5) \\
 u_j &= u_j \\
 u_{j+1} &= u_j + h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j + \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5)
 \end{aligned}$$
- Now seek a linear combination of u_{j-1} , u_j , and u_{j+1} which yields $(u_x)_j$ to $O(h^2)$ accuracy, i.e. we seek c_- , c_0 and c_+ such that

$$c_- u_{j-1} + c_0 u_j + c_+ u_{j+1} = (u_x)_j + O(h^2)$$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:15 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 20 (21 of 83) Fit Page Width

Thumbnails

16

17

18

19

20

follows

- Taylor expanding, have

$$u_{j-1} = u_j - h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j - \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5)$$

$$u_j = u_j$$

$$u_{j+1} = u_j + h(u_x)_j + \frac{1}{2}h^2(u_{xx})_j + \frac{1}{6}h^3(u_{xxx})_j + \frac{1}{24}h^4(u_{xxxx})_j + O(h^5)$$

- Now seek a linear combination of u_{j-1} , u_j , and u_{j+1} which yields $(u_x)_j$ to $O(h^2)$ accuracy, i.e. we seek c_- , c_0 and c_+ such that

$$c_- u_{j-1} + c_0 u_j + c_+ u_{j+1} = (u_x)_j + O(h^2)$$

20

Deriving Finite Difference Formulae

- Results in a system of three linear equations for u_{j-1} , u_j , and u_{j+1} :

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:16 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 21 (22 of 83) Fit Page Width

Deriving Finite Difference Formulae

- Results in a system of three linear equations for u_{j-1} , u_j , and u_{j+1} :

$$\begin{aligned} c_- + c_0 + c_+ &= 0 \\ -hc_- + hc_+ &= 1 \\ \frac{1}{2}h^2c_- + \frac{1}{2}h^2c_+ &= 0 \end{aligned}$$

which has the solution

$$\begin{aligned} c_- &= -\frac{1}{2h} \\ c_0 &= 0 \\ c_+ &= +\frac{1}{2h} \end{aligned}$$

- Thus, $O(h^2)$ FDA (finite difference approximation) for the first derivative is

$$\frac{u(x+h) - u(x-h)}{2h} = u_x(x) + O(h^2) \quad (11)$$

21

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:16 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 21 (22 of 83) Fit Page Width

Deriving Finite Difference Formulae

- Results in a system of three linear equations for u_{j-1} , u_j , and u_{j+1} :

$$\begin{aligned} c_- + c_0 + c_+ &= 0 \\ -hc_- + hc_+ &= 1 \\ \frac{1}{2}h^2c_- + \frac{1}{2}h^2c_+ &= 0 \end{aligned}$$

which has the solution

$$\begin{aligned} c_- &= -\frac{1}{2h} \\ c_0 &= 0 \\ c_+ &= +\frac{1}{2h} \end{aligned}$$

- Thus, $O(h^2)$ FDA (finite difference approximation) for the first derivative is

$$\frac{u(x+h) - u(x-h)}{2h} = u_x(x) + O(h^2) \quad (11)$$

21

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:16 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 21 (22 of 83) Fit Page Width

Deriving Finite Difference Formulae

- Results in a system of three linear equations for u_{j-1} , u_j , and u_{j+1} :

$$\begin{aligned} c_- + c_0 + c_+ &= 0 \\ -hc_- + hc_+ &= 1 \\ \frac{1}{2}h^2c_- + \frac{1}{2}h^2c_+ &= 0 \end{aligned}$$

which has the solution

$$\begin{aligned} c_- &= -\frac{1}{2h} \\ c_0 &= 0 \\ c_+ &= +\frac{1}{2h} \end{aligned}$$

- Thus, $O(h^2)$ FDA (finite difference approximation) for the first derivative is

$$\frac{u(x+h) - u(x-h)}{2h} = u_x(x) + O(h^2) \quad (11)$$

21

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:17 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 21 (22 of 83) Fit Page Width

Thumbnails

17

18

19

20

21

$$\frac{1}{2}n^-c_- + \frac{1}{2}n^-c_+ = 0$$

which has the solution

$$\begin{aligned} c_- &= -\frac{1}{2h} \\ c_0 &= 0 \\ c_+ &= +\frac{1}{2h} \end{aligned}$$

- Thus, $O(h^2)$ FDA (finite difference approximation) for the first derivative is

$$\frac{u(x+h) - u(x-h)}{2h} = u_x(x) + O(h^2) \quad (11)$$

21

Deriving Finite Difference Formulae

- May not be obvious *a priori*, that the truncation error of approximation is $O(h^2)$
- Naive consideration of the number of terms in the Taylor series expansion which can be eliminated using 2 values (namely $u(x+h)$ and $u(x-h)$)

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

$$u(t, x) \longrightarrow u_J^n = u(n \Delta t, J \Delta x)$$

$$\begin{array}{c} \Delta t \\ \hline \Delta t \end{array}$$

$$\Delta t$$

$$\Delta x$$

$$\Delta t = \lambda$$

$$\begin{array}{c} \hline \downarrow \\ h \end{array}$$

$$u(t, x) \longrightarrow u_J^n = u(n \Delta t, J \Delta x)$$

$$\left. \frac{\partial u}{\partial x} \right|_{x=x_J} = \frac{u_{J+1} - u_{J-1}}{2h}$$

$$\begin{array}{cc} \Delta t & \Delta x \\ & \Downarrow \\ \Delta t = \lambda & h \end{array}$$

$$u(t, x) \longrightarrow u_J^n = u(n \Delta t, J \Delta x)$$

$$\left. \frac{\partial u}{\partial x} \right|_{x=x_J}^n = \frac{u_{J+1}^n - u_{J-1}^n}{2h}$$

$$\left. \frac{\partial u}{\partial t} \right|_J^n = \frac{u_J^{n+1} - u_J^{n-1}}{2(\Delta t)}$$

$$\Delta t \quad \Delta x$$

$$\Delta t = \lambda \quad h$$

$$u_t = u_x \quad \text{for } u(t, x)$$

$$u(t, x) \longrightarrow u_J^n = u(n \Delta t, J \Delta x)$$

$$u|_{x=x_J} = \frac{u_J^n - u_{J-1}^n}{\Delta x}$$

$$u_t = \frac{u_J^{n+1} - u_J^n}{\Delta t}$$

$$\Delta t = \lambda \quad \Delta x = h$$

$$u_t = u_x \quad \text{for } u(t, x)$$

$$u(t, x) \longrightarrow u_J^n = u(n \Delta t, J \Delta x)$$

$$\left. \frac{\partial u}{\partial x} \right|_{x=x_J}^n = \frac{u_{J+1}^n - u_{J-1}^n}{2h}$$

$$\Delta t \quad \Delta x$$

$$\Delta t = \lambda \quad h$$

$$\left. \frac{\partial u}{\partial t} \right|_J^n = \frac{u_J^{n+1} - u_J^{n-1}}{2(\Delta t)}$$

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:19 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 22 (23 of 83) Fit Page Width

Thumbnails

18

19

20

21

22

$$\frac{u(x+h) - 2u(x) + u(x-h))}{h^2} = u_{xx}(x) + O(h^2) \quad (11)$$

Deriving Finite Difference Formulae

- May not be obvious *a priori*, that the truncation error of approximation is $O(h^2)$
- Naive consideration of the number of terms in the Taylor series expansion which can be eliminated using 2 values (namely $u(x+h)$ and $u(x-h)$) suggests that the error might be $O(h)$.
- Fact that the $O(h)$ term “drops out” a consequence of the *symmetry*, or *centering* of the stencil: common theme in such FDA, called *centred* difference approximations
- Using same technique, can easily generate $O(h^2)$ expression for the *second* derivative, which uses the same difference stencil as the above approximation for the first derivative.

$$\frac{u(x+h) - 2u(x) + u(x-h))}{h^2} = u_{xx}(x) + O(h^2) \quad (12)$$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:21 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 22 (23 of 83) Fit Page Width

Thumbnails

18

19

20

21

22

$$\frac{u(x+h) - u(x-h)}{2h} = u_x(x) + O(h^2) \quad (11)$$

Deriving Finite Difference Formulae

- May not be obvious *a priori*, that the truncation error of approximation is $O(h^2)$
- Naive consideration of the number of terms in the Taylor series expansion which can be eliminated using 2 values (namely $u(x+h)$ and $u(x-h)$) suggests that the error might be $O(h)$.
- Fact that the $O(h)$ term “drops out” a consequence of the *symmetry*, or *centering* of the stencil: common theme in such FDA, called *centred* difference approximations
- Using same technique, can easily generate $O(h^2)$ expression for the *second* derivative, which uses the same difference stencil as the above approximation for the first derivative.

$$\frac{u(x+h) - 2u(x) + u(x-h))}{h^2} = u_{xx}(x) + O(h^2) \quad (12)$$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:21 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 22 (23 of 83) Fit Page Width

Thumbnails

18

19

20

21

22

suggests that the error might be $O(h)$.

- Fact that the $O(h)$ term “drops out” a consequence of the *symmetry*, or *centering* of the stencil: common theme in such FDA, called *centred* difference approximations
- Using same technique, can easily generate $O(h^2)$ expression for the *second* derivative, which uses the same difference stencil as the above approximation for the first derivative.

$$\frac{u(x+h) - 2u(x) + u(x-h))}{h^2} = u_{xx}(x) + O(h^2) \quad (12)$$

- *Exercise:* Compute the precise form of the $O(h^2)$ terms in expressions (11) and (12).

22

Sample FDA for the 1-D Wave Equation

- Let us consider the 1-D wave equation again, but this time on the finite spatial

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:22 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 23 (24 of 83) Fit Page Width

Thumbnails

- 19
- 20
- 21
- 22
- 23

- Let us consider the 1-D wave equation again, but this time on the finite spatial domain, $0 \leq x \leq 1$, where we will prescribe fixed (Dirichlet) boundary conditions
- Then we wish to solve

$$\phi_{tt} = \phi_{xx} \quad (c = 1) \quad 0 \leq x \leq 1, \quad t \geq 0 \quad (13)$$

$$\phi(0, x) = \phi_0(x)$$

$$\phi_t(0, x) = \Pi_0(x)$$

$$\phi(t, 0) = \phi(t, 1) = 0 \quad (14)$$
- We will again require that the initial data functions, $\phi_0(x)$ and $\Pi_0(x)$ be smooth
- Moreover, in order to ensure a smooth solution everywhere, the initial values must be compatible with the boundary conditions, i.e.

$$\phi_0(0) = \phi_0(1) = \Pi_0(0) = \Pi_0(1) = 0 \quad (15)$$

23

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:22 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 23 (24 of 83) Fit Page Width

Thumbnails

- 19
- 20
- 21
- 22
- 23

- Let us consider the 1-D wave equation again, but this time on the finite spatial domain, $0 \leq x \leq 1$, where we will prescribe fixed (Dirichlet) boundary conditions
- Then we wish to solve

$$\phi_{tt} = \phi_{xx} \quad (c = 1) \quad 0 \leq x \leq 1, \quad t \geq 0 \quad (13)$$

$$\phi(0, x) = \phi_0(x)$$

$$\phi_t(0, x) = \Pi_0(x)$$

$$\phi(t, 0) = \phi(t, 1) = 0 \quad (14)$$
- We will again require that the initial data functions, $\phi_0(x)$ and $\Pi_0(x)$ be smooth
- Moreover, in order to ensure a smooth solution everywhere, the initial values must be compatible with the boundary conditions, i.e.

$$\phi_0(0) = \phi_0(1) = \Pi_0(0) = \Pi_0(1) = 0 \quad (15)$$

23

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:23 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 24 (25 of 83) Fit Page Width

Thumbnails

20

21

22

23

24

$\phi_0(0) = \phi_0(1) = \psi_0(0) = \psi_0(1) = 0$ (15)

Sample FDA for the 1-D Wave Equation

- As always, we begin the discretization process by replacing the continuum solution domain with a finite difference mesh, whose typical element (point/event) we will denote by (x_j, t^n) :

$$t^n \equiv n \Delta t, \quad n = 0, 1, 2, \dots$$

$$x_j \equiv (j-1) \Delta x, \quad j = 1, 2, \dots, J$$

$$\phi_j^n \equiv \phi(n \Delta t, (j-1) \Delta x)$$

$$\Delta x = (J-1)^{-1}$$

$$\Delta t = \lambda \Delta x \quad \lambda \equiv \text{"Courant number"}$$
- We note in passing that the quantity λ defined above is often called the Courant number or Courant factor, after the great 20th century mathematician Richard Courant who was a pioneer in the study of finite difference solutions of time dependent PDEs (in particular, in the use of FD techniques to establish existence and uniqueness of such PDEs)

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:24 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 25 (26 of 83) Fit Page Width

Thumbnails

21

22

23

24

25

Uniform Grid for 1-D Wave Equation

• When solving wave equations using FDAs, typically keep λ constant when Δx varied.

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:24 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 26 (27 of 83) Fit Page Width

Thumbnails

22

23

24

25

26

$$\Delta x \equiv h$$


$$\Delta t \equiv \lambda h$$

25

Stencil for "Standard" $O(h^2)$ Approximation of 1-D Wave Equation

Diagram illustrating the stencil for the "Standard" $O(h^2)$ Approximation of 1-D Wave Equation. The stencil shows a central point (j, n) surrounded by four points: $(j-1, n)$, $(j+1, n)$, $(j, n-1)$, and $(j, n+1)$. The points are connected by dashed lines, forming a cross shape. The labels $j-1$, j , $j+1$ are at the bottom, and $n-1$, n , $n+1$ are to the right.

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf



$$\underline{\phi_{1,t}} = \phi_{1,xx}$$



Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:26 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 26 (27 of 83) Fit Page Width

Thumbnails

22

23

24

25

26

$$\Delta x \equiv h$$


$$\Delta t \equiv \lambda h$$

25

Stencil for "Standard" $O(h^2)$ Approximation of 1-D Wave Equation

$n + 1$
 n
 $n - 1$
 $j - 1$ j $j + 1$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf


$$\square g = g_{xxx}$$


I Conf.

$$\underbrace{\phi_{144}} = \phi_{1xx}$$

$$\phi^{n+1}$$




$$\square g = g_{,xxx}$$


I Conf.

$$\phi_{,tt} = \phi_{,xx}$$

$$\frac{\phi_{J+1}^{n+1} - 2\phi_J^n + \phi_{J-1}^{n-1}}{\Delta t^2} = \frac{\phi_{J+1}^n - 2\phi_J^n + \phi_{J-1}^n}{\Delta x^2}$$

$$\square g = g_{,xx}$$



I Conf.

$$\phi_{,tt} = \phi_{,xx}$$

Knows up to level $n+1$

$$\phi_{,tt}^{n+1} = \phi_{,xx}^{n+1} = \phi_{,xx}^n + \phi_{,xx}^{n-1} - 2\phi_{,xx}^n + \phi_{,xx}^{n-1}$$

Δt^2 Δx^2

$$\square g = g_{,xx}$$


I Conf.

$$\phi_{,tt} = \phi_{,xx}$$

Knows up to level
n+1

$$\frac{\phi_{J+1}^{n+1} - 2\phi_J^n + \phi_{J-1}^{n-1}}{\Delta t^2} = \frac{\phi_{J+1}^n - 2\phi_J^n + \phi_{J-1}^n}{\Delta x^2}$$

$$\phi_J^{n+1} = \text{Something}(\phi_J^n, \phi_{J-1}^{n-1})$$

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:28 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 27 (28 of 83) Fit Page Width

Thumbnails

23

24

25

26

27

FDA for 1-D Wave Equation

- Discretized Interior equation

$$\begin{aligned}
 (\Delta t)^{-2} (\phi_j^{n+1} - 2\phi_j^n + \phi_j^{n-1}) &= (\phi_{tt})_j^n + \frac{1}{12} \Delta t^2 (\phi_{tttt})_j^n + O(\Delta t^4) \\
 &= (\phi_{tt})_j^n + O(h^2) \\
 (\Delta x)^{-2} (\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n) &= (\phi_{xx})_j^n + \frac{1}{12} \Delta x^2 (\phi_{xxxx})_j^n + O(\Delta x^4) \\
 &= (\phi_{xx})_j^n + O(h^2)
 \end{aligned}$$

Putting these two together, get $O(h^2)$ approximation

$$\frac{\phi_j^{n+1} - 2\phi_j^n + \phi_j^{n-1}}{\Delta t^2} = \frac{\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n}{\Delta x^2} \quad j = 2, 3, \dots, J-1 \quad (16)$$

- Scheme such as (16) often called a *three level scheme* since couples *three "time levels"* of data (i.e. unknowns at three distinct, discrete times t^{n-1}, t^n, t^{n+1}).

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:28 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 27 (28 of 83) Fit Page Width

Thumbnails

23

24

25

26

27

FDA for 1-D Wave Equation

- Discretized Interior equation

$$\begin{aligned}
 (\Delta t)^{-2} (\phi_j^{n+1} - 2\phi_j^n + \phi_j^{n-1}) &= (\phi_{tt})_j^n + \frac{1}{12} \Delta t^2 (\phi_{tttt})_j^n + O(\Delta t^4) \\
 &= (\phi_{tt})_j^n + O(h^2) \\
 (\Delta x)^{-2} (\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n) &= (\phi_{xx})_j^n + \frac{1}{12} \Delta x^2 (\phi_{xxxx})_j^n + O(\Delta x^4) \\
 &= (\phi_{xx})_j^n + O(h^2)
 \end{aligned}$$

Putting these two together, get $O(h^2)$ approximation

$$\frac{\phi_j^{n+1} - 2\phi_j^n + \phi_j^{n-1}}{\Delta t^2} = \frac{\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n}{\Delta x^2} \quad j = 2, 3, \dots, J-1 \quad (16)$$

- Scheme such as (16) often called a *three level scheme* since couples *three "time levels"* of data (i.e. unknowns at three distinct, discrete times t^{n-1}, t^n, t^{n+1}).

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:29 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 28 (29 of 83) Fit Page Width

Thumbnails

FDA for 1-D Wave Equation

- Discretized Boundary conditions

$$\phi_1^{n+1} = \phi_J^{n+1} = 0$$
- Discretized Initial conditions
 - Need to specify two "time levels" of data (effectively $\phi(x, 0)$ and $\phi_t(x, 0)$), i.e. we must specify

$$\phi_j^0, \quad j = 1, 2, \dots, J$$

$$\phi_j^1, \quad j = 1, 2, \dots, J$$
 ensuring that the initial values are compatible with the boundary conditions.
- Can solve (16) explicitly for ϕ_j^{n+1} :

$$\phi_j^{n+1} = 2\phi_j^n - \phi_j^{n-1} + \lambda^2 (\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n) \quad (17)$$

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:30 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 28 (29 of 83) Fit Page Width

Thumbnails

- 24
- 25
- 26
- 27
- 28

- Need to specify two "time levels" of data (effectively $\phi(x, 0)$ and $\phi_t(x, 0)$), i.e. we must specify

$$\phi_j^0, \quad j = 1, 2, \dots, J$$

$$\phi_j^1, \quad j = 1, 2, \dots, J$$
 ensuring that the initial values are compatible with the boundary conditions.
- Can solve (16) explicitly for ϕ_j^{n+1} :

$$\phi_j^{n+1} = 2\phi_j^n - \phi_j^{n-1} + \lambda^2 (\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n) \quad (17)$$

28

FDA for 1-D Wave Equation

- Also note that (17) is actually a *linear* system for the unknowns $\phi_j^{n+1}, j = 1, 2, \dots, J$; in combination with the discrete boundary conditions

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:30 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 28 (29 of 83) Fit Page Width

Thumbnails

- 24
- 25
- 26
- 27
- 28

- Need to specify two "time levels" of data (effectively $\phi(x, 0)$ and $\phi_t(x, 0)$), i.e. we must specify

$$\phi_j^0, \quad j = 1, 2, \dots, J$$

$$\phi_j^1, \quad j = 1, 2, \dots, J$$
 ensuring that the initial values are compatible with the boundary conditions.
- Can solve (16) explicitly for ϕ_j^{n+1} :

$$\phi_j^{n+1} = 2\phi_j^n - \phi_j^{n-1} + \lambda^2 (\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n) \quad (17)$$

28

FDA for 1-D Wave Equation

- Also note that (17) is actually a *linear system* for the unknowns $\phi_j^{n+1}, j = 1, 2, \dots, J$; in combination with the discrete boundary conditions

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:30 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 27 (28 of 83) Fit Page Width

Thumbnails

24

25

26

27

28

FDA for 1-D Wave Equation

- Discretized Interior equation


$$\begin{aligned}
 (\Delta t)^{-2} (\phi_j^{n+1} - 2\phi_j^n + \phi_j^{n-1}) &= (\phi_{tt})_j^n + \frac{1}{12} \Delta t^2 (\phi_{tttt})_j^n + O(\Delta t^4) \\
 &= (\phi_{tt})_j^n + O(h^2) \\
 (\Delta x)^{-2} (\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n) &= (\phi_{xx})_j^n + \frac{1}{12} \Delta x^2 (\phi_{xxxx})_j^n + O(\Delta x^4) \\
 &= (\phi_{xx})_j^n + O(h^2)
 \end{aligned}$$

Putting these two together, get $O(h^2)$ approximation

$$\frac{\phi_j^{n+1} - 2\phi_j^n + \phi_j^{n-1}}{\Delta t^2} = \frac{\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n}{\Delta x^2} \quad j = 2, 3, \dots, J-1 \quad (16)$$

- Scheme such as (16) often called a *three level scheme* since couples *three "time levels"* of data (i.e. unknowns at three distinct, discrete times t^{n-1}, t^n, t^{n+1}).

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

$$\square g = g_{,xx}$$


I Conf.

$$\phi_{,tt} = \phi_{,xx}$$

Knows up to level $n+1$

$$\frac{\phi_{,t}^{n+1} - 2\phi_{,t}^n + \phi_{,t}^{n-1}}{\Delta t^2} = \frac{(\phi_{,t}^n - 2\phi_{,t}^n + \phi_{,t}^{n-1}))}{\Delta x^2}$$

$$\phi_{,t}^{n+1} = \text{Something}(\phi_{,t}^n, \phi_{,t}^{n-1}) \left(\frac{\Delta t}{\Delta x} = \lambda \right)$$

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:31 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 28 (29 of 83) Fit Page Width

Thumbnails

24

25

26

27

28

FDA for 1-D Wave Equation

- Discretized Boundary conditions

$$\phi_1^{n+1} = \phi_J^{n+1} = 0$$
- Discretized Initial conditions
 - Need to specify two "time levels" of data (effectively $\phi(x, 0)$ and $\phi_t(x, 0)$), i.e. we must specify

$$\phi_j^0, \quad j = 1, 2, \dots, J$$

$$\phi_j^1, \quad j = 1, 2, \dots, J$$

ensuring that the initial values are compatible with the boundary conditions.

- Can solve (16) explicitly for ϕ_j^{n+1} :

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:32 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 29 (30 of 83) Fit Page Width

FDA for 1-D Wave Equation

- Also note that (17) is actually a *linear system* for the unknowns ϕ_j^{n+1} , $j = 1, 2, \dots, J$; in combination with the discrete boundary conditions can write

$$\mathbf{A} \phi^{n+1} = \mathbf{b} \quad (18)$$
 where $\mathbf{A}$ is a *diagonal* $J \times J$ matrix and ϕ^{n+1} and $\mathbf{b}$ are vectors of length J .
- Such a difference scheme for an IVP is called an *explicit* scheme.

Thumbnails

25

26

27

28

29

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

Applications Places System | Ubuntu_11.04 - VMware Pla... File Virtual Machine Help 12:33 PM ubuntu-user

Lec1-1.pdf

File Edit View Go Help

Previous Next 29 (30 of 83) Fit Page Width

Thumbnails

- 25
- 26
- 27
- 28
- 29

- Can solve (16) explicitly for ϕ_j^{n+1} :

$$\phi_j^{n+1} = 2\phi_j^n - \phi_j^{n-1} + \lambda^2 (\phi_{j+1}^n - 2\phi_j^n + \phi_{j-1}^n) \quad (17)$$

28

FDA for 1-D Wave Equation

- Also note that (17) is actually a *linear system* for the unknowns ϕ_j^{n+1} , $j = 1, 2, \dots, J$; in combination with the discrete boundary conditions can write

$$\mathbf{A} \phi^{n+1} = \mathbf{b} \quad (18)$$

where $\mathbf{A}$ is a *diagonal* $J \times J$ matrix and ϕ^{n+1} and $\mathbf{b}$ are vectors of length J .

- Such a difference scheme for an IVP is called an *explicit* scheme.

PSI: Explorat... [Update Ma... ubuntu-user... Downloads Lec1.pdf Lec1-1.pdf

