

Title: Looking for Quantum Gravity in the Space of Conformal Field Theories

Date: Jun 13, 2012 02:00 PM


URL: <http://pirsa.org/12060005>

Abstract: Conformal field theories have many applications ranging from continuous phase transitions in Statistical Mechanics to models of beyond the Standard Model physics in Particle Physics.

In this talk, I will explain another remarkable application: some conformal field theories can be used to define and study Quantum Gravity.

I will also try to give a brief summary of some of the main ideas being discussed at the conference "Back to the Bootstrap II".

# Back to the Bootstrap II






# Outline

- Why study Conformal Field Theory?
  - Continuous Phase Transitions
  - Quantum Gravity
- What we are doing
  - c-Theorems
  - Supersymmetry and Integrability
  - Numerical Bootstrap
- What we would like to do

# Continuous Phase Transitions



# Critical Exponents


Correlation length of D<sub>2</sub>O near the critical point

[Sullivan, Neilson, Fischer, Rennie '00]

$$\xi \sim \left( \frac{T - T_c}{T_c} \right)^{-\nu}$$

$$\nu = 0.62 \pm 0.03$$

3d Ising model  
universality class






# Particle Physics

Conformal field theories describe **UV** and **IR fixed points**.

They are the stepping stone for our description of dynamics in the continuum: **Quantum Field Theory**.



Could the TeV scale of **Particle Physics** require a strongly coupled CFT description?

Watch the PI colloquium of Slava Rychkov to know more about the idea of **Conformal Technicolor**.

[Luty, Okui '04]

[Rattazzi, Rychkov, Tonni, Vichi '08]

# Quantum Gravity

General Relativity is a low energy effective field theory that requires **UV completion**.

$$G_N = M_{Pl}^{-2} = L_{Pl}^2$$


**Definition:** a quantum theory of gravity is a quantum mechanical theory (unitary time evolution) whose low energy dynamics is well described by **General Relativity**.

In particular, it should have the same low lying energy **spectrum**.

## QG in a box: Anti-de Sitter

Anti-de Sitter spacetime of radius  $R$


$$ds^2 = -\cosh^2\left(\frac{r}{R}\right) dt^2 + dr^2 + R^2 \sinh^2\left(\frac{r}{R}\right) (d\theta^2 + \sin^2\theta d\phi^2)$$






Do we know any quantum  
system with this low-energy  
spectrum?

# Conformal Field Theory



$$\rho = e^{\frac{t}{R}}$$




State-operator map:  $|\mathcal{O}\rangle \leftrightarrow \mathcal{O}$


**Dilatation** operator in  $\mathbb{R}^3$  maps to the  
**Hamiltonian** on  $S^2 \times \mathbb{R}_t$

$$\Delta = ER$$

# Conformal Field Theory



$$\rho = e^{\frac{t}{R}}$$



State-operator map:  $|\mathcal{O}\rangle \leftrightarrow \mathcal{O}$

**Dilatation** operator in  $\mathbb{R}^3$  maps to the  
**Hamiltonian** on  $S^2 \times \mathbb{R}_t$

$$\Delta = ER$$




# CFT Spectrum

States created by insertions of the stress-energy tensor have a **discrete spectrum**

$$\Delta(T_{\mu\nu}) = 3$$

$$\Delta(\partial_{\alpha_1} \dots \partial_{\alpha_n} T_{\mu\nu}) = 3 + n$$




# CFT Spectrum

States created by insertions of the stress-energy tensor have a **discrete spectrum**

$$\Delta(T_{\mu\nu}) = 3$$

$$\Delta(\partial_{\alpha_1} \dots \partial_{\alpha_n} T_{\mu\nu}) = 3 + n$$



# Observables


Correlation functions are natural observables of the CFT

$$\langle T_{\mu_1 \nu_1}(x_1) \dots T_{\mu_n \nu_n}(x_n) \rangle$$

These correspond to graviton “scattering amplitudes” and form a complete set of observables.

No local observables in quantum gravity.

[See Nima's Colloquium]





# Observables


Correlation functions are natural observables of the CFT

$$\langle T_{\mu_1 \nu_1}(x_1) \dots T_{\mu_n \nu_n}(x_n) \rangle$$

These correspond to graviton “scattering amplitudes” and form a complete set of observables.

No local observables in quantum gravity.

[See Nima's Colloquium]



# Observables

Correlation functions are natural observables of the CFT

$$\langle T_{\mu_1 \nu_1}(x_1) \dots T_{\mu_n \nu_n}(x_n) \rangle$$


These correspond to graviton “scattering amplitudes” and form a complete set of observables.

No local observables in quantum gravity.

[See Nima's Colloquium]

Weak coupling implies **factorization**

$$\langle T_{\mu_1 \nu_1}(x_1) \dots T_{\mu_n \nu_n}(x_n) \rangle_c \sim \left( \frac{L_{Pl}}{R} \right)^{n-2}$$



# Quantum Gravity = CFT

Definition: quantum gravity with AdS boundary conditions is a 3-dimensional CFT obeying:

- **Large gap** in its spectrum of dimensions

$$\Delta(\mathcal{O} \neq T_{\mu\nu}) > \Lambda_{UV} R \gg 1$$

- **Factorization**

$$\langle T_{\mu_1\nu_1}(x_1) \dots T_{\mu_n\nu_n}(x_n) \rangle_c \sim (M_{Pl}R)^{2-n}$$

$$M_{Pl}R \geq \Lambda_{UV} R \gg 1$$

# Quantum Gravity = CFT

Definition: quantum gravity with AdS boundary conditions is a 3-dimensional CFT obeying:

- **Large gap** in its spectrum of dimensions

$$\Delta(\mathcal{O} \neq T_{\mu\nu}) > \Lambda_{UV} R \gg 1$$

- **Factorization**

$$\langle T_{\mu_1\nu_1}(x_1) \dots T_{\mu_n\nu_n}(x_n) \rangle_c \sim (M_{Pl}R)^{2-n}$$

$$M_{Pl}R \geq \Lambda_{UV} R \gg 1$$



# Quantum Gravity = CFT

Definition: quantum gravity with AdS boundary conditions is a 3-dimensional CFT obeying:

- **Large gap** in its spectrum of dimensions

$$\Delta(\mathcal{O} \neq T_{\mu\nu}) > \Lambda_{UV} R \gg 1$$

- **Factorization**

$$\langle T_{\mu_1\nu_1}(x_1) \dots T_{\mu_n\nu_n}(x_n) \rangle_c \sim (M_{Pl}R)^{2-n}$$

$$M_{Pl}R \geq \Lambda_{UV} R \gg 1$$

## Can we find such a CFT?

- **Conformal** QFTs in 3 dimensions are not hard to find - there are many examples based on Chern-Simons gauge fields plus charged matter (bosons and/or fermions).

- **Factorization** is easy - it follows from the large  $N$  expansion of  $SU(N)$  gauge theories.

$$T_{\mu\nu} = \frac{1}{N} \text{Tr} (\partial_\mu \Phi \partial_\nu \Phi) + \dots \quad N \sim \frac{R}{L_{Pl}}$$

- **Large gap** in the spectrum of dimensions is hard - this requires strong coupling!

$$\mathcal{O} = \frac{1}{N} \text{Tr} (\Phi^2) \quad \Delta(\mathcal{O}) = 1 + \gamma > \Lambda_{UV} R \gg 1$$

There are supersymmetric examples (ABJM).


↖ anomalous dimension

## QG in flat space

Take the zero curvature limit of AdS:  $R \rightarrow \infty$

**Main difficulty:** local bulk physics is encoded in CFT correlation functions in a non-trivial way.

How to extract the **S-matrix**?





## QG in flat space

Take the zero curvature limit of AdS:  $R \rightarrow \infty$

**Main difficulty:** local bulk physics is encoded in CFT correlation functions in a non-trivial way.


How to extract the **S-matrix**?

**Idea:** prepare wave-packets that scatter in small region of AdS

[Polchinski '99] [Susskind '99] [Gary, Giddings, JP '09]

Best language: **Mellin amplitudes**

[Mack '09]



$$\langle \mathcal{O}_1(x_1) \dots \mathcal{O}_n(x_n) \rangle = \int_{-i\infty}^{i\infty} [d\delta] \boxed{M(\delta_{ij})} \prod_{i < j}^n \Gamma(\delta_{ij}) (x_{ij}^2)^{-\delta_{ij}}$$





## c-Theorems

The space of CFTs is very large and complex.

Weakly coupled CFTs are only a small part of the landscape.

Can we use **Renormalization Group (RG) flows** to organize the landscape?






## c-Theorem in d=2

[Zamolodchikov 86']

In a Lorentz invariant and unitarity 2d QFT, there is a function  $c(g)$  that **decreases** under RG flow.

At RG fixed points (CFT),  $\beta_i(g) = 0$ , this function is **stationary** and equal to the **central charge** of the CFT.

$$c_{IR} < c_{UV}$$




## c-Theorem in d=2

[Zamolodchikov 86']

In a Lorentz invariant and unitarity 2d QFT, there is a function  $c(g)$  that **decreases** under RG flow.

At RG fixed points (CFT),  $\beta_i(g) = 0$ , this function is **stationary** and equal to the **central charge** of the CFT.



$$c_{IR} < c_{UV}$$

$$s = \frac{\pi}{3} c T$$

$$\langle T_{\mu\nu}(x) T_{\alpha\beta}(0) \rangle = \frac{c}{x^4} I_{\mu\nu, \alpha\beta}(x)$$

$$-\frac{c}{6} = \int_{S^2} \langle T_{\mu}^{\mu} \rangle$$

$$T_{\mu}^{\mu} = -\frac{c}{12} \mathcal{R}$$



## c-Theorem in d=4?


$$s = \tilde{c} T^3$$

$$\langle T_{\mu\nu}(x) T_{\alpha\beta}(0) \rangle = \frac{c}{x^4} I_{\mu\nu, \alpha\beta}(x)$$

$$-2a = \int_{S^4} \langle T^\mu_\mu \rangle$$

$$T^\mu_\mu = -a E_4 - c W^2$$

Euler density                      Weyl tensor



# a-Theorem in d=4

[Cardy '88]

[Komargodski, Schwimmer '11]

~~$$s = \frac{c}{6} T^3$$~~


$$-2a = \int_{S^4} \langle T^\mu_\mu \rangle$$

~~$$\langle T_{\mu\nu}(x) T_{\alpha\beta}(0) \rangle = \frac{c}{x^4} I_{\mu\nu,\alpha\beta}(x)$$~~

~~$$T^\mu_\mu = -a E_4 - c W^2$$~~

Two CFTs connected by  
an RG flow must satisfy

$$a_{IR} < a_{UV}$$



## c-Theorem in d=3?

No conformal anomaly

$$T_{\mu}^{\mu} = 0$$

# F-Theorem in d=3

No conformal anomaly

$$T^\mu_\mu = 0$$

Free energy on 3-sphere

$$F = -\log Z_{S^3} |_{\text{finite part}}$$

Entanglement Entropy of a circle



$$S_{EE}(A) = -\text{Tr}(\rho_A \log \rho_A)$$

$$F = S_{EE}(A) |_{\text{finite part}}$$

Two CFTs connected by an RG flow must satisfy

$$F_{IR} < F_{UV}$$

[Myers, Sinha '10]  
[Jafferis, Klebanov, Pufu, Safdi '11]  
[Casini, Huerta '12]





# Supersymmetry and Integrability

# Supersymmetry and Integrability

In **planar N=4 SYM**, we can compute the spectrum of the dilatation operator for all values of the 't Hooft coupling. The current focus is on studying higher point functions.

There is a very large class of (calculable) **Superconformal Field Theories** (N=2), some of which are not continuously connected to free theories. [Gaiotto '09]

**Strategy I:** These special theories are the perfect laboratory to develop new techniques for describing CFTs.

**Strategy II:** These special theories are calculable and closer to interesting theories than free theories.

# Supersymmetry and Integrability

In **planar N=4 SYM**, we can compute the spectrum of the dilatation operator for all values of the 't Hooft coupling. The current focus is on studying higher point functions.


There is a very large class of (calculable) **Superconformal Field Theories** (N=2), some of which are not continuously connected to free theories. [Gaiotto '09]

**Strategy I:** These special theories are the perfect laboratory to develop new techniques for describing CFTs.

**Strategy II:** These special theories are calculable and closer to interesting theories than free theories.




## CFT landscape in $d=4$





# CFT landscape in $d=4$



# Conformal Bootstrap

A CFT is a **consistent** set of correlation functions.

# Conformal Bootstrap

A CFT is a **consistent** set of correlation functions.

Operator Product Expansion (**OPE**)

$$\mathcal{O}_1(x)\mathcal{O}_2(0) = \sum_k \frac{C_{12k}}{x^{\Delta_1+\Delta_2-\Delta_k}} \mathcal{O}_k(0)$$


The **CFT data**  $\{\Delta_i, C_{ijk}\}$  defines all correlations functions by successive use of the OPE. The CFT data is constrained by unitarity and associativity of the OPE.

$$\text{Unitarity} \quad \Rightarrow \quad \begin{aligned} \Delta_i &\geq \frac{d-2}{2} \\ C_{ijk} &\in \mathbb{R} \end{aligned}$$

# Bootstrap Equations

$$\langle \mathcal{O}_1(x_1) \mathcal{O}_2(x_2) \mathcal{O}_3(x_3) \mathcal{O}_4(x_4) \rangle = \sum_k C_{12k} C_{k34}$$

Conformal  
Block





# Bootstrap Equations

$$\sum_k C_{13k} C_{k24} \begin{array}{c} 3 \\ \diagup \\ \text{---} k \text{---} \\ \diagdown \\ 1 \end{array} \begin{array}{c} 4 \\ \diagup \\ \text{---} k \text{---} \\ \diagdown \\ 2 \end{array} = \sum_k C_{12k} C_{k34} \begin{array}{c} \text{Conformal} \\ \text{Block} \\ \downarrow \\ 3 \diagup \text{---} k \text{---} \diagdown 4 \\ | \\ 1 \diagdown \text{---} k \text{---} \diagup 2 \end{array}$$

Associativity of the OPE gives rise to an infinite set of equations for the CFT data  $\{\Delta_i, C_{ijk}\}$

[Polyakov '74]

# Bootstrap Equations


$$\sum_k C_{13k} C_{k24} \begin{array}{c} 3 \\ \diagup \\ \text{---} k \text{---} \\ \diagdown \\ 1 \end{array} \begin{array}{c} 4 \\ \diagdown \\ \text{---} k \text{---} \\ \diagup \\ 2 \end{array} = \sum_k C_{12k} C_{k34} \begin{array}{c} \text{Conformal} \\ \text{Block} \\ \downarrow \\ 3 \diagup \text{---} k \text{---} \diagdown 4 \\ | \\ 1 \diagdown \text{---} k \text{---} \diagup 2 \end{array}$$

Associativity of the OPE gives rise to an infinite set of equations for the CFT data  $\{\Delta_i, C_{ijk}\}$

[Polyakov '74]


# Bounds from Bootstrap equations

[El-Showk, Paulos, Poland, Rychkov,  
Simmons-Duffin, Vichi '12]



# Bounds from Bootstrap equations


[El-Showk, Paulos, Poland, Rychkov,  
Simmons-Duffin, Vichi '12]





# Example: 3d Ising model


[El-Showk, Paulos, Poland, Rychkov, Simmons-Duffin, Vichi '12]



Operator	Spin $l$	$\mathbb{Z}_2$	$\Delta$	Exponent
$\sigma$	0	—	0.5182(3)	$\Delta = 1/2 + \eta/2$
$\sigma'$	0	—	$\gtrsim 4.5$	$\Delta = 3 + \omega_A$
$\varepsilon$	0	+	1.413(1)	$\Delta = 3 - 1/\nu$
$\varepsilon'$	0	+	3.84(4)	$\Delta = 3 + \omega$
$\varepsilon''$	0	+	4.67(11)	$\Delta = 3 + \omega_2$
$T_{\mu\nu}$	2	+	3	n/a
$C_{\mu\nu\kappa\lambda}$	4	+	5.0208(12)	$\Delta = 3 + \omega_{NR}$

# Example: 3d Ising model

[El-Showk, Paulos, Poland, Rychkov, Simmons-Duffin, Vichi '12]



Operator	Spin $l$	$\mathbb{Z}_2$	$\Delta$	Exponent
$\sigma$	0	—	0.5182(3)	$\Delta = 1/2 + \eta/2$
$\sigma'$	0	—	$\gtrsim 4.5$	$\Delta = 3 + \omega_A$
$\varepsilon$	0	+	1.413(1)	$\Delta = 3 - 1/\nu$
$\varepsilon'$	0	+	3.84(4)	$\Delta = 3 + \omega$
$\varepsilon''$	0	+	4.67(11)	$\Delta = 3 + \omega_2$
$T_{\mu\nu}$	2	+	3	n/a
$C_{\mu\nu\kappa\lambda}$	4	+	5.0208(12)	$\Delta = 3 + \omega_{NR}$

What we would like to do

## Problems for the future

- Formulate the numerical bootstrap program as an **algorithm** that finds (all) CFTs consistent with a given set of assumptions
- Interacting CFTs in  **$d > 6$** ?
- Find a CFT that UV completes pure Einstein gravity (or show that it does not exist)



## Problems for the future

- Formulate the numerical bootstrap program as an **algorithm** that finds (all) CFTs consistent with a given set of assumptions
- Interacting CFTs in  **$d > 6$** ?
- Find a CFT that UV completes pure Einstein gravity (or show that it does not exist)

## Problems for the future


- Show that all CFTs that provide UV completions of GR give rise to the exact **area law** of black hole entropy

CFT<sub>d</sub> entropy density:  $s = \tilde{c} T^{d-1}$

$$\langle T_{\mu\nu}(x) T_{\alpha\beta}(0) \rangle = \frac{c}{x^4} I_{\mu\nu, \alpha\beta}(x)$$

General Relativity

$$S = \frac{A}{4G_N} \Rightarrow c = \tilde{c}$$



## Problems for the future


- Show that all CFTs that provide UV completions of GR give rise to the exact **area law** of black hole entropy

CFT<sub>d</sub> entropy density:  $s = \tilde{c} T^{d-1}$

$$\langle T_{\mu\nu}(x) T_{\alpha\beta}(0) \rangle = \frac{c}{x^4} I_{\mu\nu, \alpha\beta}(x)$$

General Relativity

$$S = \frac{A}{4G_N} \Rightarrow c = \tilde{c}$$







***Let's keep mining the  
CFT space!***

