Title: Nonlocality, Entanglement Witnesses and Supra-Correlations

Date: Jun 25, 2012  04:00 PM

URL: http://pirsa.org/12060076

Abstract: While entanglement is believed to underlie the power of quantum computation

and communication, it is not generally well understood for multipartite

systems. Recently, it has been appreciated that there exists proper

no-signaling probability distributions derivable from operators that do not

represent valid quantum states.  Such systems exhibit supra-correlations

that are stronger than allowed by quantum mechanics, but less than the

algebraically allowed maximum in Bell-inequalities (in the bipartite case).

Some of these probability distributions are derivable from an entanglement

witness W, which is a non-positive Hermitian operator constructed such that

its expectation value with a separable quantum state (positive density

matrix) rho_sep is non-negative (so that Tr[W rho]< 0 indicates entanglement

in quantum state rho). In the bipartite case, it is known that by a

modification of the local no-signaling measurements by spacelike separated

parties A and B, the supra-correlations exhibited by any W can be modeled as

derivable from a physically realizable quantum state Ï•. However, this result

does not generalize to the n-partite case for n>2. Supra-correlations can

also be exhibited in 2- and 3-qubit systems by explicitly constructing

"states" O (not necessarily positive quantum states) that exhibit PR

correlations for a fixed, but arbitrary number, of measurements available to

each party. In this paper we examine the structure of "states" that exhibit

supra-correlations. In addition, we examine the affect upon the distribution

of the correlations amongst the parties involved when constraints of

positivity and purity are imposed. We investigate circumstances in which

such "states" do and do not represent valid

quantum states.

# Outline

- Review Bell Inequalities (BI)
- No-Signaling (NS) and Popescu-Rohrlich (PR) supra-quantum correlations (i.e. *stronger* than quantum)
- "States" (operators *O*) reproducing PR correlations
- The Bipartite case: 2-qubits, A and B perform *m* local measurements
  - The form of *O*
  - Some Linear Algebra: existence and uniqueness of solution
  - Numerical investigations (eigenvalues of *O*, correlations, …)
- The Tripartite case: 3-qubits, A,B,C perform *m* local measurements
- Effect of Positivity and Purity constraints on distribution of correlations
- Summary and Conclusion

2

# Bell Inequalities and Correlation Distances

## Bell Inequalities can be viewed as a violation of a classical quadrilateral inequality

Schumacher, PRA 44, 7047 (1991)

Consider a bipartite system $\mathcal{A}\otimes\mathcal{B}$ with measurement directions A,B$\in\mathcal{A}$ and C,D$\in\mathcal{B}$ taking values {± 1}

**Correlations** (expectation values) between A$\in\mathcal{A}$ and C$\in\mathcal{B}$

meas. value (*output*)

meas. setting (*input*)

$$E(AC) \equiv \langle AC \rangle = \sum_{a,c=\{\pm 1\}} a\,c\,P(a,c \mid A,C)$$

$$= P(+,+ \mid A,C) + P(-,- \mid A,C) - P(+,- \mid A,C) - P(-,+ \mid A,C)$$

**Define *Correlation Distance*: Δ(AC)**

$$\Delta(AC) = 1 - E(AC) = P(+,- \mid A,C) + P(-,+ \mid A,C) \geq 0$$

(Using: $\sum_{a,c=\{\pm 1\}} P(a,c \mid A,C) = 1,\ \forall A,C$)

**Then** $\Delta(AC) + \Delta(BC) + \Delta(BD) \geq \Delta(AD)$

$$\Rightarrow S \equiv E(AC) + E(BC) + E(BD) - E(AD) \leq +2$$

**Which the CHSH Bell inequality with bounds**

Classical: $|S_C| \leq 2$; QM: $|S_Q| \leq 2\sqrt{2}$; Algebraic Maximum: $|S_{AM}| \leq 4$




For a bipartite system $\mathcal{A}\otimes\mathcal{B}$ with measurement directions A,B$\in\mathcal{A}$ and C,D$\in\mathcal{B}$

**Classically:** Δ(AC) + Δ(BC) + Δ(BD) ≥ Δ(AD)

**Quantum Mechanically:** this can be violated, e.g. Δ(AC) + Δ(BC) + Δ(BD) < Δ(AD) with singlet state $(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)/\sqrt{2} = (|01\rangle - |10\rangle)/\sqrt{2}$

# Bell Inequalities and Correlation Distances

**Bell Inequalities can be viewed as a violation of a classical quadrilateral inequality**

Schumacher, PRA 44, 7047 (1991)

Consider a bipartite system $\mathcal{A}\otimes\mathcal{B}$ with measurement directions A,B$\in\mathcal{A}$ and C,D$\in\mathcal{B}$ taking values {± 1}

**Correlations** (expectation values) between A$\in\mathcal{A}$ and C$\in\mathcal{B}$

meas. value (*output*)

meas. setting (*input*)

$$E(AC) \equiv \langle AC \rangle = \sum_{a,c=\{\pm 1\}} a\,c\,P(a,c \mid A,C)$$

$$= P(+,+ \mid A,C) + P(-,- \mid A,C) - P(+,- \mid A,C) - P(-,+ \mid A,C)$$

**Define *Correlation Distance*: Δ(AC)**

$$\Delta(AC) = 1 - E(AC) = P(+,- \mid A,C) + P(-,+ \mid A,C) \geq 0$$

(Using: $\sum_{a,c=\{\pm 1\}} P(a,c \mid A,C) = 1,\ \forall A,C$)

**Then** $\Delta(AC) + \Delta(BC) + \Delta(BD) \geq \Delta(AD)$

$$\Rightarrow S \equiv E(AC) + E(BC) + E(BD) - E(AD) \leq +2$$

**Which the CHSH Bell inequality with bounds**

Classical: $|S_C| \leq 2$; QM: $|S_Q| \leq 2\sqrt{2}$; Algebraic Maximum: $|S_{AM}| \leq 4$




For a bipartite system $\mathcal{A}\otimes\mathcal{B}$ with measurement directions A,B$\in\mathcal{A}$ and C,D$\in\mathcal{B}$

**Classically:** Δ(AC) + Δ(BC) + Δ(BD) ≥ Δ(AD)

**Quantum Mechanically:** this can be violated, e.g. Δ(AC) + Δ(BC) + Δ(BD) < Δ(AD) with singlet state $(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)/\sqrt{2} = (|01\rangle - |10\rangle)/\sqrt{2}$

# No Signaling (NS) Theories

**No Signaling Theories represent valid joint probability distributions (non-local) with valid marginal distributions**

**Linear constraints on any joint NS prob. distrib.**

Normalization: $\sum_{a,b=\{0,1\}} P(a,b|x,y) = 1, \ \forall x,y$

No Signaling: $P(a_1,a_2,\ldots,a_k|x_1,\ldots,x_n)$

$= \sum_{a_{k+1},\ldots,a_n \in \{0,1\}} P(a_1,\ldots,a_n|x_1,\ldots,x_n)$

$= P(a_1,a_2,\ldots,a_k|x_1,\ldots,x_k)$

No Signaling Principle: (two inputs/outputs)

$P(a|x,y) \equiv \sum_{b=\{0,1\}} P(a,b|x,y) = P(a|x) \ \forall y$

$P(b|x,y) \equiv \sum_{a=\{0,1\}} P(a,b|x,y) = P(b|y) \ \forall x$

**PR Box:**

$$P(a,b|x,y) = \begin{cases} 1/2 & \text{if } a \oplus b = x \cdot y \\ 0 & \text{otherwise} \end{cases}$$

**Then**

$\sum_{a,b=\{0,1\}} P(a,b|x,y)$

$= \underbrace{P(0,0|x,y) + P(1,1|x,y)}_{a \oplus b = 0} + \underbrace{P(0,1|x,y) + P(1,0|x,y)}_{a \oplus b = 1}$

$= (1/2 + 1/2)\,\delta_{0,x\cdot y} + (1/2 + 1/2)\,\delta_{1,x\cdot y}$

$= \delta_{0,x\cdot y} + \delta_{1,x\cdot y}$

$= 1 \quad \forall x,y$

**Then** $P(a|x,y) \equiv \sum_{b=\{0,1\}} P(a,b|x,y)$

$= \underbrace{P(a,0|x,y)}_{a \oplus b = a \oplus 0 = a} + \underbrace{P(a,1|x,y)}_{a \oplus b = a \oplus 1 \equiv \bar{a}}$

$= 1/2\ \delta_{a,\,x\cdot y} + 1/2\ \delta_{\bar{a},\,x\cdot y}$

$= \begin{cases} 1/2+0 \ (\text{if } a=0 \,\&\, x\cdot y = 0),\ 0+1/2 \ (\text{if } a=0 \,\&\, x\cdot y = 1) \\ 0+1/2 \ (\text{if } a=1 \,\&\, x\cdot y = 0),\ 1/2+0 \ (\text{if } a=1 \,\&\, x\cdot y = 1) \end{cases}$

$= 1/2 \quad \forall a,x,y$

$= 1/2$

$= P(a|x) \ \forall a,x$

(this is *Isotropic*, i.e. $P(a|x) = 1/2$ indep of $x$)

5

# No Signaling (NS) Theories

**No Signaling Theories represent valid joint probability distributions (non-local) with valid marginal distributions**

**Linear constraints on any joint NS prob. distrib.**

$$\text{Normalization: } \sum_{a,b=\{0,1\}} P(a,b \mid x,y) = 1, \ \forall x,y$$

$$\text{No Signaling: } P(a_1,a_2,\ldots,a_k \mid x_1,\ldots,x_n) \equiv \sum_{a_{k+1},\ldots,a_n \in \{0,1\}} P(a_1,\ldots,a_n \mid x_1,\ldots,x_n) = P(a_1,a_2,\ldots,a_k \mid x_1,\ldots,x_k)$$

No Signaling Principle: (two inputs/outputs)

$$P(a \mid x,y) \equiv \sum_{b=\{0,1\}} P(a,b \mid x,y) = P(a \mid x) \ \forall y$$

$$P(b \mid x,y) \equiv \sum_{a=\{0,1\}} P(a,b \mid x,y) = P(b \mid y) \ \forall x$$

**PR Box:**

$$P(a,b \mid x,y) = \begin{cases} 1/2 & \text{if } a \oplus b = x \cdot y \\ 0 & \text{otherwise} \end{cases}$$

**Then**

$$\sum_{a,b=\{0,1\}} P(a,b \mid x,y)$$
$$= \underbrace{P(0,0 \mid x,y) + P(1,1 \mid x,y)}_{a \oplus b = 0} \ \underbrace{+P(0,1 \mid x,y) + P(1,0 \mid x,y)}_{a \oplus b = 1}$$
$$= (1/2 + 1/2)\ \delta_{0,x\cdot y} + (1/2 + 1/2)\ \delta_{1,x\cdot y}$$
$$= \delta_{0,x\cdot y} + \delta_{1,x\cdot y}$$
$$= 1 \quad \forall x,y$$

**Then**

$$P(a \mid x,y) \equiv \sum_{b=\{0,1\}} P(a,b \mid x,y)$$
$$= \underbrace{P(a,0 \mid x,y)}_{a \oplus b = a \oplus 0 = a} + \underbrace{P(a,1 \mid x,y)}_{a \oplus b = a \oplus 1 \equiv \bar{a}}$$
$$= 1/2\ \delta_{a,\,x\cdot y} + 1/2\ \delta_{\bar{a},\,x\cdot y}$$
$$= \begin{cases} 1/2+0 \ (\text{if } a=0 \,\&\, x\cdot y=0),\ 0+1/2 \ (\text{if } a=0 \,\&\, x\cdot y=1) \\ 0+1/2 \ (\text{if } a=1 \,\&\, x\cdot y=0),\ 1/2+0 \ (\text{if } a=1 \,\&\, x\cdot y=1) \end{cases}$$
$$= 1/2 \quad \forall a,x,y$$
$$= 1/2$$
$$= P(a \mid x) \ \forall a,x$$

(this is *Isotropic*, i.e. $P(a \mid x) = 1/2$ indep of $x$)

# Structure of NS probability distributions

## Consider a set of *n*-spacelike separated measurements

(Acin *et al.*, PRL 104, 140404 (2011)

**No-Signaling (local) measurements**

measurement setting (*input*)

$$M_{\text{non-sig}} = M_{a_1}^{x_1} \otimes \cdots \otimes M_{a_n}^{x_n}$$

measurement value (*output*)

**Probability Distribution from "Trace Rule" for *fixed* set of measurements**

$$P_O \equiv P_O(a_1, \ldots, a_n \mid x_1, \ldots, x_n) = Tr[O\, M_{a_1}^{x_1} \otimes \cdots \otimes M_{a_n}^{x_n}] \geq 0$$

projectors: $M_a^x = \Pi_a^x = |a\rangle_x \langle a|$

**Probability Distribution from "Trace Rule" for *all* measurements**

$$P_W \equiv P(a_1, \ldots, a_N \mid x_1, \ldots, x_n) = Tr[W\, M_{a_1}^{x_1} \otimes \cdots \otimes M_{a_n}^{x_n}] \geq 0$$

(*Gleason correlations*) (W is an *Entanglement Witness*)

$$n = 2\ only: \ P_W(a,b \mid x,y) = Tr[W\, M_a^x \otimes M_b^y] = Tr[\rho_{|\Phi_{BP}\rangle}\, M_a^x \otimes \bar{M}_b^y)],$$

(Barnum *et al.*, PRL 104, 140401 (2011)

**Entanglement Witness**

- A *quantum state* ρ is positive semidefinite (ρ≥0) Hermitian matrix with Tr[ρ]=1
- A *separable state* (classical correlations) has the form $\rho^{sep} = \sum_i p_i \rho_i^{A_1} \otimes \rho_i^{A_2} \otimes \cdots \otimes \rho_i^{A_N}$
- An *entangled state* is a state that cannot be written as separable: $\rho \neq \rho^{sep}$
- An *entanglement witness* W is constructed to be positive on all separable states $\langle a_1, a_2, \ldots | W | a_1, a_2 \ldots \rangle \geq 0$
  i.e. $Tr[\rho^{sep} W] \geq 0$ (follows from using $\rho_i^{A_k} = \sum_j p_{ij}^k |\psi_{ij}^k\rangle\langle\psi_{ij}^k|$ )
- Hence, if Tr[ρ W] ≤ 0, then the quantum state ρ is entangled
- W is generally not a quantum state ρ≥0, since W can have negative eigenvalues

# Current Understanding

1. It is known that for N=2 qubits, one can always write a probability distribution derived from a witness W, from one derived from a valid quantum state using modified measurements. (Barnum, et al. PRL 104, 1040401 (2011))
2. However, this does not generalize to the case of N>2 (Acin, *et al.*, PRL 104, 140404 (2011))

1. Proof: By the Choi-Jamiolkowski isomorphism (CJI), any 2-party witness W can be written as

CJI $\Rightarrow W = (I_A \otimes \Lambda_B)(\rho_{|\Phi_{BP}\rangle})$, where $\Lambda_B$ is positive, trace preserving map,

and $\rho_{|\Phi_{BP}\rangle} = |\Phi_{BP}\rangle\langle\Phi_{BP}|$ is the density matrix for a maximally entangled pure bipartite state.

Then: $P_W(a,b|x,y) = Tr[W\, M_a^x \otimes M_b^y] = Tr[(I \otimes \Lambda)(\rho_{|\Phi_{BP}\rangle})\, M_a^x \otimes M_b^y]$ (acts to right)

$= Tr[M_a^x \otimes M_b^y (I \otimes \Lambda)(\rho_{|\Phi_{BP}\rangle})] = Tr[M_a^x \otimes \Lambda^*(M_b^y)\, \rho_{|\Phi_{BP}\rangle}] = Tr[\rho_{|\Phi_{BP}\rangle}\, M_a^x \otimes M_b'^y)]$, (act to left)

where $M_b'^y = \Lambda^*(M_b^y)$, and $\Lambda^*$ is the dual map to $\Lambda$, i.e Tr[A $\Lambda$(B)]=Tr[$\Lambda^*$(A) B].

2. The above made explicit use of the CJI, in particular $W = (I_A \otimes \Lambda_B)(\rho_{|\Phi_{BP}\rangle})$, which does not extend in general to the multipartite case (N>2).

3. The extension of the CJI holds in the N-party case only for those W with the form

$W = \sum_k p_k \Lambda_{A_1}^x \otimes \ldots \otimes \Lambda_{A_N}^y(\rho_k)$ where $\rho_k$ are N-party quantum states, $p_k$ probs., $\Lambda_{A_i}^x$ pos. maps

$\Rightarrow P_W(a_1,\ldots,a_N|x_1,\ldots,x_N) = Tr[W\, M_{a_1}^{x_1} \otimes \ldots \otimes M_{a_N}^{x_N}] = \sum_k p_k Tr[\rho_k \Lambda_{A_1}^x(M_{a_1}^{x_1}) \otimes \ldots \otimes \Lambda_{A_N}^y(M_{a_N}^{x_N})]$

# Structure of NS probability distributions

Consider a set of *n*-spacelike separated measurements

(Acin *et al.*, PRL 104, 140404 (2011)

measurement setting (*input*)

No-Signaling (local) measurements

$$M_{\text{non-sig}} = M_{a_1}^{x_1} \otimes \cdots \otimes M_{a_n}^{x_n}$$

measurement value (*output*)

Probability Distribution from "Trace Rule" for *fixed* set of measurements

$$P_O \equiv P_O(a_1,\ldots,a_n \mid x_1,\ldots,x_n) = Tr[O\, M_{a_1}^{x_1} \otimes \cdots \otimes M_{a_n}^{x_n}] \geq 0$$

projectors: $M_a^x = \Pi_a^x = |a\rangle_x \langle a|$

Probability Distribution from "Trace Rule" for *all* measurements

$$P_W \equiv P(a_1,\ldots,a_N \mid x_1,\ldots,x_n) = Tr[W\, M_{a_1}^{x_1} \otimes \cdots \otimes M_{a_n}^{x_n}] \geq 0$$

(*Gleason correlations*) (W is an *Entanglement Witness*)

Entanglement Witness

$$n = 2 \text{ only}: \; P_W(a,b \mid x,y) = Tr[W\, M_a^x \otimes M_b^y] = Tr[\rho_{|\Phi_{BP}\rangle}\, M_a^x \otimes \bar{M}_b^y)],$$

(Barnum *et al.*, PRL 104, 140401 (2011)

- A *quantum state* ρ is positive semidefinite (ρ≥0) Hermitian matrix with Tr[ρ]=1
- A *separable state* (classical correlations) has the form $\rho^{sep} = \sum_i p_i \rho_i^{A_1} \otimes \rho_i^{A_2} \otimes \cdots \otimes \rho_i^{A_N}$
- An *entangled state* is a state that cannot be written as separable: $\rho \neq \rho^{sep}$
- An *entanglement witness* W is constructed to be positive on all separable states $\langle a_1, a_2, \ldots | W | a_1, a_2 \ldots \rangle \geq 0$ i.e. $Tr[\rho^{sep} W] \geq 0$ (follows from using $\rho_i^{A_k} = \sum_j p_{ij}^k |\psi_{ij}^k\rangle\langle\psi_{ij}^k|$ )
- Hence, if $Tr[\rho W] \leq 0$, then the quantum state ρ is entangled
- W is generally not a quantum state ρ≥0, since W can have negative eigenvalues

# Current Understanding

1. It is known that for N=2 qubits, one can always write a probability distribution derived from a witness W, from one derived from a valid quantum state using modified measurements. (Barnum, et al. PRL 104, 1040401 (2011))
2. However, this does not generalize to the case of N>2 (Acin, *et al.*, PRL 104, 140404 (2011))

1. Proof: By the Choi-Jamiolkowski isomorphism (CJI), any 2-party witness W can be written as

CJI $\Rightarrow W = (I_A \otimes \Lambda_B)(\rho_{|\Phi_{BP}\rangle})$, where $\Lambda_B$ is positive, trace preserving map,

and $\rho_{|\Phi_{BP}\rangle} = |\Phi_{BP}\rangle\langle\Phi_{BP}|$ is the density matrix for a maximally entangled pure bipartite state.

Then: $P_W(a,b|x,y) = Tr[W M_a^x \otimes M_b^y] = Tr[(I \otimes \Lambda)(\rho_{|\Phi_{BP}\rangle}) M_a^x \otimes M_b^y]$ (acts to right)

$= Tr[M_a^x \otimes M_b^y (I \otimes \Lambda)(\rho_{|\Phi_{BP}\rangle})] = Tr[M_a^x \otimes \Lambda^*(M_b^y)\rho_{|\Phi_{BP}\rangle}] = Tr[\rho_{|\Phi_{BP}\rangle} M_a^x \otimes M_b'^y)]$, (act to left)

where $M_b'^y = \Lambda^*(M_b^y)$, and $\Lambda^*$ is the dual map to $\Lambda$, i.e Tr[A $\Lambda$(B)]=Tr[$\Lambda^*$(A) B].

2. The above made explicit use of the CJI, in particular $W = (I_A \otimes \Lambda_B)(\rho_{|\Phi_{BP}\rangle})$, which does not extend in general to the multipartite case (N>2).

3. The extension of the CJI holds in the N-party case only for those W with the form

$W = \sum_k p_k \Lambda_{A_1}^x \otimes \ldots \otimes \Lambda_{A_N}^y(\rho_k)$ where $\rho_k$ are N-party quantum states, $p_k$ probs., $\Lambda_{A_i}^x$ pos. maps

$\Rightarrow P_W(a_1,\ldots,a_N|x_1,\ldots,x_N) = Tr[W M_{a_1}^{x_1} \otimes \ldots \otimes M_{a_N}^{x_N}] = \sum_k p_k Tr[\rho_k \Lambda_{A_1}^x(M_{a_1}^{x_1}) \otimes \ldots \otimes \Lambda_{A_N}^y(M_{a_N}^{x_N})]$

# Bipartite (2-qubits) case: *m*-inputs (x,y), *r* = 2 outputs (a,b)

Let us specialize to the bipartite (Alice-A and Bob-B), 2-qubit case (i.e. $a,b = \{0,1\}$). We let A, B each have *m* possible measurements (i.e. $x,y = \{0,1,\ldots,m-1\}$)

$$P(a,b|x,y) = Tr[O M_a^x \otimes M_b^y] \geq 0$$

Alice and Bob share paired (generalized) PR Boxes with

$$P(a,b|x,y) = 1/2 \quad \text{iff } a \oplus b = x \cdot y \bmod 2, \quad (0 \text{ otherwise})$$




Viewpoint: (i) restrict measurements to NS-type $M_a^x \otimes M_b^y$, (ii) fix probability distribution to $P(a,b|x,y)$, (iii) find $W$.

In general: for each value of x = {0,...,m-1}, define a complete set of projection measurements $\{M_a^x\}$

$$M_a^x = \Pi_a^x = |a\rangle_x \langle a| \qquad \sum_{a=0}^{r-1} M_a^x \equiv I_{r\times r} \qquad M_{a=r-1}^x \equiv I_{r\times r} - \sum_{a=0}^{r-2} M_a^x$$

8

# Bipartite (2-qubits) case: *m*-inputs (x,y), *r* = 2 outputs (a,b)

**For qubits:** $M^x_{a=0} = |0\rangle_x\langle 0| = 1/2(I + \vec{m}_x \cdot \vec{\sigma})$, $\vec{m}_x = m_x(\sin\theta_x\cos\phi_x, \sin\theta_x\cos\phi_x, \cos\theta_x)$

$|\vec{m}_x| \le 1$, (density matrix on Bloch Sphere)

$M^x_{a=1} = |1\rangle_x\langle 1| = 1/2(I - \vec{m}_x \cdot \vec{\sigma})$, $(I = I_{2\times 2}$ identity matrix)

**Solve for dual matrices** $\tilde{M}^x_a$ **that satisfy** $Tr[M^x_a \tilde{M}^{x'}_{a'}] = \delta_{x,x'}\delta_{a,a'}$

**O:**

$$O = \sum_{a,b=0}^{r-1=0}\sum_{x,y=0}^{m-1} P(a,b|x,y)\tilde{M}^x_a \otimes \tilde{M}^y_b + \sum_{a=0}^{r-1=0}\sum_{x=0}^{m-1} P(a|x)\tilde{M}^x_a \otimes \tilde{I} + \sum_{b=0}^{r-1=0}\sum_{y}^{m-1} P(b|y)\tilde{I} \otimes \tilde{M}^y_b + \tilde{I} \otimes \tilde{I},$$

$$= \sum_{x,y=0}^{m-1} P(a=0,b=0|x,y)\tilde{M}^x_0 \otimes \tilde{M}^y_0 + \sum_{x=0}^{m-1} P(a=0|x)\tilde{M}^x_0 \otimes \tilde{I} + \sum_{y=0}^{m-1} P(b=0|y)\tilde{I} \otimes \tilde{M}^y_0 + +\tilde{I} \otimes \tilde{I}$$

**Simplify Notation:** let $\{I, M^x_a; a = 0,\ldots,r-2;\ x = 0,\cdots,m-1\} \equiv \{M_0 \equiv I, \{M_{j\ge 1}\} = \{M_1, M_2, \ldots\}\} = \{M_{\alpha=\{0,j\ge 1\}}\}$

$\{\tilde{M}_{\beta=\{0,j\ge 1\}}\} \equiv \{\tilde{M}_0 \equiv \tilde{I}, \tilde{M}_1, \tilde{M}_2 \ldots\}$, such that $\mathrm{Tr}[M_\alpha \tilde{M}_\beta] = \delta_{\alpha,\beta}$ $\quad \left(\{I, \{M^B_{j\ge 1}\}\} \to \{N_{\beta=\{0,j\ge 1\}}\}\right)$

$$O = \sum_{i,j=0}^{m-1} P^{0,0}_{i,j}\tilde{M}_i \otimes \tilde{N}_j + \sum_{i=0}^{m-1} P^{0,\bullet}_{i}\tilde{M}_i \otimes \tilde{I} + \sum_{j=0}^{m-1} P^{\bullet,0}_{j}\tilde{I} \otimes \tilde{N}_j + \tilde{I} \otimes \tilde{I}$$

9

# Bipartite (2-qubits) case: *m*-inputs (x,y), *r* = 2 outputs (a,b)

**Solution for qubits: (for A; with M,m→N,n for B)**

$$M_0 = I = I_{2\times 2}$$

$$M_i = 1/2(I + \vec{m}_i \cdot \vec{\sigma}),\ |\vec{m}_i| \le 1$$

$$\tilde{M}_0 \equiv \tilde{I} = 1/2(I + \sum_i \vec{\tilde{m}}_i \cdot \vec{\sigma}) \equiv 1/2(I + \vec{\tilde{m}} \cdot \vec{\sigma});$$

$$\tilde{M}_i = \vec{\tilde{m}}_i \cdot \vec{\sigma}$$

$$\vec{m}_i \cdot \vec{\tilde{m}}_j = \delta_{i,j}$$

$$|\vec{m}_i| \le 1,\ |\vec{\tilde{m}}_i| \ge 1$$

**In general**

$$O = \frac{1}{4}[\sum_{i,j=0}^{m-1} (4P_{i,j}^{0,0} - 2(P_i^{0,\bullet} + P_j^{\bullet,0}) + 1)(\vec{\tilde{m}}_i \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_j \cdot \vec{\sigma}) + \sum_{i=0}^{m-1} (2P_i^{0,\bullet} - 1)(\vec{\tilde{m}}_i \cdot \vec{\sigma}) \otimes I + \sum_{j=0}^{m-1} (2P_j^{\bullet,0} - 1)\, I \otimes (\vec{\tilde{n}}_j \cdot \vec{\sigma}) + I \otimes I\ ]$$

**Using PR-Box with** $P(a,b \mid x=i, y=j) = 1/2\ \delta_{a\oplus b,\, i\cdot j \bmod 2} \Rightarrow P_i^{0,\bullet} = P_j^{\bullet,0} = 1/2\ \forall i,j;\ P_{i,j}^{0,0} = 1/2\,\delta_{0,\, i\cdot j \bmod 2}$

***O*: 2-qubits**

$$O = \frac{1}{4}[\ (\vec{\tilde{m}}_e \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_e \cdot \vec{\sigma}) + (\vec{\tilde{m}}_e \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_o \cdot \vec{\sigma}) + (\vec{\tilde{m}}_o \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_e \cdot \vec{\sigma}) - (\vec{\tilde{m}}_o \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_o \cdot \vec{\sigma}) + I \otimes I\ ]$$

where $\vec{\tilde{m}}_e = \sum_{\text{even}\ i=0,1,2,\ldots} \vec{\tilde{m}}_{2i}$, $\vec{\tilde{m}}_o = \sum_{\text{odd}\ i=0,1,2,\ldots} \vec{\tilde{m}}_{2i+1}$, $\vec{\tilde{n}}_e = \sum_{\text{even}\ j=0,1,2,\ldots} \vec{\tilde{n}}_{2j}$, $\vec{\tilde{n}}_o = \sum_{\text{odd}\ j=0,1,2,\ldots} \vec{\tilde{n}}_{2j+1}$

# Bipartite (2-qubits) case: *m*-inputs (x,y), *r* = 2 outputs (a,b)

**Solution for qubits: (for A; with M,m→N,n for B)**

$$M_0 = I = I_{2\times 2}$$

$$M_i = 1/2(I + \vec{m}_i \cdot \vec{\sigma}),\ |\vec{m}_i| \le 1$$

$$\tilde{M}_0 \equiv \tilde{I} = 1/2(I + \sum_i \vec{\tilde{m}}_i \cdot \vec{\sigma}) \equiv 1/2(I + \vec{\tilde{m}} \cdot \vec{\sigma});$$

$$\tilde{M}_i = \vec{\tilde{m}}_i \cdot \vec{\sigma}$$

$$\vec{m}_i \cdot \vec{\tilde{m}}_j = \delta_{i,j}$$

$$|\vec{m}_i| \le 1,\ |\vec{\tilde{m}}_i| \ge 1$$

**In general**

$$O = \frac{1}{4}[\sum_{i,j=0}^{m-1} (4P_{i,j}^{0,0} - 2(P_i^{0,\bullet} + P_j^{\bullet,0}) + 1)(\vec{\tilde{m}}_i \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_j \cdot \vec{\sigma}) + \sum_{i=0}^{m-1} (2P_i^{0,\bullet} - 1)(\vec{\tilde{m}}_i \cdot \vec{\sigma}) \otimes I + \sum_{j=0}^{m-1} (2P_j^{\bullet,0} - 1)\, I \otimes (\vec{\tilde{n}}_j \cdot \vec{\sigma}) + I \otimes I\ ]$$

**Using PR-Box with** $P(a,b|x=i,y=j) = 1/2\ \delta_{a\oplus b,\, i\cdot j \bmod 2} \Rightarrow P_i^{0,\bullet} = P_j^{\bullet,0} = 1/2\ \forall i,j;\ P_{i,j}^{0,0} = 1/2\,\delta_{0,\, i\cdot j \bmod 2}$

***O*: 2-qubits**

$$O = \frac{1}{4}[(\vec{\tilde{m}}_e \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_e \cdot \vec{\sigma}) + (\vec{\tilde{m}}_e \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_o \cdot \vec{\sigma}) + (\vec{\tilde{m}}_e \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_e \cdot \vec{\sigma}) - (\vec{\tilde{m}}_o \cdot \vec{\sigma}) \otimes (\vec{\tilde{n}}_o \cdot \vec{\sigma}) + I \otimes I\ ]$$

where $\vec{\tilde{m}}_e = \sum_{\text{even}\ i=0,1,2,\ldots} \vec{\tilde{m}}_{2i}$, $\vec{\tilde{m}}_o = \sum_{\text{odd}\ i=0,1,2,\ldots} \vec{\tilde{m}}_{2i+1}$, $\vec{\tilde{n}}_e = \sum_{\text{even}\ j=0,1,2,\ldots} \vec{\tilde{n}}_{2j}$, $\vec{\tilde{n}}_o = \sum_{\text{odd}\ j=0,1,2,\ldots} \vec{\tilde{n}}_{2j+1}$

# Bipartite (2-qubits) case: *m*-inputs (x,y), *r* = 2 outputs (a,b)

**For qubits:** $M^x_{a=0} = |0\rangle_x\langle 0| = 1/2(I + \vec{m}_x \cdot \vec{\sigma})$, $\vec{m}_x = m_x(\sin\theta_x\cos\phi_x, \sin\theta_x\cos\phi_x, \cos\theta_x)$

$|\vec{m}_x| \le 1$, (density matrix on Bloch Sphere)

$M^x_{a=1} = |1\rangle_x\langle 1| = 1/2(I - \vec{m}_x \cdot \vec{\sigma})$, $(I = I_{2\times 2}$ identity matrix)

Solve for dual matrices $\tilde{M}^x_a$ that satisfy $Tr[M^x_a \tilde{M}^{x'}_{a'}] = \delta_{x,x'}\delta_{a,a'}$

**O:**

$$O = \sum_{a,b=0}^{r-1=0}\sum_{x,y=0}^{m-1} P(a,b|x,y)\tilde{M}^x_a \otimes \tilde{M}^y_b + \sum_{a=0}^{r-1=0}\sum_{x=0}^{m-1} P(a|x)\tilde{M}^x_a \otimes \tilde{I} + \sum_{b=0}^{r-1=0}\sum_{y}^{m-1} P(b|y)\tilde{I} \otimes \tilde{M}^y_b + \tilde{I} \otimes \tilde{I},$$

$$= \sum_{x,y=0}^{m-1} P(a=0,b=0|x,y)\tilde{M}^x_0 \otimes \tilde{M}^y_0 + \sum_{x=0}^{m-1} P(a=0|x)\tilde{M}^x_0 \otimes \tilde{I} + \sum_{y=0}^{m-1} P(b=0|y)\tilde{I} \otimes \tilde{M}^y_0 + + \tilde{I} \otimes \tilde{I}$$

**Simplify Notation:** let $\{I, M^x_a; a = 0,\ldots,r-2;\ x = 0,\cdots,m-1\} \equiv \{M_0 \equiv I, \{M_{j\ge 1}\} = \{M_1, M_2, \ldots\}\} = \{M_{a=\{0,j\ge 1\}}\}$

$\{\tilde{M}_{\beta=\{0,j\ge 1\}}\} \equiv \{\tilde{M}_0 \equiv \tilde{I}, \tilde{M}_1, \tilde{M}_2 \ldots\}$, such that $\mathrm{Tr}[M_\alpha \tilde{M}_\beta] = \delta_{\alpha,\beta}$ $\quad \left(\{I, \{M^B_{j\ge 1}\}\} \to \{N_{\beta=\{0,j\ge 1\}}\}\right)$

$$O = \sum_{i,j=0}^{m-1} P^{0,0}_{i,j}\tilde{M}_i \otimes \tilde{N}_j + \sum_{i=0}^{m-1} P^{0,\bullet}_i \tilde{M}_i \otimes \tilde{I} + \sum_{j=0}^{m-1} P^{\bullet,0}_j \tilde{I} \otimes \tilde{N}_j + \tilde{I} \otimes \tilde{I}$$

9

# Bipartite (2-qubits) case: *m*-inputs (x,y), *r* = 2 outputs (a,b)

**For qubits:** $\boxed{M^x_{a=0} = |0\rangle_x\langle 0| = 1/2(I + \vec{m}_x \cdot \vec{\sigma})}$, $\vec{m}_x = m_x(\sin\theta_x\cos\phi_x, \sin\theta_x\cos\phi_x, \cos\theta_x)$

$|\vec{m}_x| \le 1$, (density matrix on Bloch Sphere)

$M^x_{a=1} = |1\rangle_x\langle 1| = 1/2(I - \vec{m}_x \cdot \vec{\sigma})$, $(I = I_{2\times 2}$ identity matrix)

**Solve for dual matrices** $\tilde{M}^x_a$ **that satisfy** $\boxed{Tr[M^x_a \tilde{M}^{x'}_{a'}] = \delta_{x,x'}\delta_{a,a'}}$

**O:**

$$\boxed{\begin{aligned} O &= \sum_{a,b=0}^{r-1=0}\sum_{x,y=0}^{m-1} P(a,b|x,y)\tilde{M}^x_a \otimes \tilde{M}^y_b + \sum_{a=0}^{r-1=0}\sum_{x=0}^{m-1} P(a|x)\tilde{M}^x_a \otimes \tilde{I} + \sum_{b=0}^{r-1=0}\sum_{y}^{m-1} P(b|y)\tilde{I} \otimes \tilde{M}^y_b + \tilde{I} \otimes \tilde{I}, \\ &= \sum_{x,y=0}^{m-1} P(a=0,b=0|x,y)\tilde{M}^x_0 \otimes \tilde{M}^y_0 + \sum_{x=0}^{m-1} P(a=0|x)\tilde{M}^x_0 \otimes \tilde{I} + \sum_{y=0}^{m-1} P(b=0|y)\tilde{I} \otimes \tilde{M}^y_0 + +\tilde{I} \otimes \tilde{I} \end{aligned}}$$

**Simplify Notation:** let $\{I, M^x_a; a = 0,\ldots,r-2;\ x = 0,\cdots,m-1\} \equiv \{M_0 \equiv I, \{M_{j\ge 1}\} = \{M_1, M_2, \ldots\}\} = \{M_{\alpha=\{0,j\ge 1\}}\}$

$\{\tilde{M}_{\beta=\{0,j\ge 1\}}\} \equiv \{\tilde{M}_0 = \tilde{I}, \tilde{M}_1, \tilde{M}_2 \ldots\}$, such that $\mathrm{Tr}[M_\alpha \tilde{M}_\beta] = \delta_{\alpha,\beta}$ $\quad \left(\{I, \{M^B_{j\ge 1}\}\} \to \{N_{\beta=\{0,j\ge 1\}}\}\right)$

$$\boxed{O = \sum_{i,j=0}^{m-1} P^{0,0}_{i,j}\tilde{M}_i \otimes \tilde{N}_j + \sum_{i=0}^{m-1} P^{0,\bullet}_i \tilde{M}_i \otimes \tilde{I} + \sum_{j=0}^{m-1} P^{\bullet,0}_j \tilde{I} \otimes \tilde{N}_j + \tilde{I} \otimes \tilde{I}}$$

9

# Bipartite (2-qubits) case: Properties
## *m*-inputs (x,y), *r* = 2 outputs (a,b)

$$O=\frac{1}{4}[\,(\vec{\tilde{m}}_e\cdot\vec{\sigma})\otimes(\vec{\tilde{n}}_e\cdot\vec{\sigma})+(\vec{\tilde{m}}_e\cdot\vec{\sigma})\otimes(\vec{\tilde{n}}_o\cdot\vec{\sigma})+(\vec{\tilde{m}}_o\cdot\vec{\sigma})\otimes(\vec{\tilde{n}}_e\cdot\vec{\sigma})-(\vec{\tilde{m}}_o\cdot\vec{\sigma})\otimes(\vec{\tilde{n}}_o\cdot\vec{\sigma})+I\otimes I\,]$$

where $\vec{\tilde{m}}_e=\sum_{\text{even}\ i=0,1,2,\ldots}\vec{\tilde{m}}_{2i}$, $\vec{\tilde{m}}_o=\sum_{\text{odd}\ i=0,1,2,\ldots}\vec{\tilde{m}}_{2i+1}$, $\vec{\tilde{n}}_e=\sum_{\text{even}\ j=0,1,2,\ldots}\vec{\tilde{n}}_{2j}$, $\vec{\tilde{n}}_o=\sum_{\text{odd}\ j=0,1,2,\ldots}\vec{\tilde{n}}_{2j+1}$




**m=3**

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ |
|---|---|---|---|
| $\vec{m}_0$ | 1/2 | 1/2 | 1/2 |
| $\vec{m}_1$ | 1/2 | 0 | 1/2 |
| $\vec{m}_2$ | 1/2 | 1/2 | 1/2 |

$P(a=0,b=0\,|\,x=\vec{m}_i,y=\vec{n}_j)$,

$a\oplus b=0,\ P=1/2$

$(x,y)\in\{(e,e),(e,o),(o,e)\}$.

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ |
|---|---|---|---|
| $\vec{m}_0$ | 0 | 0 | 0 |
| $\vec{m}_1$ | 0 | −1/2 | 0 |
| $\vec{m}_2$ | 0 | 0 | 0 |

$P(a=0,b=1\,|\,\vec{m}_i,\vec{n}_j)$,

$a\oplus b=1,P=1/2$

$(x,y)\in\{(o,o)\}$.

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ |
|---|---|---|---|
| $\vec{m}_0$ | 1 | 1 | 1 |
| $\vec{m}_1$ | 1 | −1 | 1 |
| $\vec{m}_2$ | 1 | 1 | 1 |

$E(\vec{m}_i,\vec{n}_j)$

correlations

**m=4**

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ | $\vec{n}_3$ |
|---|---|---|---|---|
| $\vec{m}_0$ | 0.237 | 0.395 | 0.072 | 0.406 |
| $\vec{m}_1$ | 0.162 | 0.018 | 0.381 | 0.004 |
| $\vec{m}_2$ | 0.469 | 0.449 | 0.341 | 0.457 |
| $\vec{m}_3$ | 0.249 | 0.038 | 0.481 | 0.024 |

$P(a=0,b=0\,|\,x=\vec{m}_i,y=\vec{n}_j)$,

probabilities

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ | $\vec{n}_3$ |
|---|---|---|---|---|
| $\vec{m}_0$ | −0.052 | 0.581 | −0.710 | 0.623 |
| $\vec{m}_1$ | −0.350 | −0.927 | 0.522 | −0.982 |
| $\vec{m}_2$ | 0.875 | 0.796 | 0.365 | 0.829 |
| $\vec{m}_3$ | −0.004 | −0.847 | 0.923 | −0.905 |

$E(\vec{m}_i,\vec{n}_j)$

correlations

12

# Bipartite (2-qubits) case: Properties
## *m*-inputs (x,y), *r* = 2 outputs (a,b)

$$O=\frac{1}{4}[\ (\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})+(\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})+(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})-(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})+I\otimes I\ ]$$

where $\underset{\text{even}}{\vec{m}_e}=\sum_{i=0,1,2,\ldots}\vec{m}_{2i}$, $\underset{\text{odd}}{\vec{m}_o}=\sum_{i=0,1,2,\ldots}\vec{m}_{2i+1}$, $\underset{\text{even}}{\vec{n}_e}=\sum_{j=0,1,2,\ldots}\vec{n}_{2j}$, $\underset{\text{odd}}{\vec{n}_o}=\sum_{j=0,1,2,\ldots}\vec{n}_{2j+1}$




**m=3**

$$\begin{array}{c|ccc} & \vec{n}_0 & \vec{n}_1 & \vec{n}_2 \\ \vec{m}_0 & 1/2 & 1/2 & 1/2 \\ \vec{m}_1 & 1/2 & 0 & 1/2 \\ \vec{m}_2 & 1/2 & 1/2 & 1/2 \end{array}$$

$P(a=0,b=0\,|x=\vec{m}_i, y=\vec{n}_j)$,
$a\oplus b=0,\ P=1/2$
$(x,y)\in\{(e,e),(e,o),(o,e)\}$.

$$\begin{array}{c|ccc} & \vec{n}_0 & \vec{n}_1 & \vec{n}_2 \\ \vec{m}_0 & 0 & 0 & 0 \\ \vec{m}_1 & 0 & -1/2 & 0 \\ \vec{m}_2 & 0 & 0 & 0 \end{array}$$

$P(a=0,b=1\,|\vec{m}_i,\vec{n}_j)$,
$a\oplus b=1, P=1/2$
$(x,y)\in\{(o,o)\}$.

$$\begin{array}{c|ccc} & \vec{n}_0 & \vec{n}_1 & \vec{n}_2 \\ \vec{m}_0 & 1 & 1 & 1 \\ \vec{m}_1 & 1 & -1 & 1 \\ \vec{m}_2 & 1 & 1 & 1 \end{array}$$

$E(\vec{m}_i,\vec{n}_j)$
correlations

**m=4**

$$\begin{array}{c|cccc} & \vec{n}_0 & \vec{n}_1 & \vec{n}_2 & \vec{n}_3 \\ \vec{m}_0 & 0.237 & 0.395 & 0.072 & 0.406 \\ \vec{m}_1 & 0.162 & 0.018 & 0.381 & 0.004 \\ \vec{m}_2 & 0.469 & 0.449 & 0.341 & 0.457 \\ \vec{m}_3 & 0.249 & 0.038 & 0.481 & 0.024 \end{array}$$

$P(a=0,b=0\,|\,x=\vec{m}_i, y=\vec{n}_j)$.
probabilities

$$\begin{array}{c|cccc} & \vec{n}_0 & \vec{n}_1 & \vec{n}_2 & \vec{n}_3 \\ \vec{m}_0 & -0.052 & 0.581 & -0.710 & 0.623 \\ \vec{m}_1 & -0.350 & -0.927 & 0.522 & -0.982 \\ \vec{m}_2 & 0.875 & 0.796 & 0.365 & 0.829 \\ \vec{m}_3 & -0.004 & -0.847 & 0.923 & -0.905 \end{array}$$

$E(\vec{m}_i,\vec{n}_j)$
correlations

12

# Bipartite (2-qubits) case: Properties
## *m*-inputs (x,y), *r* = 2 outputs (a,b)

$$O=\frac{1}{4}[\,(\vec{\tilde{m}}_e\cdot\vec{\sigma})\otimes(\vec{\tilde{n}}_e\cdot\vec{\sigma})+(\vec{\tilde{m}}_e\cdot\vec{\sigma})\otimes(\vec{\tilde{n}}_o\cdot\vec{\sigma})+(\vec{\tilde{m}}_o\cdot\vec{\sigma})\otimes(\vec{\tilde{n}}_e\cdot\vec{\sigma})-(\vec{\tilde{m}}_o\cdot\vec{\sigma})\otimes(\vec{\tilde{n}}_o\cdot\vec{\sigma})+I\otimes I\,]$$

where $\vec{\tilde{m}}_e=\sum_{\text{even}\ i=0,1,2,\ldots}\vec{\tilde{m}}_{2i}$, $\vec{\tilde{m}}_o=\sum_{\text{odd}\ i=0,1,2,\ldots}\vec{\tilde{m}}_{2i+1}$, $\vec{\tilde{n}}_e=\sum_{\text{even}\ j=0,1,2,\ldots}\vec{\tilde{n}}_{2j}$, $\vec{\tilde{n}}_o=\sum_{\text{odd}\ j=0,1,2,\ldots}\vec{\tilde{n}}_{2j+1}$






**m=3**

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ |
|---|---|---|---|
| $\vec{m}_0$ | 1/2 | 1/2 | 1/2 |
| $\vec{m}_1$ | 1/2 | 0 | 1/2 |
| $\vec{m}_2$ | 1/2 | 1/2 | 1/2 |

$P(a=0,b=0\,|\,x=\vec{m}_i,y=\vec{n}_j)$,
$a\oplus b=0,\ P=1/2$
$(x,y)\in\{(e,e),(e,o),(o,e)\}$,

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ |
|---|---|---|---|
| $\vec{m}_0$ | 0 | 0 | 0 |
| $\vec{m}_1$ | 0 | −1/2 | 0 |
| $\vec{m}_2$ | 0 | 0 | 0 |

$P(a=0,b=1\,|\,\vec{m}_i,\vec{n}_j)$,
$a\oplus b=1,P=1/2$
$(x,y)\in\{(o,o)\}$.

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ |
|---|---|---|---|
| $\vec{m}_0$ | 1 | 1 | 1 |
| $\vec{m}_1$ | 1 | −1 | 1 |
| $\vec{m}_2$ | 1 | 1 | 1 |

$E(\vec{m}_i,\vec{n}_j)$
correlations

**m=4**

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ | $\vec{n}_3$ |
|---|---|---|---|---|
| $\vec{m}_0$ | 0.237 | 0.395 | 0.072 | 0.406 |
| $\vec{m}_1$ | 0.162 | 0.018 | 0.381 | 0.004 |
| $\vec{m}_2$ | 0.469 | 0.449 | 0.041 | 0.457 |
| $\vec{m}_3$ | 0.249 | 0.038 | 0.481 | 0.024 |

$P(a=0,b=0\,|\,x=\vec{m}_i,y=\vec{n}_j)$,
probabilities

| | $\vec{n}_0$ | $\vec{n}_1$ | $\vec{n}_2$ | $\vec{n}_3$ |
|---|---|---|---|---|
| $\vec{m}_0$ | −0.052 | 0.581 | −0.710 | 0.623 |
| $\vec{m}_1$ | −0.350 | −0.927 | 0.522 | −0.982 |
| $\vec{m}_2$ | 0.875 | 0.796 | 0.365 | 0.829 |
| $\vec{m}_3$ | −0.004 | −0.847 | 0.923 | −0.905 |

$E(\vec{m}_i,\vec{n}_j)$
correlations

12

# Bipartite (2-qubits) case: *m*-inputs (*x*,*y*), *r* = 2 outputs (*a*,*b*)

**PR Box state is of the form:**

$$O_{PR}=\frac{1}{4}\left[I\otimes I+\sum_{\alpha,\beta=1}^{3}C_{\alpha\beta}^{PR}\,\sigma_\alpha\otimes\sigma_\beta\right],\qquad C_{\alpha\beta}^{PR}=\begin{array}{c}\\ e\\ o\\ e\end{array}\begin{array}{c}\begin{array}{ccc}e & o & e\end{array}\\ \begin{bmatrix}1 & 1 & 1\\ 1 & -1 & 1\\ 1 & 1 & 1\end{bmatrix}\end{array},\qquad \Rightarrow E(\vec{m},\vec{n})=\sum_{\alpha,\beta=1}^{3}m_\alpha C_{\alpha\beta}^{PR}\otimes n_\beta\equiv\vec{m}\cdot\mathbf{C}^{PR}\cdot\vec{n}$$

$$P(a,b\,|\,x,y)=1/2\quad \text{if } a\oplus b=x\cdot y \bmod 2$$

**Most general 2-qubit state:**

max. mixed; Alice Bloch vector; Bob Bloch vector; 2-party Alice-Bob correlations

$$O^{(2)}=\frac{1}{4}\left[I\otimes I+(\vec{s}^{(1)}\cdot\vec{\sigma})\otimes I+I\otimes(\vec{s}^{(2)}\cdot\vec{\sigma})+\sum_{\alpha,\beta=1}^{3}C_{\alpha\beta}\,\sigma_\alpha\otimes\sigma_\beta\right]$$

---

**Quantum Singlet (Pure) State (Bell)** $\quad |\psi_{\text{singlet}}\rangle=(|01\rangle-|10\rangle)/\sqrt{2}$

$$\rho^{\text{singlet}}=\frac{1}{4}\left[I\otimes I+\sum_{\alpha,\beta=1}^{3}C_{\alpha\beta}^{\text{singlet}}\,\sigma_\alpha\otimes\sigma_\beta\right],\qquad C_{\alpha\beta}^{\text{singlet}}=-\delta_{\alpha\beta}=\begin{bmatrix}-1 & 0 & 0\\ 0 & -1 & 0\\ 0 & 0 & -1\end{bmatrix},\qquad \Rightarrow E^{\text{singlet}}(\vec{m},\vec{n})=-\vec{m}\cdot\vec{n}$$

**Bell states** $\quad C_{\alpha\beta}^{\text{Bell States}}=\left\{\begin{bmatrix}-1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 1\end{bmatrix},\begin{bmatrix}1 & 0 & 0\\ 0 & -1 & 0\\ 0 & 0 & 1\end{bmatrix},\begin{bmatrix}1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & -1\end{bmatrix},\begin{bmatrix}-1 & 0 & 0\\ 0 & -1 & 0\\ 0 & 0 & -1\end{bmatrix}\right\}$

**What determines the structure of** $\vec{s}^{(1)}$, $\vec{s}^{(2)}$, $C_{\alpha\beta}$ ?

13

# Bipartite (2-qubits) case: *m*-inputs (*x*,*y*), *r* = 2 outputs (*a*,*b*)

**PR Box state is of the form:**

$$O_{PR}=\frac{1}{4}\left[I\otimes I+\sum_{\alpha,\beta=1}^{3}C_{\alpha\beta}^{PR}\,\sigma_\alpha\otimes\sigma_\beta\right],\qquad C_{\alpha\beta}^{PR}=\begin{array}{c}\\ e\\ o\\ e\end{array}\begin{array}{c}\begin{array}{ccc}e & o & e\end{array}\\ \begin{bmatrix}1 & 1 & 1\\ 1 & -1 & 1\\ 1 & 1 & 1\end{bmatrix}\end{array},\qquad \Rightarrow E(\vec{m},\vec{n})=\sum_{\alpha,\beta=1}^{3}m_\alpha C_{\alpha\beta}^{PR}\otimes n_\beta\equiv\vec{m}\cdot\mathbf{C}^{PR}\cdot\vec{n}$$

$$P(a,b\,|\,x,y)=1/2\quad \text{if } a\oplus b=x\cdot y \bmod 2$$

**Most general 2-qubit state:**

max. mixed; Alice Bloch vector; Bob Bloch vector; 2-party Alice-Bob correlations

$$O^{(2)}=\frac{1}{4}\left[I\otimes I+(\vec{s}^{(1)}\cdot\vec{\sigma})\otimes I+I\otimes(\vec{s}^{(2)}\cdot\vec{\sigma})+\sum_{\alpha,\beta=1}^{3}C_{\alpha\beta}\,\sigma_\alpha\otimes\sigma_\beta\right]$$

---

**Quantum Singlet (Pure) State (Bell)** $\quad |\psi_{\text{singlet}}\rangle=(|01\rangle-|10\rangle)/\sqrt{2}$

$$\rho^{\text{singlet}}=\frac{1}{4}\left[I\otimes I+\sum_{\alpha,\beta=1}^{3}C_{\alpha\beta}^{\text{singlet}}\,\sigma_\alpha\otimes\sigma_\beta\right],\qquad C_{\alpha\beta}^{\text{singlet}}=-\delta_{\alpha\beta}=\begin{bmatrix}-1 & 0 & 0\\ 0 & -1 & 0\\ 0 & 0 & -1\end{bmatrix},\qquad \Rightarrow E^{\text{singlet}}(\vec{m},\vec{n})=-\vec{m}\cdot\vec{n}$$

**Bell states** $C_{\alpha\beta}^{\text{Bell States}}=\left\{\begin{bmatrix}-1&0&0\\0&1&0\\0&0&1\end{bmatrix},\begin{bmatrix}1&0&0\\0&-1&0\\0&0&1\end{bmatrix},\begin{bmatrix}1&0&0\\0&1&0\\0&0&-1\end{bmatrix},\begin{bmatrix}-1&0&0\\0&-1&0\\0&0&-1\end{bmatrix}\right\}$

**What determines the structure of** $\vec{s}^{(1)}$, $\vec{s}^{(2)}$, $C_{\alpha\beta}$ ?

13

# No Signaling (NS) Theories
# Extension to Tripartite case: 3-qubits

- No Signaling Theories can be extended to tripartite systems of 3-qubits, each with *m*-inputs (measurements) to measure pure tripartite entanglement.
- The generalization of the bipartite CHSH inequality was given by Svetlichny

Svetlichny, PRD 35, 3066 (1987)

First, consider correlations E(a,b,c|x,y,z) between 3 observers A,B,C with

inputs: $x, y, z \in \{0,1\}$, and outputs: $a, b, c \in \{0,1\}$

The relevant inequality to compute is the *Svetlichny inequality* (SI)

$$S \equiv | E(a,b,c|0,0,0) + E(a,b,c|0,1,0) + E(a,b,c|1,0,0) - E(a,b,c|1,1,0) + E(a,b,c|0,0,1) - E(a,b,c|0,1,1) - E(a,b,c|1,0,1) - E(a,b,c|1,1,1) |$$

The bounds on the Svetlichny inequality are

Classical: $|S_C| \le 4$; QM: $|S_Q| \le 4\sqrt{2}$; Algebraic Maximum: $|S_{AM}| \le 8$

A Tripartite No Signaling (TNS) Box yielding the algebraic maximum of S is given by probabilities

TPR Box:

$$P(a,b,c|x,y,z) = \begin{cases} 1/4 & \text{if } a \oplus b \oplus c = x \cdot y \oplus y \cdot z \oplus x \cdot z \\ 0 & \text{otherwise} \end{cases}$$

with measurement settings $x, y, z$ and outcomes $a, b, c$ as *bits*,

i.e $a, b, c, x, y, z \in \{0,1\}$ (Note: $\{0,1\} \leftrightarrow \{+1,-1\}$)

With a TPR Box:

$S = 1 + 1 + 1 - (-1) + 1 - (-1) - (-1) - (-1) = 8$

14

# Tripartite (3-qubits) case: *m*-inputs (*x*,*y*), *r* = 2 outputs (*a*,*b*)

We generalize this to *m* measurements settings (inputs), still with binary outputs

**TPR Box:**

$$P(a,b,c\,|\,x,y,z)=\begin{cases}1/4 & \text{if } a\oplus b\oplus c = x\cdot y\oplus y\cdot z\oplus x\cdot z\\ 0 & \text{otherwise}\end{cases}$$

with $m$ measurement settings $x, y, z$ and binary outcomes $a, b, c$, (i.e. qubits)

$\Rightarrow a,b,c,\in\{0,1\}, x,y,z\in\{0,1,\ldots,m-1\}$

and find the 3-qubit entanglement witness exhibiting TPR correlations:

$$O_{TPR}=\frac{1}{8}[I\otimes I\otimes I+$$

$O_{PR}$ 2-qubit term

$$+\{(\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})+(\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})+(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})-(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})\}\otimes(\vec{r}_e\cdot\vec{\sigma})$$

$$-\{(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})+(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})+(\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})-(\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})\}\otimes(\vec{r}_o\cdot\vec{\sigma})]$$

where $\underset{\text{even}}{\vec{q}_e}=\sum_{i=0,1,2,\ldots}\vec{q}_{2i}$, $\underset{\text{odd}}{\vec{q}_o}=\sum_{i=0,1,2,\ldots}\vec{q}_{2i+1}$, $\vec{q}=\{\vec{m},\vec{n},\vec{r}\}$

15

# Tripartite (3-qubits) case: *m*-inputs (*x*,*y*), *r* = 2 outputs (*a*,*b*)

We generalize this to *m* measurements settings (inputs), still with binary outputs

**TPR Box:**

$$P(a,b,c\,|\,x,y,z)=\begin{cases}1/4 & \text{if } a\oplus b\oplus c = x\cdot y\oplus y\cdot z\oplus x\cdot z\\ 0 & \text{otherwise}\end{cases}$$

with $m$ measurement settings $x, y, z$ and binary outcomes $a, b, c$, (i.e. qubits)

$\Rightarrow a,b,c,\in\{0,1\}, x, y, z\in\{0,1,\ldots,m-1\}$

and find the 3-qubit entanglement witness exhibiting TPR correlations:

$$O_{TPR}=\frac{1}{8}[I\otimes I\otimes I+$$

$O_{PR}$ 2-qubit term

$$+\{(\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})+(\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})+(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})-(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})\}\otimes(\vec{r}_e\cdot\vec{\sigma})$$

$$-\{(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})+(\vec{m}_o\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})+(\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_o\cdot\vec{\sigma})-(\vec{m}_e\cdot\vec{\sigma})\otimes(\vec{n}_e\cdot\vec{\sigma})\}\otimes(\vec{r}_o\cdot\vec{\sigma})]$$

where $\vec{q}_e=\sum_{i=0,1,2,\ldots}\vec{q}_{2i}$ (even), $\vec{q}_o=\sum_{i=0,1,2,\ldots}\vec{q}_{2i+1}$ (odd), $\vec{q}=\{\vec{m},\vec{n},\vec{r}\}$

15

# Positivity Constraints

**Traceless part of state** $O^{(n)} = \frac{1}{N}[I_N - K]$ Define: $K \equiv [I_N - NO^{(n)}]$. If $O^{(n)} \geq 0 \Rightarrow K \leq 1$.

Characteristic Polynomial $\chi_A(\lambda) = \prod_{i=1}^{N}(\lambda - \lambda_i)$, $\{\lambda_{i=1,\ldots,N} \leq 1\}$ Cayley-Hamilton Theorem $\chi_A(K) = 0$




$$\chi_A(\lambda) = \prod_{i=1}^{N}(\lambda - \lambda_i)\Big|_{\lambda \geq 1} \geq 0 \Rightarrow \frac{d^k \chi(\lambda)}{d\lambda^k}\Big|_{\lambda \geq 1} \geq 0 \Rightarrow \frac{d^k \chi(\lambda)}{d\lambda^k}\Big|_{\lambda = 1} \geq 0$$

General: Characteristic Polynomial in terms of Traces of Matrix A: $T_m \equiv Tr[A^m]$

$$\chi_A(\lambda) = \lambda^N + D_1\lambda^{N-1} + \ldots + D_{N-1}\lambda + D_N$$
$$\Rightarrow mD_m + D_{m-1}T_1 + D_{m-2}T_2 + \ldots + D_1T_{m-1} + T_m = 0,\ m = 1,2,\ldots,N.$$

$$\sum_{r=2}^{k}\binom{N-r}{N-k}(-D_r) \leq \binom{N}{k} = \frac{N!}{k!(N-k)!},$$

$$T_2 \leq N(N-1),$$

$$(N-2)(T_2/2) + (T_3/3) \leq \binom{N}{3},$$

$$\frac{(N-2)(N-3)}{2}(T_2/2) + (N-3)(T_3/3) + 1/4(T_4 - 1/2T_2^2) \leq \binom{N}{4}$$

**Results:**
**2 & 3 Qubits**

$$T_2 = Tr[K^2] = N\left[\sum_{i=1}^{n}\left(\vec{s}^{(i)}\right)^2 + \sum_{i<j}^{n}\sum_{\alpha_i,\alpha_j=1}^{3}\left(C^{(i,j)}_{\alpha_i\alpha_j}\right)^2 + \ldots + \sum_{\alpha_1,\alpha_2,\ldots,\alpha_n=1}^{3}\left(C^{(1,2,\ldots,n)}_{\alpha_1\alpha_2,\ldots,\alpha_n}\right)^2\right] \leq N(N-1)$$

$$\Rightarrow n = 2:\ \vec{s}^{2(1)} + \vec{s}^{2(2)} + \sum_{\alpha_1,\alpha_2=1}^{3} C^2_{\alpha_1\alpha_2} \leq (N-1)_{N=2^2} = 3,$$

$$\Rightarrow n = 3:\ \vec{s}^{2(1)} + \vec{s}^{2(2)} + \vec{s}^{2(3)} + \sum_{\alpha_1,\alpha_2=1}^{3} A^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2=1}^{3} B^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2=1}^{3} C^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2,\alpha_3=1}^{3} C^2_{\alpha_1\alpha_2\alpha_3} \leq (N-1)_{N=2^3} = 7,$$

16

# Positivity Constraints

**Traceless part of state** $O^{(n)} = \frac{1}{N}[I_N - K]$ Define: $K \equiv [I_N - NO^{(n)}]$. If $O^{(n)} \geq 0 \Rightarrow K \leq 1$.

Characteristic Polynomial $\chi_A(\lambda) = \prod_{i=1}^{N}(\lambda - \lambda_i)$, $\{\lambda_{i=1,\ldots,N} \leq 1\}$ Cayley-Hamilton Theorem $\chi_A(K) = 0$




$$\chi_A(\lambda) = \prod_{i=1}^{N}(\lambda - \lambda_i)\Big|_{\lambda \geq 1} \geq 0 \Rightarrow \frac{d^k \chi(\lambda)}{d\lambda^k}\Big|_{\lambda \geq 1} \geq 0 \Rightarrow \frac{d^k \chi(\lambda)}{d\lambda^k}\Big|_{\lambda = 1} \geq 0$$

General: Characteristic Polynomial in terms of Traces of Matrix A: $T_m \equiv Tr[A^m]$

$$\chi_A(\lambda) = \lambda^N + D_1\lambda^{N-1} + \ldots + D_{N-1}\lambda + D_N$$
$$\Rightarrow mD_m + D_{m-1}T_1 + D_{m-2}T_2 + \ldots + D_1T_{m-1} + T_m = 0,\ m = 1, 2, \ldots, N.$$

$$\sum_{r=2}^{k}\binom{N-r}{N-k}(-D_r) \leq \binom{N}{k} = \frac{N!}{k!(N-k)!},$$

$$T_2 \leq N(N-1),$$

$$(N-2)(T_2/2) + (T_3/3) \leq \binom{N}{3},$$

$$\frac{(N-2)(N-3)}{2}(T_2/2) + (N-3)(T_3/3) + 1/4(T_4 - 1/2T_2^2) \leq \binom{N}{4}$$

**Results:**
**2 & 3 Qubits**

$$T_2 = Tr[K^2] = N\left[\sum_{i=1}^{n}\left(\vec{s}^{(i)}\right)^2 + \sum_{i<j}^{n}\sum_{\alpha_i,\alpha_j=1}^{3}\left(C^{(i,j)}_{\alpha_i\alpha_j}\right)^2 + \ldots + \sum_{\alpha_1,\alpha_2,\ldots,\alpha_n=1}^{3}\left(C^{(1,2,\ldots,n)}_{\alpha_1,\alpha_2,\ldots,\alpha_n}\right)^2\right] \leq N(N-1)$$

$$\Rightarrow n = 2:\ \vec{s}^{2(1)} + \vec{s}^{2(2)} + \sum_{\alpha_1,\alpha_2=1}^{3} C^2_{\alpha_1\alpha_2} \leq (N-1)_{N=2^2} = 3,$$

$$\Rightarrow n = 3:\ \vec{s}^{2(1)} + \vec{s}^{2(2)} + \vec{s}^{2(3)} + \sum_{\alpha_1,\alpha_2=1}^{3} A^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2=1}^{3} B^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2=1}^{3} C^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2,\alpha_3=1}^{3} C^2_{\alpha_1\alpha_2\alpha_3} \leq (N-1)_{N=2^3} = 7,$$

16

# Positivity Constraints

**Traceless part of state** $O^{(n)} = \frac{1}{N}[I_N - K]$ Define: $K \equiv [I_N - NO^{(n)}]$. If $O^{(n)} \geq 0 \Rightarrow K \leq 1$.

Characteristic Polynomial $\chi_A(\lambda) = \prod_{i=1}^{N}(\lambda - \lambda_i)$, $\{\lambda_{i=1,\ldots,N} \leq 1\}$ Cayley-Hamilton Theorem $\chi_A(K) = 0$




$$\chi_A(\lambda) = \prod_{i=1}^{N}(\lambda - \lambda_i)\Big|_{\lambda \geq 1} \geq 0 \Rightarrow \frac{d^k \chi(\lambda)}{d\lambda^k}\Big|_{\lambda \geq 1} \geq 0 \Rightarrow \frac{d^k \chi(\lambda)}{d\lambda^k}\Big|_{\lambda = 1} \geq 0$$

General: Characteristic Polynomial in terms of Traces of Matrix A: $T_m \equiv Tr[A^m]$

$$\chi_A(\lambda) = \lambda^N + D_1\lambda^{N-1} + \ldots + D_{N-1}\lambda + D_N$$
$$\Rightarrow mD_m + D_{m-1}T_1 + D_{m-2}T_2 + \ldots + D_1T_{m-1} + T_m = 0,\ m = 1, 2, \ldots, N.$$

$$\sum_{r=2}^{k}\binom{N-r}{N-k}(-D_r) \leq \binom{N}{k} = \frac{N!}{k!(N-k)!},$$

$$T_2 \leq N(N-1),$$

$$(N-2)(T_2/2) + (T_3/3) \leq \binom{N}{3},$$

$$\frac{(N-2)(N-3)}{2}(T_2/2) + (N-3)(T_3/3) + 1/4(T_4 - 1/2T_2^2) \leq \binom{N}{4}$$

**Results:**
**2 & 3 Qubits**

$$T_2 = Tr[K^2] = N\left[\sum_{i=1}^{n}\left(\vec{s}^{(i)}\right)^2 + \sum_{i<j}^{n}\sum_{a_i,a_j=1}^{3}\left(C^{(i,j)}_{a_i a_j}\right)^2 + \ldots + \sum_{a_1,a_2,\ldots,a_n=1}^{3}\left(C^{(1,2,\ldots,n)}_{a_1,a_2,\ldots,a_n}\right)^2\right] \leq N(N-1)$$

$$\Rightarrow n = 2:\ \vec{s}^{2(1)} + \vec{s}^{2(2)} + \sum_{a_1,a_2=1}^{3} C^2_{a_1 a_2} \leq (N-1)_{N=2^2} = 3,$$

$$\Rightarrow n = 3:\ \vec{s}^{2(1)} + \vec{s}^{2(2)} + \vec{s}^{2(3)} + \sum_{a_1,a_2=1}^{3} A^2_{a_1 a_2} + \sum_{a_1,a_2=1}^{3} B^2_{a_1 a_2} + \sum_{a_1,a_2=1}^{3} C^2_{a_1 a_2} + \sum_{a_1,a_2,a_3=1}^{3} C^2_{a_1 a_2 a_3} \leq (N-1)_{N=2^3} = 7,$$

# Positivity Constraints

**3-Qubit states**

$$O^{(3)} = \frac{1}{8}[I\otimes I\otimes I + (\vec{s}^{(1)}\cdot\vec{\sigma})\otimes I\otimes I + I\otimes(\vec{s}^{(2)}\cdot\vec{\sigma})\otimes I + I\otimes I\otimes(\vec{s}^{(3)}\cdot\vec{\sigma}) + \sum_{\alpha,\beta=1}^{3} A_{\alpha\beta}\,\sigma_\alpha\otimes\sigma_\beta\otimes I + \sum_{\alpha,\beta=1}^{3} B_{\alpha\beta}\,\sigma_\alpha\otimes I\otimes\sigma_\beta + \sum_{\alpha,\beta=1}^{3} C_{\alpha\beta}\,I\otimes\sigma_\alpha\otimes\sigma_\beta + \sum_{\alpha,\beta,\gamma=1}^{3} C_{\alpha\beta\gamma}\,\sigma_\alpha\otimes\sigma_\beta\otimes\sigma_\gamma].$$

**$T_2$ Positivity Constraint**

$$T_2 = Tr[K^2];\ n=3:\ \vec{s}^{2(1)} + \vec{s}^{2(2)} + \vec{s}^{2(3)} + \sum_{\alpha_1,\alpha_2=1}^{3} A^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2=1}^{3} B^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2=1}^{3} C^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2,\alpha_3=1}^{3} C^2_{\alpha_1\alpha_2\alpha_3} \le (N-1)_{N=2^3} = 7$$

**Tripartite PR Box States are of the form:**

$$O_{TPR} = \frac{1}{8}\left[I\otimes I\otimes I + \sum_{\alpha,\beta,\gamma=1}^{3} C^{TPR}_{\alpha\beta\gamma}\,\sigma_\alpha\otimes\sigma_\beta\otimes\sigma_\gamma\right],$$

Example: $C^{TPR;\varepsilon}_{\alpha\beta\gamma} \equiv \varepsilon_{\alpha\beta\gamma}\ (\varepsilon_{123}=1)$

(i) NOT a quantum state

(ii) NOT a pure state

But odd *n*-qubit states of solely $n=2n'+1$-party correlations have $T_{2n'+1} \equiv 0$ ($T_3 \equiv 0$); Consider $D_3$ constraint:

$$(N-2)(T_2/2) + (T_3/3) \le \binom{N}{3} \Rightarrow \boxed{T_2 \le N(N-1)/3}$$

$$\Rightarrow \sum_{\alpha,\beta,\gamma=1}^{3}\left(C_{\alpha\beta\gamma}\right)^2 \le 7/3,\ \text{but}\ \sum_{\alpha,\beta,\gamma=1}^{3}\left(C^{TPR;\varepsilon}_{\alpha\beta\gamma}\right)^2 = 6 > 7/3$$

$\varepsilon_{123} = \varepsilon_{231} = \varepsilon_{312} = 1$

$\varepsilon_{213} = \varepsilon_{321} = \varepsilon_{132} = -1$

$\boxed{T_2 = 6 < 7}$

# Positivity Constraints

**3-Qubit states**

$$O^{(3)} = \frac{1}{8}[I\otimes I\otimes I + (\vec{s}^{(1)}\cdot\vec{\sigma})\otimes I\otimes I + I\otimes(\vec{s}^{(2)}\cdot\vec{\sigma})\otimes I + I\otimes I\otimes(\vec{s}^{(3)}\cdot\vec{\sigma}) + \sum_{\alpha,\beta=1}^{3} A_{\alpha\beta}\,\sigma_\alpha\otimes\sigma_\beta\otimes I + \sum_{\alpha,\beta=1}^{3} B_{\alpha\beta}\,\sigma_\alpha\otimes I\otimes\sigma_\beta + \sum_{\alpha,\beta=1}^{3} C_{\alpha\beta}\,I\otimes\sigma_\alpha\otimes\sigma_\beta + \sum_{\alpha,\beta,\gamma=1}^{3} C_{\alpha\beta\gamma}\,\sigma_\alpha\otimes\sigma_\beta\otimes\sigma_\gamma].$$

**$T_2$ Positivity Constraint**

$$T_2 = Tr[K^2];\ n=3:\ \vec{s}^{2(1)} + \vec{s}^{2(2)} + \vec{s}^{2(3)} + \sum_{\alpha_1,\alpha_2=1}^{3} A^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2=1}^{3} B^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2=1}^{3} C^2_{\alpha_1\alpha_2} + \sum_{\alpha_1,\alpha_2,\alpha_3=1}^{3} C^2_{\alpha_1\alpha_2\alpha_3} \le (N-1)_{N=2^3} = 7$$

**Tripartite PR Box States are of the form:**

$$O_{TPR} = \frac{1}{8}\left[I\otimes I\otimes I + \sum_{\alpha,\beta,\gamma=1}^{3} C^{TPR}_{\alpha\beta\gamma}\,\sigma_\alpha\otimes\sigma_\beta\otimes\sigma_\gamma\right],$$

Example: $C^{TPR;\varepsilon}_{\alpha\beta\gamma} \equiv \varepsilon_{\alpha\beta\gamma}\ (\varepsilon_{123}=1)$

(i) NOT a quantum state

(ii) NOT a pure state

But odd $n$-qubit states of solely $n=2n'+1$-party correlations have $T_{2n'+1} \equiv 0$ ($T_3 \equiv 0$); Consider $D_3$ constraint:

$$(N-2)(T_2/2) + (T_3/3) \le \binom{N}{3} \Rightarrow \boxed{T_2 \le N(N-1)/3}$$

$$\Rightarrow \sum_{\alpha,\beta,\gamma=1}^{3}\left(C_{\alpha\beta\gamma}\right)^2 \le 7/3,\ \text{but}\ \sum_{\alpha,\beta,\gamma=1}^{3}\left(C^{TPR;\varepsilon}_{\alpha\beta\gamma}\right)^2 = 6 > 7/3$$

$\varepsilon_{123} = \varepsilon_{231} = \varepsilon_{312} = 1$

$\varepsilon_{213} = \varepsilon_{321} = \varepsilon_{132} = -1$

$\boxed{T_2 = 6 < 7}$

# Purity Constraints

**Pure State**

$$O_{pure}^{(n)} = |\Psi_N\rangle\langle\Psi_N| \quad O_{pure}^{2(n)} = O_{pure}^{(n)} \quad O_{pure}^{(n)}(O_{pure}^{(n)} - I_N) = 0. \quad (N = 2^n)$$

**Pure State Constraint**

$$O^{(n)} = 1/N[I_N - K]: \; O_{pure}^{(2)}(O_{pure}^{(2)} - I_N) = 0 \;\Rightarrow\; K^2 = (N-1)I_N - (N-2)K$$

$$\Rightarrow T_{m+2} = (N-1)T_m - (N-2)T_{m+1}: \quad \boxed{T_3 = -N(N-1)(N-2),} \quad T_4 = N(N-1)^2 + N(N-1)(N-2)^2,$$

**An Immediate Consequence**

States of solely odd n=2n'+1-party correlations cannot be pure due to $T_{2n'+1} \equiv 0$ ($T_3 \equiv 0$)

$$O^{(2n'+1)} = \frac{1}{N}[I \otimes I \otimes \cdots \otimes I + \sum_{\alpha_1,\alpha_2,\ldots,\alpha_{2n'+1}=1}^{3} C_{\alpha_1,\alpha_2,\ldots,\alpha_{2n'+1}}^{(1,2,\ldots,n)} \sigma_{\alpha_1} \otimes \sigma_{\alpha_2} \otimes \cdots \otimes \sigma_{\alpha_{2n'+1}}]$$

**Tripartite PR Box States are not pure states:**

$$O_{TPR} = \frac{1}{8}\left[I \otimes I \otimes I + \sum_{\alpha,\beta,\gamma=1}^{3} C_{\alpha\beta\gamma}^{TPR} \sigma_\alpha \otimes \sigma_\beta \otimes \sigma_\gamma\right],$$

# Purity Constraints: 2-Qubits

**Most general 2-qubit state:**

max. mixed; Alice Bloch vector; Bob Bloch vector; 2-party Alice-Bob correlations

$$O^{(2)} = \frac{1}{4}\left[ I \otimes I + (\vec{s}^{(1)} \cdot \vec{\sigma}) \otimes I + I \otimes (\vec{s}^{(2)} \cdot \vec{\sigma}) + \sum_{\alpha,\beta=1}^{3} C_{\alpha\beta}\, \sigma_\alpha \otimes \sigma_\beta \right]$$

**Pure State** $O^{(n)} = 1/N[I_N - K]:\ O^{(2)}_{pure}(O^{(2)}_{pure} - I_N) = 0 \ \Rightarrow\ K^2 = (N-1)I_N - (N-2)K$

**2-Qubit Pure State**

$$\rho^{(2)}_{pure} = 1/4[I \otimes I + p\,\sigma_1 \otimes I + p\, I \otimes \sigma_1 - \sigma_1 \otimes \sigma_1 - q\,\sigma_2 \otimes \sigma_2 - \sigma_3 \otimes \sigma_3] \geq 0$$

$0 \leq p \leq 1$ ($p = 0$ : max ent. Bell States), $\quad q = Tr[\rho^{(2)^*}_{pure}\rho^{(2)}_{pure}] = \sqrt{1-p^2} \geq 0$ (concurrence)

**Purity Constraint**

$$\vec{s}^{(1)} = \mathbf{C} \cdot \vec{s}^{(2)}, \qquad \vec{s}^{(2)} = \vec{s}^{(1)} \cdot \mathbf{C},$$

$$C_{\alpha\beta} = s^{(1)}_\alpha s^{(2)}_\beta - C^{(sub)}_{\alpha\beta} \qquad C^{(sub)}_{\alpha\beta} \equiv 1/2 \sum\nolimits_{\mu\nu\mu'\nu'=1}^{3} C_{\mu\nu} C_{\mu'\nu'} \varepsilon_{\mu\mu'\alpha} \varepsilon_{\nu\nu'\beta}$$

# Purity Constraints: 3-Qubits (cont)

**General 3-Qubit**

$$O^{(3)} = \frac{1}{8}[I\otimes I\otimes I + (\vec{s}^{(1)}\cdot\vec{\sigma})\otimes I\otimes I + I\otimes(\vec{s}^{(2)}\cdot\vec{\sigma})\otimes I + I\otimes I\otimes(\vec{s}^{(3)}\cdot\vec{\sigma}) + \sum_{\alpha,\beta=1}^{3} A_{\alpha\beta}\,\sigma_\alpha\otimes\sigma_\beta\otimes I + \sum_{\alpha,\beta=1}^{3} B_{\alpha\beta}\,\sigma_\alpha\otimes I\otimes\sigma_\beta + \sum_{\alpha,\beta=1}^{3} C_{\alpha\beta}\,I\otimes\sigma_\alpha\otimes\sigma_\beta + \sum_{\alpha,\beta,\gamma=1}^{3} C_{\alpha\beta\gamma}\,\sigma_\alpha\otimes\sigma_\beta\otimes\sigma_\gamma].$$

**notation:** $s_i^2 = \vec{s}^{2(i)}$, $C_{12}^2 = \sum_{\alpha\beta=1}^{3} A_{\alpha\beta}^2 = Tr[A^T A]$, $C_{13}^2 = Tr[B^T B]$, $C_{23}^2 = Tr[C^T C]$, $C_{123}^2 = \sum_{\alpha\beta\gamma=1}^{3} C_{\alpha\beta\gamma}^2$

**Maximal Slice States** $S_{\max}^{(MS)} = 4\sqrt{1+\tau^{(MS)}}$

$$|MS_2\rangle = \frac{1}{\sqrt{2}}\left(|000\rangle + \cos\theta_2\,|101\rangle + \sin\theta_2\,|111\rangle\right), \quad |MS_3\rangle = \frac{1}{\sqrt{2}}\left(|000\rangle + \cos\theta_3\,|110\rangle + \sin\theta_3\,|111\rangle\right)$$

| | | | |
|---|---|---|---|
| | $s_1^2 = s_2^2 = 0,\ s_3^2 = 1-\tau,$ | $C_{12}^2 = 3-2\tau,\ C_{13}^2 = C_{23}^2 = \tau,$ | $C_{123}^2 = 3+\tau,$ |
| Max tangle $\tau = 1$ on GHZ state | $s_1^2 = s_2^2 = s_3^2 = 0,$ | $C_{12}^2 = C_{13}^2 = C_{23}^2 = 1$ | $C_{123}^2 = 4$ |

**Generalized GHZ State** $S_{\max}^{(GGHZ)} = 4\sqrt{1-\tau^{(GGHZ)}}$ for $\tau \le 1/3$ $\quad S_{\max}^{(GGHZ)} = 4\sqrt{2\tau^{(GGHZ)}}$ for $\tau \ge 1/3$.

$$|GGHZ\rangle = \cos\theta_1\,|000\rangle + \sin\theta_1\,|111\rangle, \text{ with } \tau_{MS_2}^2 = \sin^2\theta_2,\ \tau_{MS_3}^2 = \sin^2\theta_3,\ \tau_{GGHZ}^2 = \sin^2 2\theta_1.$$

$s_3^{(1)} = s_3^{(2)} = s_3^{(3)} = \sqrt{1-\tau}$, $s_1^2 + s_2^2 + s_3^2 = 3-3\tau$, $C_{12} = C_{13} = C_{23} = diag(0,0,1)$, $C_{12}^2 + C_{13}^2 + C_{23}^2 = 3$, $C_{123}^2 = 1+3\tau$.

cusp at $\tau = 1/3$ occurs at $s_1^2 + s_2^2 + s_3^2 = C_{123}^2$.


TER
UTE
PHYSICS