


Title: Integral Transforms & Green's Functions - Lecture 3

Date: Aug 29, 2012 09:00 AM

URL: <http://pirsa.org/12080030>

Abstract:

CLASSICAL ELECTRODYNAMA



CLASSICAL ELECTRODYNAMICS

CLASSICAL ELECTRODYNAMICS

MAXWELL EQUATIONS

$$\nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} = 4\pi \mathbf{j}$$

$$\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$$

MAXWELL EQUATIONS

$$\nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} = 4\pi \mathbf{j} \quad (1)$$

$$\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0 \quad (2)$$

$$\nabla \cdot \mathbf{E} = 4\pi \rho \quad (3)$$

$$\nabla \cdot \mathbf{B} = 0 \quad (4)$$

CLASSICAL ELECTRODYNAMICS

a) MAXWELL EQUATIONS

$$\nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} = 4\pi \mathbf{j} \quad (1)$$

$$\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0 \quad (2)$$

$$\nabla \cdot \mathbf{E} = 4\pi \rho \quad (3)$$

$$\nabla \cdot \mathbf{B} = 0 \quad (4)$$

i) CONTING UNKNOWN: E, B

CLASSICAL ELECTRODYNAMICS

a) MAXWELL EQUATIONS

$$\begin{aligned}\nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} &= 4\pi \mathbf{j} & (1) \\ \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0 & (2) \\ \nabla \cdot \mathbf{E} &= 4\pi \rho & (3) \\ \nabla \cdot \mathbf{B} &= 0 & (4)\end{aligned}$$

i) CONTINUOUS UNKNOWN: $\mathbf{E}, \mathbf{B}, \rho, \mathbf{j}$, 8 EQUATIONS
(3) & (4) $\approx$ "INITIAL COND"

CLASSICAL ELECTRODYNAMICS

a) MAXWELL EQUATIONS

$$\begin{aligned}\nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} &= 4\pi \mathbf{j} & (1) \\ \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0 & (2) \\ \nabla \cdot \mathbf{E} &= 4\pi \rho & (3) \\ \nabla \cdot \mathbf{B} &= 0 & (4)\end{aligned}$$

i) CONTINUOUS UNKNOWNS: $\mathbf{E}, \mathbf{B}, \rho, \mathbf{j}$, 8 EQUATIONS
(3) & (4) $\approx$ "INITIAL CONDITIONS"
 $\nabla \cdot (2) \quad \nabla \cdot \nabla \times \mathbf{E} + \frac{\partial}{\partial t} \nabla \cdot \mathbf{B} = 0$

CLASSICAL ELECTRODYNAMICS

a) MAXWELL EQUATIONS

$$\begin{aligned}\nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} &= 4\pi \mathbf{j} & (1) \\ \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0 & (2) \\ \nabla \cdot \mathbf{E} &= 4\pi \rho & (3) \\ \nabla \cdot \mathbf{B} &= 0 & (4)\end{aligned}$$

i) CONTINUOUS UNKNOWNS: $\mathbf{E}, \mathbf{B}, \rho, \mathbf{j}$ 8 EQUATIONS
(3) & (4) $\approx$ "INITIAL CONDITIONS"

$$\nabla \cdot (2) \quad \nabla \cdot \underbrace{\nabla \times \mathbf{E}}_0 + \frac{\partial}{\partial t} \nabla \cdot \mathbf{B} = 0$$

CLASSICAL ELECTRODYNAMICS

a) MAXWELL EQUATIONS

$$\begin{aligned}\nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} &= 4\pi \mathbf{j} \\ \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0 \\ \nabla \cdot \mathbf{E} &= 4\pi \rho \\ \nabla \cdot \mathbf{B} &= 0\end{aligned}$$

(1)

i) CONTINUOUS UNKNOWN: $\mathbf{E}, \mathbf{B}, \rho, \mathbf{j}$ 8 EQUATIONS
(3) & (4) $\approx$ "INITIAL CONDITIONS"

$$\nabla \cdot (\nabla \times \mathbf{E}) + \frac{\partial}{\partial t} \nabla \cdot \mathbf{B} = 0$$

CLASSICAL ELECTRODYNAMICS

a) MAXWELL EQUATIONS

$$\begin{aligned}\nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} &= 4\pi \mathbf{j} & (1) \\ \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0 & (2) \\ \nabla \cdot \mathbf{E} &= 4\pi \rho & (3) \\ \nabla \cdot \mathbf{B} &= 0 & (4)\end{aligned}$$

i) CONTINUOUS UNKNOWN: $\mathbf{E}, \mathbf{B}$ 6, 8 EQUATIONS
(3) & (4) $\approx$ "INITIAL CONDITIONS"

$$\nabla \cdot (2) \quad \nabla \cdot \nabla \times \mathbf{E} + \frac{\partial}{\partial t} \nabla \cdot \mathbf{B} = 0$$

$$\nabla \cdot (1) \Rightarrow -\frac{\partial}{\partial t} \nabla \cdot \mathbf{E} = 4\pi \nabla \cdot \mathbf{j}$$

CLASSICAL ELECTRODYNAMICS

a) MAXWELL EQUATIONS

$$\begin{aligned} \nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} &= 4\pi \mathbf{j} & (1) \\ \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0 & (2) \\ \nabla \cdot \mathbf{E} &= 4\pi \rho & (3) \\ \nabla \cdot \mathbf{B} &= 0 & (4) \end{aligned}$$

i) CONTINUOUS UNKNOWNS: $\mathbf{E}, \mathbf{B}, \rho, \mathbf{j}$ 8 EQUATIONS
(3) & (4) $\approx$ "INITIAL CONDITIONS"

$$\nabla \cdot (2) \quad \nabla \cdot \nabla \times \mathbf{E} + \frac{\partial}{\partial t} \nabla \cdot \mathbf{B} = 0$$

$$\nabla \cdot (1) \Rightarrow -\frac{\partial}{\partial t} \nabla \cdot \mathbf{E} = 4\pi \nabla \cdot \mathbf{j} = -4\pi \frac{\partial \rho}{\partial t}$$

CONTINUITY EQ.

CLASSICAL ELECTRODYNAMICS

a) MAXWELL EQUATIONS

$$\begin{aligned} \nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} &= 4\pi \mathbf{j} & (1) \\ \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0 & (2) \\ \nabla \cdot \mathbf{E} &= 4\pi \rho & (3) \\ \nabla \cdot \mathbf{B} &= 0 & (4) \end{aligned}$$

i) CONTINUOUS UNKNOWNS: $\mathbf{E}, \mathbf{B}, \rho, \mathbf{j}$ 8 EQUATIONS
(3) & (4) $\approx$ "INITIAL CONDITIONS"

$$\nabla \cdot (2) \quad \nabla \cdot \nabla \times \mathbf{E} + \frac{\partial}{\partial t} \nabla \cdot \mathbf{B} = 0$$

$$\nabla \cdot (1) \Rightarrow -\frac{\partial}{\partial t} \nabla \cdot \mathbf{E} = 4\pi \nabla \cdot \mathbf{j} = -4\pi \frac{\partial \rho}{\partial t}$$

$-\frac{\partial \rho}{\partial t}$ CONTINUITY EQ.

MAXWELL EQUATIONS

$$\begin{aligned} \nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} &= 4\pi \mathbf{j} & (1) \\ \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0 & (2) \\ \nabla \cdot \mathbf{E} &= 4\pi \rho & (3) \\ \nabla \cdot \mathbf{B} &= 0 & (4) \end{aligned}$$

i) (CONTINUITY EQUATION)

(3) & (4) $\approx$ "INITIAL CONDITIONS"

$$\nabla \cdot (2) \quad \nabla \cdot \underbrace{\nabla \times \mathbf{E}}_0 + \frac{\partial}{\partial t} \nabla \cdot \mathbf{B} = 0$$

$$\nabla \cdot (1) \Rightarrow -\frac{\partial}{\partial t} \nabla \cdot \mathbf{E} = 4\pi \underbrace{\nabla \cdot \mathbf{j}}_{-\frac{\partial \rho}{\partial t}} = -4\pi \frac{\partial \rho}{\partial t}$$

CONTINUITY EQ

ii) INTERPRETATION - RELATIVITY

$$\text{Diff}(\text{FIELD}) = \text{SOURCE}$$

11) INTERPRETATION — RELATIVITY. $\text{Diff}(\text{FIELD}) = \text{SOURCE}$
EVOLUTION EQ

$$\frac{\partial E}{\partial t} = \dots$$

$$\frac{\partial B}{\partial t} = \dots$$

EVOLUTION EQ

$$G_{\mu\nu} = T_{\mu\nu}$$

$$\frac{\partial E}{\partial t} = \dots$$

$$\frac{\partial B}{\partial t} = \dots$$

III) OTHER FL

EVOLUTION EQ

$$G_{\mu\nu} = T_{\mu\nu}$$

$$\frac{\partial E}{\partial t} = \dots$$

$$\frac{\partial B}{\partial t} = \dots$$

iii) OTHER FORMS.

$$\nabla_\mu F = 4\pi j$$

EVOLUTION EQ

$$\frac{\partial E}{\partial t} = \dots$$
$$\frac{\partial B}{\partial t} = \dots$$

$$G_{\mu\nu} = T_{\mu\nu}$$

iii) OTHER FORMS.

$$\nabla_\mu F^{\mu\nu} = 4\pi j^\nu$$

$$F_{(\mu\nu;\alpha)}{}_{;\alpha} = 0$$

(SEPTEMBER)

iii) OTHER FORMS,

$$\nabla_\mu F^{\mu\nu} = 4\pi j^\nu$$

$$F_{[\mu\nu;\alpha]}_{\text{cycl}} = 0$$

(SEPTEMBER)

$$\begin{aligned} \mathcal{A} * F &= 4\pi j \\ \mathcal{A} F &= 0 \end{aligned}$$

11.11) OTHER FORMS.

$$\nabla_\mu F^{\mu\nu} = 4\pi j^\nu$$
$$F_{(\mu\nu;\alpha\beta)}_{\text{cyc}} = 0$$

(SEPTEMBER)

$\Leftrightarrow$

$$d * F = 4\pi j$$
$$dF = 0$$

(JANUARY)

b) ELECTROMAGNETIC POTENTIALS:

(4)

b) ELECTROMAGNETIC POTENTIALS:

$$(4) \quad \Leftrightarrow \quad B = \nabla \times A$$

(2)

b) ELECTROMAGNETIC POTENTIALS:

$$(4) \quad \Leftrightarrow \quad B = \nabla \times A$$

$$(2) \quad \Leftrightarrow \quad \nabla \times E + \frac{\partial}{\partial t} \nabla \times A = 0$$

$$\nabla \times \left(E + \frac{\partial A}{\partial t} \right) = 0$$

b) ELECTROMAGNETIC POTENTIALS:

$$(4) \quad \Leftrightarrow \quad B = \nabla \times A$$

$$(2) \quad \Leftrightarrow \quad \nabla \times E + \frac{\partial}{\partial t} \nabla \times A = 0$$

$$\nabla \times \left(\underbrace{E + \frac{\partial A}{\partial t}}_{-\nabla \phi} \right) = 0 \quad \Rightarrow$$

$$E = -\nabla \phi - \frac{\partial A}{\partial t}$$

b) ELECTROMAGNETIC POTENTIALS

$$(4) \Leftrightarrow \boxed{B = \nabla \times A}$$

$$\Leftrightarrow \nabla \times E + \frac{\partial}{\partial t} \nabla \times A = 0$$

$$\nabla \times \left(\underbrace{E + \frac{\partial A}{\partial t}}_{-\nabla \phi} \right) = 0 \Rightarrow$$

$$\boxed{E = -\nabla \phi - \frac{\partial A}{\partial t}}$$

↳ ELECTROMAGNETIC POTENTIALS

$$(1) \Leftrightarrow \boxed{B = \nabla \times A}$$

$$(2) \Leftrightarrow \nabla \times E + \frac{\partial}{\partial t} \nabla \times A = 0$$

$$\nabla \times \left(\underbrace{E + \frac{\partial A}{\partial t}}_{-\nabla \phi} \right) = 0 \Rightarrow$$

$$\boxed{E = -\nabla \phi - \frac{\partial A}{\partial t}}$$

$$A^\mu = (\phi, A) \Leftrightarrow E, B$$

5) ELECTROMAGNETIC POTENTIALS

$$(4) \Leftrightarrow \boxed{B = \nabla \times A}$$

$$(2) \Leftrightarrow \nabla \times E + \frac{\partial}{\partial t} \nabla \times A = 0$$

$$\nabla \times \left(\underbrace{E + \frac{\partial A}{\partial t}}_{-\nabla \phi} \right) = 0 \Rightarrow$$

$$\boxed{E = -\nabla \phi - \frac{\partial A}{\partial t}}$$

• GAUGE DEGREE OF FREEDOM

ϕ

$$(2) \Leftrightarrow \nabla \times \mathbf{E} + \frac{\partial}{\partial t} \nabla \times \mathbf{A} = 0$$


$$\nabla \times \left(\mathbf{E} + \frac{\partial \mathbf{A}}{\partial t} \right) = 0 \Rightarrow$$

$$\boxed{\mathbf{E} = -\nabla \phi - \frac{\partial \mathbf{A}}{\partial t}}$$

$$\phi \rightarrow \phi + \frac{\partial \Lambda}{\partial t} \quad \Lambda \text{ ARBITRARY}$$

$$\mathbf{A} \rightarrow \mathbf{A} + \nabla \Lambda$$

$\Rightarrow \mathbf{E}, \mathbf{B}$ DOES NOT CHANGE



CAN IMPOSE $\partial_t \phi + \nabla \cdot \mathbf{A} = 0 \Leftrightarrow \partial_\mu A^\mu = 0$

CAN IMPOSE $\boxed{\partial_t \phi + \nabla \cdot \mathbf{A} = 0}$ $\Leftrightarrow \partial_\mu A^\mu = 0$
LORENTZ GAUGE COND.

CAN IMPOSE $\boxed{\partial_t \phi + \nabla \cdot \mathbf{A} = 0}$ $\Leftrightarrow \partial_\mu A^\mu = 0$
LORENTZ GAUGE COND.

C) WAVE EQ. (4) & (2) SOLVED BY A, ϕ
(3)

CAN IMPOSE $\boxed{\partial_t \phi + \nabla \cdot \mathbf{A} = 0}$ $\Leftrightarrow \partial_\mu A^\mu = 0$
LORENTZ GAUGE COND.

C) WAVE EQ. (4) & (2) SOLVED BY A, ϕ

$$(3) \quad \nabla \cdot \left(-\nabla \phi - \frac{\partial \mathbf{A}}{\partial t} \right) = 4\pi \rho$$

$$-\Delta \phi - \frac{\partial}{\partial t} \underbrace{\nabla \cdot \mathbf{A}} = 4\pi \rho$$

CAN IMPOSE $\boxed{\partial_t \phi + \nabla \cdot \mathbf{A} = 0}$ $\Leftrightarrow \partial_\mu A^\mu = 0$
LORENTZ GAUGE COND.

C) WAVE EQ. (4) & (2) SOLVED BY A, ϕ

$$(3) \quad \nabla \cdot \left(-\nabla \phi - \frac{\partial \mathbf{A}}{\partial t} \right) = 4\pi \rho$$
$$-\Delta \phi - \frac{\partial}{\partial t} \underbrace{\nabla \cdot \mathbf{A}}_{-\partial_t \phi} = 4\pi \rho$$

$$\boxed{\square \phi = -4\pi \rho}$$

CAN IMPOSE $\partial_\mu \phi + \partial_\mu A = 0$ (LORENTZ GAUGE COND.)

LORENTZ GAUGE COND.

C) WAVE EQ. (4) & (2) SOLVED BY A, ϕ


$$(3) \nabla \cdot \left(-\nabla \phi - \frac{\partial \mathbf{A}}{\partial t} \right) = 4\pi \rho$$

$$-\Delta \phi - \underbrace{\frac{\partial}{\partial t} \nabla \cdot \mathbf{A}}_{-\partial_t \phi} = 4\pi \rho$$

$$\square \phi = -4\pi \rho$$

$$\square = -\frac{\partial^2}{\partial t^2} + \Delta$$

D'ALAMBERT OP.



CAN IMPOSE $\partial_\mu \phi + \partial_\mu A_\mu = 0$ (LORENTZ GAUGE COND.)

LORENTZ GAUGE COND.

C) WAVE EQ. (4) & (2) SOLVED BY A_μ, ϕ

$$(3) \nabla \cdot (-\nabla \phi - \frac{\partial \mathbf{A}}{\partial t}) = 4\pi \rho$$


$$-\Delta \phi - \frac{\partial}{\partial t} \underbrace{\nabla \cdot \mathbf{A}}_{-\partial_t \phi} = 4\pi \rho$$

$$\square \phi = -4\pi \rho$$

WAVE EQ.

$$\square = -\frac{\partial^2}{\partial t^2} + \Delta$$

D'ALAMBERT OP.



LORENTZ GAUGE COND.

C) WAVE EQ. (1) & (2) SOLVED BY A, ϕ

$$\begin{aligned} (3) \quad \nabla \cdot \left(-\nabla \phi - \frac{\partial \mathbf{A}}{\partial t} \right) &= 4\pi \rho \\ -\Delta \phi - \frac{\partial}{\partial t} \underbrace{\nabla \cdot \mathbf{A}}_{=0} &= 4\pi \rho \end{aligned}$$

$$\boxed{\square \phi = -4\pi \rho}$$

$$\boxed{\square = -\frac{\partial^2}{\partial t^2} + \Delta}$$

D'ALAMBERT OP.

WAVE EQ.

$$(4) \Leftrightarrow \boxed{\square \mathbf{A} = -4\pi \mathbf{j}}$$

d) GREEN'S FUNCTION :

$$\square G(\vec{R}, T) = \delta(\vec{R}) \delta(t)$$

d) GREEN'S FUNCTION :

$$\square G(\vec{R}, T) = -\delta(\vec{R})\delta(T) \quad (6)$$

$$\square G(\vec{R}, T) = -\delta(\vec{R})\delta(T) \quad (6)$$

$$\frac{1}{(2\pi)^4} \int d\vec{k} d\omega G_{\vec{k}\omega} \square e^{i\vec{k}\cdot\vec{R} - i\omega T} = \frac{1}{(2\pi)^4}$$

(SEPTEMBER)


EVOLUTION EQ

$$\frac{\partial E}{\partial t} = \dots$$

d) GREEN'S FUNCTION :

$$\square G(\vec{R}, T) = -\delta(\vec{R})\delta(T) \quad (6)$$

$$\frac{1}{(2\pi)^4} \int d\vec{k} d\omega \, G_{\vec{k}\omega} \underbrace{\square e^{i\vec{k}\cdot\vec{R} - i\omega T}}_{= -\frac{1}{(2\pi)^4} \int d\vec{k} d\omega}$$



$$\frac{\partial E}{\partial t} =$$

d) GREEN'S FUNCTION.

$$\square G(\vec{R}, T) = -\delta(\vec{R})\delta(T) \quad (6)$$

$$\frac{1}{(2\pi)^4} \int d\vec{k} d\omega \, G_{\vec{k}\omega} \square e^{i\vec{k}\cdot\vec{R} - i\omega T} = -\frac{1}{(2\pi)^4} \int d\vec{k} d\omega \, e^{i\vec{k}\cdot\vec{R} - i\omega T}$$

$$\frac{1}{\omega^2 - \vec{k}^2} e^{i\omega T + (i\vec{k}\cdot\vec{R})} e^{-\epsilon T}$$

$$\frac{\partial E}{\partial t} =$$

d) GREEN'S FUNCTION.

$$\square G(\vec{R}, T) = -\delta(\vec{R})\delta(T) \quad (6)$$

$$\frac{1}{(2\pi)^4} \int d\vec{k} d\omega \, G_{\vec{k}\omega} \square e^{i\vec{k}\cdot\vec{R} - i\omega T} = -\frac{1}{(2\pi)^4} \int d\vec{k} d\omega \, e^{i\vec{k}\cdot\vec{R} - i\omega T}$$

$$\frac{[-(-i\omega)^2 + (i\vec{k})^2]}{[\omega^2 - \vec{k}^2]} e^{i\vec{k}\cdot\vec{R} - i\omega T}$$

(SEPTEMBER)

$$\frac{\partial E}{\partial t} =$$

d) GREEN'S FUNCTION.

$$\square G(\vec{R}, T) = -\delta(\vec{R})\delta(T) \quad (6)$$

$$\frac{1}{(2\pi)^4} \int d\vec{k} d\omega \, G_{\vec{k}\omega} \square e^{i\vec{k}\cdot\vec{R} - i\omega T} = -\frac{1}{(2\pi)^4} \int d\vec{k} d\omega \, e^{i\vec{k}\cdot\vec{R} - i\omega T}$$

$$\frac{[-(-i\omega)^2 + (i\vec{k})^2]}{[\omega^2 - \vec{k}^2]} e^{i\vec{k}\cdot\vec{R} - i\omega T} \Rightarrow G_{\vec{k}\omega} =$$

(SEPTEMBER)

$$\frac{\partial E}{\partial t} =$$

d) GREEN'S FUNCTION.

$$\square G(\vec{R}, T) = -\delta(\vec{R})\delta(T) \quad (6)$$

$$\frac{1}{(2\pi)^4} \int d\vec{k} d\omega \, G_{\vec{k}\omega} \square e^{i\vec{k}\cdot\vec{R} - i\omega T} = -\frac{1}{(2\pi)^4} \int d\vec{k} d\omega \, e^{i\vec{k}\cdot\vec{R} - i\omega T}$$

$$\frac{[-(-i\omega)^2 + (i\vec{k})^2]}{[\omega^2 - \vec{k}^2]} e^{i\vec{k}\cdot\vec{R} - i\omega T} \Rightarrow \boxed{G_{\vec{k}\omega} = \frac{1}{\vec{k}^2 - \omega^2}}$$

(SEPTEMBER)

$$(2\pi)^4 \int d\vec{k} d\omega \underbrace{G_{\vec{k}\omega}}_{\text{Green's function}} \cdot \frac{1}{\omega^2 - \vec{k}^2}$$

$$\frac{[-(-i\omega)^2 + (i\vec{k})^2]}{[\omega^2 - \vec{k}^2]} e^{-i\omega t + i\vec{k} \cdot \vec{r}}$$

$$\Rightarrow G_{\vec{k}\omega} = \frac{1}{\vec{k}^2 - \omega^2}$$

$$G(\vec{R}, T) =$$

$$(2\pi)^4 \int d\vec{k} d\omega \underbrace{G_{\vec{k}\omega}}_{\substack{[-(-i\omega)^2 + (i\vec{k})^2]e \\ [\omega^2 - \vec{k}^2]e}}$$

$$\Rightarrow \boxed{G_{\vec{k}\omega} = \frac{1}{\vec{k}^2 - \omega^2}}$$

$$G(\vec{R}, T) = \frac{1}{(2\pi)^4} \int \frac{1}{\vec{k}^2 - \omega^2} e^{i\vec{k} \cdot \vec{R} - i\omega T} d\vec{k} d\omega$$

$$(2\pi)^4 \int d^3k d\omega \underbrace{G_{\vec{k}\omega}}_{\substack{[-(-i\omega)^2 + (i\vec{k})^2]e \\ [\omega^2 - \vec{k}^2]e}}$$

$$\Rightarrow G_{\vec{k}\omega} = \frac{1}{\vec{k}^2 - \omega^2}$$

$$G(\vec{R}, T) = \frac{1}{(2\pi)^4} \int \frac{1}{\vec{k}^2 - \omega^2} e^{i\vec{k} \cdot \vec{R} - i\omega T} d^3k d\omega$$

$$d^3k = d^3\vec{k} = d^3k$$


(2.11) ...

$$\frac{[-(-i\omega)^2 + (i\vec{k})^2]}{[\omega^2 - \vec{k}^2]} e^{i\vec{k} \cdot \vec{R} - i\omega T}$$

$$\Rightarrow G_{\vec{k}\omega} = \frac{1}{\vec{k}^2 - \omega^2}$$

$$G(\vec{R}, T) = \frac{1}{(2\pi)^4} \int \frac{1}{\vec{k}^2 - \omega^2} e^{i\vec{k} \cdot \vec{R} - i\omega T} d\vec{k} d\omega$$

$$d\vec{k} = d^3k = d(\cos\theta) 2\pi k^2 dk$$



(2.11) ...


$$\frac{[-(-i\omega)^2 + (i\vec{k})^2]}{[\omega^2 - k^2]} e^{i\vec{k} \cdot \vec{R} - i\omega T}$$


$$\Rightarrow G_{\vec{k}\omega} = \frac{1}{k^2 - \omega^2}$$


$$G(\vec{R}, T) = \frac{1}{(2\pi)^4} \int \frac{1}{k^2 - \omega^2} e^{i\vec{k} \cdot \vec{R} - i\omega T} d\vec{k} d\omega$$


$$= \int \frac{1}{k^2 - \omega^2} \hat{\ell}$$


$$d\vec{k} = d^3k = d(\cos\theta) 2\pi k^2 dk$$











$$(2\pi)^4 \int \frac{1}{k^2 - \omega^2} \delta(kR - \omega R) d^3k d\omega$$

$$= \frac{1}{(2\pi)^3} \int_{-\infty}^{\infty} d\omega \int_0^{\infty} \frac{e^{ikR} - e^{-ikR}}{k^2 - \omega^2} \frac{1}{ikR} k^2 dk e^{-i\omega R}$$

111) OTHER FORMS.


$$\boxed{\begin{aligned} \nabla_\mu F^{\mu\nu} &= 4\pi j^\nu \\ F_{(\mu\nu;\alpha\epsilon)}_{cycl} &= 0 \end{aligned}}$$

(SEPTEMBER)

(=)


$$\boxed{\begin{aligned} F &= 4\pi j \\ MF &= 0 \end{aligned}}$$

(JANUARY)




$$= \frac{1}{(2\pi)^3} \int_{-\infty}^{\infty} d\omega \int_0^{\infty} \frac{e^{ikR} - e^{-ikR}}{k^2 - \omega^2} \frac{1}{ikR} k^2 dk e^{-i\omega T}$$

$$I = \int_0^{\infty} \frac{e^{ikR} - e^{-ikR}}{k^2 - \omega^2} k dk = \int_{-\infty}^{\infty} \underbrace{\frac{k e^{ikR}}{k^2 - \omega^2}}_{f(k)} dk$$




$$= \frac{1}{(2\pi)^3} \int_{-\infty}^{\infty} d\omega \int_0^{\infty} \frac{e^{ikR} - e^{-ikR}}{k^2 - \omega^2} \frac{1}{ikR} k^2 dk e^{-i\omega T}$$

$$I = \int_0^{\infty} \frac{e^{ikR} - e^{-ikR}}{k^2 - \omega^2} k dk = \int_{-\infty}^{\infty} \underbrace{\frac{k e^{ikR}}{k^2 - \omega^2}}_{f(k)} dk$$




$$I = \int_0^{\infty} \frac{e^{ikR} - e^{-ikR}}{k^2 - \omega^2} k dk = \int_{-\infty}^{\infty} \frac{k e^{ikR}}{k^2 - \omega^2} dk$$

$$= I_1 + I_2 = i\pi \left(\underbrace{\text{Res}_{-\omega} f(k)}_{\frac{-\omega e^{-ik\omega}}{-2\omega}} + \underbrace{\text{Res}_{\omega} f(k)}_{\frac{\omega e^{ik\omega}}{2\omega}} \right)$$




$$I = \int_0^{\infty} \frac{e^{ikR} - e^{-ikR}}{k^2 - \omega^2} k dk = \int_{-\infty}^{\infty} \frac{k e^{ikR}}{k^2 - \omega^2} dk = \int_{-\infty}^{\infty} \frac{f(k)}{f(k)} dk$$

$$= I_1 + I_2 = i\pi \left(\underbrace{\text{Res}_{/\omega} f(k)}_{\frac{-\omega e^{-ik\omega}}{-2\omega}} + \underbrace{\text{Res}_{/\omega} f(k)}_{\frac{\omega e^{ik\omega}}{2\omega}} \right)$$



$$I = \int_0^{\infty} \frac{e^{ikR} - e^{-ikR}}{k^2 - \omega^2} k dk = \int_{-\infty}^{\infty} \frac{k e^{i k R}}{k^2 - \omega^2} dk = \int_{-\infty}^{\infty} f(k) dk$$

$$= I_1 + I_2 = i\pi \left(\underbrace{\text{Res}_{\omega} f(k)}_{\frac{-\omega e^{-iR\omega}}{-2\omega}} + \underbrace{\text{Res}_{-\omega} f(k)}_{\frac{\omega e^{iR\omega}}{2\omega}} \right)$$




$$I - I_1 - I_2 = 0$$

$$= i\pi e^{-iR\omega}$$

$$I = \int_0^{\infty} \frac{e^{ikR} - e^{-ikR}}{k^2 - \omega^2} k dk = \int_{-\infty}^{\infty} \frac{k e^{ikR}}{k^2 - \omega^2} dk = \int_{-\infty}^{\infty} \frac{f(k)}{k^2 - \omega^2} dk$$

$$= I_1 + I_2 = i\pi \left(\underbrace{\text{Res}_{\omega} f(k)}_{\frac{-\omega e^{-iR\omega}}{-2\omega}} + \underbrace{\text{Res}_{-\omega} f(k)}_{\frac{\omega e^{iR\omega}}{2\omega}} \right)$$



$$I - I_1 - I_2 = 0$$

$$= \frac{i\pi}{2} \left(e^{-iR\omega} + e^{iR\omega} \right)$$

$$\frac{-\omega e^{-iR\omega}}{-2\omega} + \frac{\omega e^{iR\omega}}{2\omega} = \frac{i\pi}{2} (e^{-iR\omega} + e^{iR\omega})$$

$$G = \frac{1}{(2\pi)^3} \frac{1}{iR} \int_{-\infty}^{\infty} d\omega \frac{i\pi}{2} (e^{-iR\omega} + e^{iR\omega}) d\omega e^{-i\omega T}$$

$$= \frac{1}{8\pi} \frac{1}{2\pi} \frac{1}{R} \int d\omega \left[e^{-i\omega(R+T)} + e^{-i\omega(T-R)} \right]$$

$$\frac{-\omega e^{-iR\omega}}{-2\omega} + \frac{\omega e^{iR\omega}}{2\omega} = \frac{i}{2} (e^{-iR\omega} + e^{iR\omega})$$


$$\begin{aligned} G &= \frac{1}{(2\pi)^3} \frac{1}{iR} \int_{-\infty}^{\infty} d\omega \frac{i\pi}{2} (e^{-iR\omega} + e^{iR\omega}) d\omega e^{-i\omega T} \\ &= \frac{1}{8\pi} \frac{1}{2\pi} \frac{1}{R} \int d\omega [e^{-i\omega(R+T)} + e^{-i\omega(T-R)}] \\ &= \frac{1}{8\pi} \frac{1}{R} [\delta(R+T) + \delta(T-R)] \end{aligned}$$


$$\delta(x) = \frac{1}{2\pi} \int dk e^{-ikx}$$

$$\frac{-\omega e^{-iR\omega}}{-2\omega} + \frac{\omega e^{iR\omega}}{2\omega} = \frac{i\pi}{2} (e^{-iR\omega} + e^{iR\omega})$$

$$\begin{aligned} G &= \frac{1}{(2\pi)^3} \frac{1}{iR} \int_{-\infty}^{\infty} d\omega \frac{i\pi}{2} (e^{-iR\omega} + e^{iR\omega}) d\omega e^{-i\omega T} \\ &= \frac{1}{8\pi (2\pi)} \frac{1}{R} \int d\omega [e^{-i\omega(R+T)} + e^{-i\omega(T-R)}] \\ &= \frac{1}{8\pi} \frac{1}{R} [\delta(R+T) + \delta(T-R)] = \frac{1}{4\pi R} \end{aligned}$$

$$\delta(x) = \frac{1}{2\pi} \int dk e^{-ikx}$$





$$\frac{-\omega e^{-iR\omega}}{-2\omega} + \frac{\omega e^{iR\omega}}{2\omega} = \frac{i\pi}{2} (e^{-iR\omega} + e^{iR\omega})$$

$$\begin{aligned} G &= \frac{1}{(2\pi)^3} \frac{1}{iR} \int_{-\infty}^{\infty} d\omega \frac{i\pi}{2} (e^{-iR\omega} + e^{iR\omega}) d\omega e^{-i\omega T} \\ &= \frac{1}{8\pi} \frac{1}{2\pi} \frac{1}{R} \int d\omega [e^{-i\omega(R+T)} + e^{-i\omega(T-R)}] \\ &= \frac{1}{8\pi} \frac{1}{R} [\delta(R+T) + \delta(T-R)] = \frac{1}{4\pi} \frac{\delta(R^2 - T^2)}{R} \end{aligned}$$

$$\delta(x) = \frac{1}{2\pi} \int dk e^{-ikx}$$

$$f'(a_n) = \sum \frac{\partial(x-a_n)}{|f'(a_n)|}$$


$$\frac{-\omega e^{-iR\omega}}{-2\omega} + \frac{\omega e^{iR\omega}}{2\omega} = \frac{i\pi}{2} (e^{-iR\omega} + e^{iR\omega})$$

$$\begin{aligned} G &= \frac{1}{(2\pi)^3} \frac{1}{iR} \int_{-\infty}^{\infty} d\omega \frac{i\pi}{2} (e^{-iR\omega} + e^{iR\omega}) d\omega e^{-i\omega T} \\ &= \frac{1}{8\pi} \frac{1}{2\pi} \frac{1}{R} \int d\omega [e^{-i\omega(R+T)} + e^{-i\omega(T-R)}] \\ &= \frac{1}{8\pi} \frac{1}{R} [\underbrace{\delta(R+T)}_{\text{ADVANCED}} + \underbrace{\delta(T-R)}_{\text{RETARDED}}] = \frac{1}{4\pi} \frac{\delta(R^2 - T^2)}{R} \end{aligned}$$

$$\delta(x) = \frac{1}{2\pi} \int dk e^{-ikx}$$

$$\delta(f(x)) = \sum_n \frac{\delta(x-a_n)}{|f'(a_n)|}$$

$f(x) = 0$



LORENTZ GAUGE COND.

C) WAVE EQ. (1) & (2) SOLVED BY A, ϕ

$$\begin{aligned} (3) \quad \nabla \cdot \left(-\nabla \phi - \frac{\partial \mathbf{A}}{\partial t} \right) &= 4\pi \rho \\ -\Delta \phi - \frac{\partial}{\partial t} \underbrace{\nabla \cdot \mathbf{A}}_{=0} &= 4\pi \rho \\ -\Delta \phi &= 4\pi \rho \end{aligned}$$

$$\boxed{\square \phi = -4\pi \rho}$$

$$\boxed{\square = -\frac{\partial^2}{\partial t^2} + \Delta}$$

D'ALAMBERT OP.

WAVE EQ.

IN CLASSICAL ELECTRODYNAMICS WE TAKE ONLY RETARDED

LORENZ GAUGE COND.

C) WAVE EQ. (1) & (2) SOLVED BY A, ϕ

$$\begin{aligned} (3) \quad \nabla \cdot \left(-\nabla \phi - \frac{\partial \mathbf{A}}{\partial t} \right) &= 4\pi \rho \\ -\Delta \phi - \frac{\partial}{\partial t} \underbrace{\nabla \cdot \mathbf{A}}_{=0} &= 4\pi \rho \\ -\Delta \phi &= 4\pi \rho \end{aligned}$$

$$\boxed{\square \phi = -4\pi \rho}$$

$$\boxed{\square = -\frac{\partial^2}{\partial t^2} + \Delta}$$

D'ALAMBERT OP.

WAVE EQ.

IN CLASSICAL ELECTRODYNAMICS WE TAKE ONLY RETARDED
"THROWING AWAY $\frac{1}{2}$ PHYSICS"

$$\phi(\mathbf{r}, t) = \int G(\mathbf{r}, \mathbf{r}', t, t')$$

LORENTZ GAUGE COND.

C) WAVE EQ. (4) & (2) SOLVED BY A, ϕ

$$\square = -\frac{\partial^2}{\partial t^2} + \Delta$$

IN CLASSICAL ELECTRODYNAMICS WE TAKE ONLY RETARDED
"THROWING AWAY $\frac{1}{2}$ PHYSICS"

$$\begin{aligned}\phi(\mathbf{r}, t) &= \int G(\mathbf{r}, \mathbf{r}', t, t') 4\pi \rho(\mathbf{r}', t') d\mathbf{r}' dt' \\ &= \int \mathcal{D}\end{aligned}$$

LORENZ GAUGE COND.

C) WAVE EQ. (1) & (2) SOLVED BY A.ϕ

$$\square = -\frac{\partial^2}{\partial t^2} + \Delta$$

IN CLASSICAL ELECTRODYNAMICS WE TAKE ONLY RETARDED
"THROWING AWAY $\frac{1}{2}$ PHYSICS"

$$\begin{aligned}\Phi(\mathbf{r}, t) &= \int G(\mathbf{r}, \mathbf{r}', t, t') 4\pi \rho(\mathbf{r}', t') d\mathbf{r}' dt' \\ &= \int \frac{\delta(t - t' - |\mathbf{r} - \mathbf{r}'|)}{|\mathbf{r} - \mathbf{r}'|} \rho(\mathbf{r}', t') d\mathbf{r}' dt'\end{aligned}$$

LORENZ GAUGE COND.

C) WAVE EQ. (1) & (2) SOLVED BY A. & ϕ

$$\square = -\frac{\partial^2}{\partial t^2} + \Delta$$


IN CLASSICAL ELECTRODYNAMICS WE TAKE ONLY RETARDED
"THROWING AWAY $\frac{1}{2}$ PHYSICS"

$$\begin{aligned}\phi(\mathbf{r}, t) &= \int G(\mathbf{r}, \mathbf{r}', t, t') 4\pi \rho(\mathbf{r}', t') d\mathbf{r}' dt' \\ &= \int \frac{\delta(t - t' - |\mathbf{r} - \mathbf{r}'|)}{|\mathbf{r} - \mathbf{r}'|} \rho(\mathbf{r}', t') d\mathbf{r}' dt' = \int d\mathbf{r}'\end{aligned}$$

$$\begin{aligned}\phi(r,t) &= \int G(r,r',t,t') 4\pi g(r',t') dr' dt' \\ &= \int \frac{\delta(t-t' - |r-r'|)}{|r-r'|} g(r',t') dr' dt' = \boxed{\int dr' \frac{g(r',t-|r-r'|)}{|r-r'|} = \phi(r,t)}\end{aligned}$$

$$(4) \Leftrightarrow \boxed{\square A = -4\pi g}$$

$$\boxed{\square A^\mu = -4\pi g^\mu}$$



FEYNMAN'S STORY:
PROBLEM OF ELECTRON



FEYNMAN'S STORY:

PROBLEM OF ELECTRON




NEGATIVE PRESS.

SELF-FORCE

FEYNMAN'S STORY:

PROBLEM OF ELECTRON



NEGATIVE PRESS.

'SELF-FORCE' ON ITSELF

$$F_{\text{SELF}} = -\frac{U_{\text{SELF}}}{c} \alpha + \frac{2}{3}$$

FEYNMAN'S STORY

PROBLEM OF ELECTRON



NEGATIVE PRESS.

'SELF-FORCE' ON ITSELF

$$F_{\text{SELF}} = -U_{\text{SELF}} a + \frac{2}{3} e^2 \frac{da}{dt} + \sum_{n=2}^{\infty} (-1)^n \frac{d^n a}{dt^n}$$

FEYNMAN'S STORY:

PROBLEM OF ELECTRON



NEGATIVE PRESS.

'SELF-FORCE' ON ITSELF

$$F_{\text{SELF}} = -U_{\text{SELF}} a + \frac{2}{3} e^2 \frac{da}{dt} + \sum_{n=2}^{\infty} (-1)^n \frac{d^n a}{dt^n} O(R^{n-1})$$

FEYNMAN'S STORY

PROBLEM OF ELECTRON



NEGATIVE CHARGES

'SELF-FORCE' ON ITSELF


$$F_{\text{SELF}} = -\frac{1}{2} \frac{d}{dt} \left(\frac{2}{3} R^2 \frac{da}{dt} \right) + \sum_{n=2}^{\infty} (-1)^n \frac{d^n a}{dt^n} O(R^{n-1})$$

2 POSS - R-FINITE . DISASTER

FEYNMAN'S STORY.

PROBLEM OF ELECTRON

$V_{\text{SELF}} =$



NEGATIVE PRESS.


'SELF-FORCE' ON ITSELF

$$F_{\text{SELF}} = -V_{\text{SELF}} \dot{a} + \frac{2}{3} \dot{a}^2 \frac{da}{dt} + \sum_{n=2}^{\infty} (-1)^n \frac{d^n a}{dt^n} O(R^{n-1})$$

2 POSS. $\left\{ \begin{array}{l} R\text{-FINITE} \quad \text{DISASTER} \\ R \rightarrow 0 \end{array} \right.$

FEYNMAN'S STORY.

PROBLEM OF ELECTRON



'SELF-FORCE' ON ITSELF


$$F_{\text{SELF}} = -V_{\text{SELF}} \dot{a} + \frac{2}{3} \dot{a}^2 \frac{da}{dt} + \sum_{n=2}^{\infty} (-1)^n \frac{d^n a}{dt^n} O(R^{n-1})$$

2 POSS: $\left\{ \begin{array}{l} R \text{ - FINITE} \quad \text{DISASTER} \\ R \rightarrow 0 \end{array} \right.$

FEYNMAN'S STORY.

PROBLEM OF ELECTRON

$$U_{\text{SELF}} = \int E^2 dV$$

$$\sim \int \frac{1}{r^2} r^2 dr$$


NEGATIVE PRESS.

'SELF-FORCE' ON ITSELF

$$F_{\text{SELF}} = -U_{\text{SELF}} a + \frac{2}{3} e^2 \frac{da}{dt} + \sum_{n=2}^{\infty} (-1)^n \frac{d^n a}{dt^n} O(R^{n-1})$$

2 POSS: $\left\{ \begin{array}{l} R\text{-FINITE} \quad \text{DISASTER} \\ R \rightarrow 0 \end{array} \right.$



FEYNMAN'S STORY.

PROBLEM OF ELECTRON

$$U_{\text{SELF}} = \frac{1}{2} \epsilon_0 \int E^2 dV$$

$$\sim \int \frac{1}{r^2} r^2 dr$$

$$\sim \frac{1}{R}$$


NEGATIVE PRESS.

'SELF-FORCE' ON ITSELF

$$F_{\text{SELF}} = -U_{\text{SELF}} a + \frac{2}{3} e^2 \frac{da}{dt} + \sum_{n=2}^{\infty} (-1)^n \frac{d^n a}{dt^n} O(R^{n-1})$$

2 POSS: R-FINITE . DISASTER

$R \rightarrow 0$

$U_{\text{SELF}} \rightarrow \infty$

FEYNMAN'S STORY


PROBLEM OF ELECTRON

$$U_{\text{SELF}} = \int E^2 dV$$

$$\sim \int \frac{1}{r^2} r^2 dr$$

$$\sim \frac{1}{R}$$

NEGATIVE PROBS.



'SELF-FORCE' ON ITSELF

$$F_{\text{SELF}} = -U_{\text{SELF}} \dot{a} + \frac{2}{3} \dot{a}^2 \frac{da}{dt} + \sum_{n=2}^{\infty} (-1)^n \frac{d^n a}{dt^n} O(R^{n-1})$$

2 POSS: R-FINITE DISASTER

$$R \rightarrow 0 \quad U_{\text{SELF}} \rightarrow \infty$$

FEYNMAN'S STORY.

PROBLEM OF ELECTRON

$$U_{\text{SELF}} = \int E^2 dV$$

$$\sim \int \frac{1}{r^2} r^2 dr$$

$$\sim \frac{1}{R}$$

NEGATIVE PRESS.




'SELF-FORCE' ON ITSELF


$$F_{\text{SELF}} = -U_{\text{SELF}} a + \frac{2}{3} e^2 \frac{da}{dt} + \sum_{n=2}^{\infty} (-1)^n \frac{d^n a}{dt^n} O(R^{n-1})$$

2 POSS. $\left\{ \begin{array}{l} R \text{ FINITE} \quad \text{DISASTER} \\ R \rightarrow 0 \end{array} \right.$


$$F = ma$$

$U_{\text{SELF}} \rightarrow \infty$
(RENORMALIZATION)









"MAKING THE THEORY"


E: • ACTION AT A DISTANCE DOESN'T NEED FIELD $\Rightarrow$
 $\Rightarrow$ NO SELF ENERGY PROBLEM

• HOW EXPLAIN RADIATION DAMPING FORCE ?



W: • F_D WOULD DEPEND ON $q_2, m_2, \frac{1}{R_2}$, THICKNESS OF LAYER
 AND OCCURS AT THE WRONG TIME

"YOU HAVE DESCRIBED ORDINARY REFLECTED LIGHT"



• LET ALL CHARGES PRODUCE


$$F_{\mu\nu} = R F_{\mu\nu}^{\text{RET}} + (1-R) F_{\mu\nu}^{\text{ADV}}$$

• FOR $R = 1/2$ TP SUFFERS JUST
 FROM RETARDED POTENTIALS

• FOR LIMITED NUMBER OF CHARGES

W: • F_D WOULD DEPEND ON $q_2, m_2, \frac{1}{R_2}$, THICKNESS OF LAYER
AND OCCURS AT THE WRONG TIME

"YOU HAVE DESCRIBED ORDINARY REFLECTED LIGHT"



• LET ALL CHARGES PRODUCE

$$F_{\mu\nu} = k F_{\mu\nu}^{\text{RET}} + (1-k) F_{\mu\nu}^{\text{ADV}}$$

• FOR $k = 1/2$ TP SUFFERS JUST
FROM RETARDED POTENTIALS

• FOR LIMITED NUMBER OF CHARGES
IN UNIVERSE ADVANCED EFFECTS


OCCUR EXPLICITLY $\Rightarrow$ NO SEPARATION

BETWEEN PAST AND FUTURE

• BUT FOR PERFECT ABSORBER THE REACTION BACK IS INDEPENDENT
OF IT'S STRUCTURE

W: • F_D WOULD DEPEND ON $q_2, m_2, \frac{1}{R_2}$, THICKNESS OF LAYER
AND OCCURS AT THE WRONG TIME

"YOU HAVE DESCRIBED ORDINARY REFLECTED LIGHT"



• LET ALL CHARGES PRODUCE
 $F_{\mu\nu} = k F_{\mu\nu}^{\text{RET}} + (1-k) F_{\mu\nu}^{\text{ADV}}$


• FOR $k = 1/2$ TP SUFFERS JUST
FROM RETARDED POTENTIALS

• FOR LIMITED NUMBER OF CHARGES
IN UNIVERSE ADVANCED EFFECTS

OCCUR EXPLICITLY $\Rightarrow$ NO SEPARATION

BETWEEN PAST AND FUTURE

• BUT FOR PERFECT ABSORBER THE REACTION BACK IS INDEPENDENT
OF IT'S STRUCTURE



- NOTIONS:
- 1) THEORY IS MICROSCOPICALLY REVERSIBLE (TIME SYMMETRIC)
 - 2) DESPITE THIS IT IS ABLE TO DESCRIBE DAMPING FORCE
(ABSORBER ENTROPY IS INCREASING)
 - 3) CAN BE DESCRIBED BY FOKKER'S ACTION

$$A_F = - \sum_{i=1}^N m_i \int (\dot{x}_i^M \dot{x}_i^M)^{1/2} d\tau_i + \frac{1}{2} \sum_{i \neq j} e_i e_j \int \int \delta((x_{iM} - x_{jM})^2) \dot{x}_i^M \dot{x}_j^M d\tau_i d\tau_j$$

$\Rightarrow$ PURE PARTICLE-PARTICLE INTERACTION

• THANKS TO $\delta((x_{iM} - x_{jM})^2)$ PARTICLES INTERACT
ONLY AT ZERO FOUR-DIMENSIONAL DISTANCE

- 4) IT IS ADJUNCT FIELD THEORY:

FIELD THEORY $\approx$ HAMILTONIAN METHOD (TIME DIF. METH.)

• I HAVE CONCEPT OF FIELD ($\approx$ MATHEMATICAL DESCR.)

- THIS FIELD HAS TO BE GIVEN EVERYWHERE AT $\pm$

DIF. EQUATIONS GIVE ME THE FIELD AT THE NEXT TIME

(ABSORBER ENTROPY IS INCREASING)

3) CAN BE DESCRIBED BY FOKKER'S ACTION

$$A_F = - \sum_{i=1}^N m_i \int (\dot{x}_i^M \dot{x}_i^M)^{1/2} d\tau_i + \frac{4}{2} \sum_{i \neq j} e_i e_j \iint \delta((x_{i1} - x_{j1})^2) \dot{x}_{i1}^M \dot{x}_{j1}^M d\tau_i d\tau_j$$

⇒ PURE PARTICLE-PARTICLE INTERACTION

• THANKS TO $\delta((x_{i1} - x_{j1})^2)$ PARTICLES INTERACT
ONLY AT ZERO FOUR-DIMENSIONAL DISTANCE


4) IT IS ADJUNCT FIELD THEORY:

FIELD THEORY ≈ HAMILTONIAN METHOD (TIME DIF. METH.)

• I HAVE CONCEPT OF FIELD (≈ MATHEMATICAL DESCR.)

- THIS FIELD HAS TO BE GIVEN EVERYWHERE AT ±
DIF. EQUATIONS GIVE ME THE FIELD AT THE NEXT TIME

NOTE: 'PHYSICAL FIELD' IS A
SOLENOID:



Einstein says, "Nooooooooooooo," a nice, German-sounding "No,"—very polite. "I find only that it would be very difficult to make a corresponding theory for gravitational interaction." He meant for the general theory of relativity, which was his baby. He continued: "Since we have at this time not a great deal of experimental evidence, I am not absolutely sure of the correct gravitational theory." Einstein appreciated that things might be different from what his theory stated; he was very tolerant of other ideas.

I wish I had remembered what Pauli said, because I discovered years later that the theory was not satisfactory when it came to making the quantum theory. It's possible that that great man noticed the difficulty immediately and explained it to me in the question, but I was so relieved at not having to answer the questions that I didn't really listen to them carefully. I do remember walking up the steps of Palmer Library with Pauli, who said to me, "What is Wheeler going to say about the quantum theory when he gives his talk?"

I said, "I don't know. He hasn't told me. He's working it out himself."

"Oh?" he said. "The man works and doesn't tell his assistant what he's doing on the quantum theory?" He came closer to me and said in a low, secretive voice, "Wheeler will never give that seminar."

And it's true. Wheeler didn't give the seminar. He thought it would be easy to work out the quantum part; he thought he had it, almost. But he didn't. And by the time the seminar came around, he realized he didn't know how to do it, and therefore didn't have anything to say.

I never solved it, either—a quantum theory of half-advanced, half-retarded potentials—and I worked on it for years.

got written—that never struck me! So I developed a negative attitude toward the guy who studies French literature, or studies too much music or poetry—all those "fancy" things. I admired better the steel-worker, the welder, or the machine shop man. I always thought the guy who worked in the machine shop and could make things, now *he* was a *real* guy! That was my attitude. To be a practical man was, to me, always somehow a positive virtue, and to be "cultured" or "intellectual" was not. The first was right, of course, but the second was crazy.

I still had this feeling when I was doing my graduate study at Princeton, as you'll see. I used to eat often in a nice little restaurant called Papa's Place. One day, while I was eating there, a painter in his painting clothes came down from an upstairs room he'd been painting, and sat near me. Somehow we struck up a conversation and he started talking about how you've got to learn a lot to be in the painting business. "For example," he said, "in this restaurant, what colors would you use to paint the walls, if you had the job to do?"

I said I didn't know, and he said, "You have a dark band up to such-and-such a height, because, you see, people who sit at the tables rub their elbows against the walls, so you don't want a nice, white wall

Mixing Paints