

Title: 12/13 PSI - Found Quantum Mechanics Lecture 14

Date: Jan 24, 2013 11:30 AM

URL: <http://pirsa.org/13010083>

Abstract:

Inconsistencies of the orthodox interpretation

By the collapse postulate
(applied to the system)

By unitary evolution postulate
(applied to isolated system that
includes the apparatus)

Indeterministic and
discontinuous evolution

Deterministic and
continuous evolution

Determinate properties

Indeterminate properties

Responses to the measurement problem

1. Deny universality of quantum dynamics
 - Quantum-classical hybrid models
 - Collapse models
2. Deny representational completeness of ψ
 - ψ -ontic hidden variable models (e.g. deBroglie-Bohm)
 - ψ -epistemic hidden variable models
3. Deny that there is a unique outcome
 - Everett's relative state interpretation (many worlds)
4. Deny some aspect of classical logic or classical probability theory
 - Quantum logic and quantum Bayesianism
5. Deny some other feature of the realist framework?

Collapse theories

Posit a new dynamical evolution law:
either **nonlinear** or **indeterministic** or both

Recover unitary evolution and the collapse dynamics as special cases

Microscopic systems obey **unitary** dynamics to good approximation
Macroscopic systems obey **collapse** dynamics to good approximation

Collapse theories

Posit a new dynamical evolution law:
either **nonlinear** or **indeterministic** or both

Recover unitary evolution and the collapse dynamics as special cases

Microscopic systems obey **unitary** dynamics to good approximation
Macroscopic systems obey **collapse** dynamics to good approximation

Motivations:

- Achieves realism
- Maintains representational completeness of ψ
- No “cut”, i.e. one universal dynamics (unlike a hybrid model)

Nonlinear deterministic models

$$|\uparrow\rangle|“ready”\rangle \rightarrow |\uparrow\rangle|“up”\rangle$$

$$|\downarrow\rangle|“ready”\rangle \rightarrow |\downarrow\rangle|“down”\rangle$$

$$(a|\uparrow\rangle + b|\downarrow\rangle)|“ready”\rangle \rightarrow a|\uparrow\rangle|“up”\rangle + b|\downarrow\rangle|“down”\rangle$$

~~$a|\uparrow\rangle|“up”\rangle + b|\downarrow\rangle|“down”\rangle$~~

rather $(a|\uparrow\rangle + b|\downarrow\rangle)|“ready”\rangle \rightarrow |\uparrow\rangle|“up”_{a,b}\rangle$

or

$$|\downarrow\rangle|“down”_{a,b}\rangle$$

Final state depends on details of the initial state

Ignorance of those details implies subjective indeterminism

Linear indeterministic models

The goal:

$$(a| \uparrow \rangle + b| \downarrow \rangle)| \text{"ready"} \rangle \rightarrow | \uparrow \rangle| \text{"up"} \rangle \text{ with probability } |a|^2$$
$$\rightarrow | \downarrow \rangle| \text{"down"} \rangle \text{ with probability } |b|^2$$

The preferred decomposition issue

Into what states do collapses occur?

The trigger issue

When and how do collapses occur?

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) = H\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t + dt) = \frac{1}{\sqrt{p(\mathbf{q}_k)}} G_{\mathbf{q}_k}(\mathbf{r}_k) \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{"Collapse"}$$

where $G_{\mathbf{q}_k}(\mathbf{r}_k) = K \exp\left(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{2\sigma^2}\right)$

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) = H\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t + dt) = \frac{1}{\sqrt{p(\mathbf{q}_k)}} G_{\mathbf{q}_k}(\mathbf{r}_k) \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{"Collapse"}$$

where $G_{\mathbf{q}_k}(\mathbf{r}_k) = K \exp\left(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{2\sigma^2}\right)$

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) = H\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t + dt) = \frac{1}{\sqrt{p(\mathbf{q}_k)}} G_{\mathbf{q}_k}(\mathbf{r}_k) \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{"Collapse"}$$

where $G_{\mathbf{q}_k}(\mathbf{r}_k) = K \exp\left(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{2\sigma^2}\right)$

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) = H\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t + dt) = \frac{1}{\sqrt{p(\mathbf{q}_k)}} G_{\mathbf{q}_k}(\mathbf{r}_k) \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{"Collapse"}$$

where $G_{\mathbf{q}_k}(\mathbf{r}_k) = K \exp\left(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{2\sigma^2}\right)$

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) = H\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t + dt) = \frac{1}{\sqrt{p(\mathbf{q}_k)}} G_{\mathbf{q}_k}(\mathbf{r}_k) \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{"Collapse"}$$

where $G_{\mathbf{q}_k}(\mathbf{r}_k) = K \exp\left(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{2\sigma^2}\right)$

$$p(\mathbf{q}_k) = \int d\mathbf{r}_1 \dots d\mathbf{r}_N |G_{\mathbf{q}_k}(\mathbf{r}_k) \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t)|^2$$

k is chosen uniformly at random

\mathbf{q}_k is chosen by sampling from $p(\mathbf{q}_k)$

Two new fundamental constants:

$\tau \approx 10^{15} \text{ s} \approx 100 \text{ million years}$ mean time between collapses for one particle

$\sigma \approx 10^{-7} \text{ m} \approx \text{size of large molecule}$ Localization width

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) = H\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$\psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t + dt) = \frac{1}{\sqrt{p(\mathbf{q}_k)}} G_{\mathbf{q}_k}(\mathbf{r}_k) \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad \text{"Collapse"}$$

where $G_{\mathbf{q}_k}(\mathbf{r}_k) = K \exp\left(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{2\sigma^2}\right)$

$$p(\mathbf{q}_k) = \int d\mathbf{r}_1 \dots d\mathbf{r}_N |G_{\mathbf{q}_k}(\mathbf{r}_k) \psi(\mathbf{r}_1, \dots, \mathbf{r}_N, t)|^2$$

k is chosen uniformly at random

\mathbf{q}_k is chosen by sampling from $p(\mathbf{q}_k)$

Two new fundamental constants:

$\tau \approx 10^{15} \text{ s} \approx 100 \text{ million years}$ mean time between collapses for one particle

$\sigma \approx 10^{-7} \text{ m} \approx \text{size of large molecule}$ Localization width

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} |\psi(t)\rangle = H |\psi(t)\rangle \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$|\psi(t+dt)\rangle = \frac{1}{\sqrt{p(\mathbf{q}_k)}} \sqrt{E^{(k)}(\mathbf{q}_k)} |\psi(t)\rangle \quad \text{"Collapse"}$$

where $E^{(k)}(\mathbf{q}_k) = \int d\mathbf{r}_k K \exp(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{\sigma^2}) |\mathbf{r}_k\rangle \langle \mathbf{r}_k|$

$$p(\mathbf{q}_k) = \langle \psi(t) | E^{(k)}(\mathbf{q}_k) | \psi(t) \rangle$$

k is chosen uniformly at random

\mathbf{q}_k is chosen by sampling from $p(\mathbf{q}_k)$

Two new fundamental constants:

$\tau \approx 10^{15} \text{ s} \approx 100 \text{ million years}$ mean time between collapses for one particle

$\sigma \approx 10^{-7} \text{ m} \approx \text{size of large molecule}$ Localization width

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} |\psi(t)\rangle = H |\psi(t)\rangle \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$|\psi(t+dt)\rangle = \frac{1}{\sqrt{p(\mathbf{q}_k)}} \sqrt{E^{(k)}(\mathbf{q}_k)} |\psi(t)\rangle \quad \text{"Collapse"}$$

where $E^{(k)}(\mathbf{q}_k) = \int d\mathbf{r}_k K \exp(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{\sigma^2}) |\mathbf{r}_k\rangle \langle \mathbf{r}_k|$

$$p(\mathbf{q}_k) = \langle \psi(t) | E^{(k)}(\mathbf{q}_k) | \psi(t) \rangle$$

k is chosen uniformly at random

\mathbf{q}_k is chosen by sampling from $p(\mathbf{q}_k)$

Two new fundamental constants:

$\tau \approx 10^{15} \text{ s} \approx 100 \text{ million years}$ mean time between collapses for one particle

$\sigma \approx 10^{-7} \text{ m} \approx \text{size of large molecule}$ Localization width

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} |\psi(t)\rangle = H |\psi(t)\rangle \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$|\psi(t+dt)\rangle = \frac{1}{\sqrt{p(\mathbf{q}_k)}} \sqrt{E^{(k)}(\mathbf{q}_k)} |\psi(t)\rangle \quad \text{"Collapse"}$$

where $E^{(k)}(\mathbf{q}_k) = \int d\mathbf{r}_k K \exp(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{\sigma^2}) |\mathbf{r}_k\rangle \langle \mathbf{r}_k|$

$$p(\mathbf{q}_k) = \langle \psi(t) | E^{(k)}(\mathbf{q}_k) | \psi(t) \rangle$$

k is chosen uniformly at random

\mathbf{q}_k is chosen by sampling from $p(\mathbf{q}_k)$

Two new fundamental constants:

$\tau \approx 10^{15} \text{ s} \approx 100 \text{ million years}$ mean time between collapses for one particle

$\sigma \approx 10^{-7} \text{ m} \approx \text{size of large molecule}$ Localization width

In terms of density operators

$$\rho \rightarrow \frac{\sqrt{E^{(k)}(\mathbf{q}_k)}\rho\sqrt{E^{(k)}(\mathbf{q}_k)}}{\text{Tr}(E^{(k)}(\mathbf{q}_k)\rho)}$$

Essentially a POVM measurement of approximate position

The Ghirardi-Rimini-Weber model

At most times:

$$i\hbar \frac{\partial}{\partial t} |\psi(t)\rangle = H |\psi(t)\rangle \quad \text{Schrödinger's equation}$$

Every τ/N time interval on average

$$|\psi(t+dt)\rangle = \frac{1}{\sqrt{p(\mathbf{q}_k)}} \sqrt{E^{(k)}(\mathbf{q}_k)} |\psi(t)\rangle \quad \text{"Collapse"}$$

where $E^{(k)}(\mathbf{q}_k) = \int d\mathbf{r}_k K \exp(-\frac{(\mathbf{r}_k - \mathbf{q}_k)^2}{\sigma^2}) |\mathbf{r}_k\rangle \langle \mathbf{r}_k|$

$$p(\mathbf{q}_k) = \langle \psi(t) | E^{(k)}(\mathbf{q}_k) | \psi(t) \rangle$$

k is chosen uniformly at random

\mathbf{q}_k is chosen by sampling from $p(\mathbf{q}_k)$

Two new fundamental constants:

$\tau \approx 10^{15} \text{ s} \approx 100 \text{ million years}$ mean time between collapses for one particle

$\sigma \approx 10^{-7} \text{ m} \approx \text{size of large molecule}$ Localization width

In terms of density operators

$$\rho \rightarrow \frac{\sqrt{E^{(k)}(\mathbf{q}_k)}\rho\sqrt{E^{(k)}(\mathbf{q}_k)}}{\text{Tr}(E^{(k)}(\mathbf{q}_k)\rho)}$$


Essentially a POVM measurement of approximate position

If \mathbf{k} and \mathbf{q}_k are unknown, then the effective evolution is

$$\rho \rightarrow \sum_k \int dq_k \sqrt{E^{(k)}(\mathbf{q}_k)} \rho \sqrt{E^{(k)}(\mathbf{q}_k)}$$

Single particle in 1D

$$\psi(x) = \frac{\sqrt{3}}{2}\phi_a(x) + \frac{1}{2}\phi_b(x)$$



Single particle in 1D

$$\psi(x) = \frac{\sqrt{3}}{2}\phi_a(x) + \frac{1}{2}\phi_b(x)$$



$$G_q(x) = K \exp\left(-\frac{(x-q)^2}{2\sigma^2}\right)$$

\times

=

Two particles in 1D

$$\psi(x_1, x_2) = \frac{\sqrt{3}}{2}\phi_a(x_1)\chi_a(x_2) + \frac{1}{2}\phi_b(x_1)\chi_b(x_2)$$




$$\psi'(x_1, x_2) \approx \phi_a(x_1)\chi'_a(x_2)$$



=

Single particle in 1D

$$\psi(x) = \frac{\sqrt{3}}{2}\phi_a(x) + \frac{1}{2}\phi_b(x)$$





$$G_q(x) = K \exp\left(-\frac{(x-q)^2}{2\sigma^2}\right)$$





Two particles in 1D

$$\psi(x_1, x_2) = \frac{\sqrt{3}}{2}\phi_a(x_1)\chi_a(x_2) + \frac{1}{2}\phi_b(x_1)\chi_b(x_2)$$




$$\psi'(x_1, x_2) \approx \phi_a(x_1)\chi'_a(x_2)$$





$$\psi = \alpha \phi_a(\mathbf{r}_1) \chi_a(\mathbf{r}_2, \dots, \mathbf{r}_M) + \beta \phi_b(\mathbf{r}_1) \chi_b(\mathbf{r}_2, \dots, \mathbf{r}_M)$$




$$\psi = \alpha \phi_a(\mathbf{r}_1) \chi_a(\mathbf{r}_2, \dots, \mathbf{r}_M) + \beta \phi_b(\mathbf{r}_1) \chi_b(\mathbf{r}_2, \dots, \mathbf{r}_M)$$

Suppose $\chi_a(\dots \mathbf{r}_k \dots) \chi_b(\dots \mathbf{r}_k \dots) \approx 0$ for macroscopic # of components

One particle is hit \rightarrow wavefn' is localized in all coordinates

$$\psi' = \phi_a(\mathbf{r}_1) \chi'_a(\mathbf{r}_2, \dots, \mathbf{r}_M) \text{ with probability } |\alpha|^2$$

$$\psi' = \phi_b(\mathbf{r}_1) \chi'_b(\mathbf{r}_2, \dots, \mathbf{r}_M) \text{ with probability } |\beta|^2$$



$$\psi = \alpha \phi_a(\mathbf{r}_1) \chi_a(\mathbf{r}_2, \dots, \mathbf{r}_M) + \beta \phi_b(\mathbf{r}_1) \chi_b(\mathbf{r}_2, \dots, \mathbf{r}_M)$$

Suppose $\chi_a(\dots \mathbf{r}_k \dots) \chi_b(\dots \mathbf{r}_k \dots) \approx 0$ for macroscopic # of components

One particle is hit \rightarrow wavefn' is localized in all coordinates

$$\psi' = \phi_a(\mathbf{r}_1) \chi'_a(\mathbf{r}_2, \dots, \mathbf{r}_M) \text{ with probability } |\alpha|^2$$

$$\psi' = \phi_b(\mathbf{r}_1) \chi'_b(\mathbf{r}_2, \dots, \mathbf{r}_M) \text{ with probability } |\beta|^2$$

For $M \approx 10^{20}$ particles

This happens every $\frac{10^{15} \text{ s}}{10^{20}} \approx 10^{-5} \text{ s}$

Constraints on parameters

- τ too big \rightarrow persistence of coherence of macro objects
- τ too small \rightarrow loss of coherence of micro objects

Constraints on parameters

- τ too big \rightarrow persistence of coherence of macro objects
- τ too small \rightarrow loss of coherence of micro objects

Constraints on parameters

τ too big \rightarrow persistence of coherence of macro objects

τ too small \rightarrow loss of coherence of micro objects

σ too big \rightarrow delocalized macro objects

σ too small \rightarrow excitation and heating

Experimental status

Difficult to distinguish fundamental collapse from decoherence

Difficult to detect anomalous heating

Continuous Spontaneous localization

Philip Pearle

Collapse is a continuous process governed by a randomly fluctuating field
"gambler's ruin"

Criticisms

What is the ontology in space-time? The "tails" problem.
Bell's "flashes" ontology

Failure of energy conservation

Criticisms

What is the ontology in space-time? The "tails" problem.
Bell's "flashes" ontology

Failure of energy conservation

Unclear whether one can maintain Lorentz invariance for
relativistic models

Criticisms

What is the ontology in space-time? The "tails" problem.
Bell's "flashes" ontology

Failure of energy conservation

Unclear whether one can maintain Lorentz invariance for
relativistic models

Any anomalous decoherence is also consistent with unitary coupling
to novel degrees of freedom

Criticisms

What is the ontology in space-time? The "tails" problem.
Bell's "flashes" ontology

Failure of energy conservation

Unclear whether one can maintain Lorentz invariance for
relativistic models

Any anomalous decoherence is also consistent with unitary coupling
to novel degrees of freedom

An alternative way of accounting for any "anomalous decoherence"

The map corresponding to the "anomalous decoherence" is


$$\rho \rightarrow \sum_k \int dq_k \sqrt{E^{(k)}(\mathbf{q}_k)} \rho \sqrt{E^{(k)}(\mathbf{q}_k)}$$

This is a **trace-preserving completely positive linear map**


$$\rho \rightarrow \mathcal{T}(\rho) = \sum_{\mu} A_{\mu} \rho A_{\mu}^{\dagger}$$

where the A_{μ} are linear operators satisfying $\sum_{\mu} A_{\mu}^{\dagger} A_{\mu} = I$

Church of the larger Hilbert space



Church of the larger Hilbert space



GRW and CSL:
of fundamental constituents in an object → rate of collapse

Other ideas for what induces dynamical collapse in macroscopic systems but not microscopic systems:

gravity

Complexity/entanglement

GRW and CSL:
of fundamental constituents in an object → rate of collapse

Other ideas for what induces dynamical collapse in macroscopic systems but not microscopic systems:

gravity

Complexity/entanglement