Title: Cosmic Tides


Date: Jan 29, 2013 11:00 AM

URL: http://www.pirsa.org/13010114

Abstract: We apply CMB lensing techniques to large scale structure and solve for the 3-D cosmic tidal field. We use small scale filamentary structures to solve for the large scale tidal shear and gravitational potential.

By comparing this to the redshift space density field, one can measure the gravitational growth factor on large scales without cosmic variance. This potentially enables accurate measurements of neutrino masses and reconstruction of radial modes lost in 21 cm intensity mapping, which are essential for CMB and other cross correlations. We relate the tidal fields to the squeezed limit bispectrum, and present initial results from simulations and data from the SDSS.

Pirsa: 13010114 Page 1/23



Pirsa: 13010114 Page 2/23

U. Pen

Introduction Tides Cosmic Variance Summary Overview ► What is a tide? ► Tides and large scale structure Overcoming cosmic variance ► Neutrinos and more


Pirsa: 13010114 Page 3/23

U. Pen

Beating Cosmic Variance with Co

Reconstruction Data

Tides



Graphic of tidal forces; the gravity field is generated by a body to the right. The top picture shows the gravitational forces; the bottom shows their residual once the field of the sphere is subtracted; this is the tidal force. (From Wikipedia)


U. Pen

Beating Cosmic Variance with Cosmic Tides

Pirsa: 13010114 Page 4/23


Reconstruction Data

Tides on Earth



The Moon's gravity differential field at the surface of the Earth is known as the Tide Generating Force. This is the primary mechanism driving tidal action, explaining two tidal equipotential bulges, and accounting for two high tides per day. In this figure, the Moon is either on the right side or on the left side of the Earth (at center). (truncated from Wikipedia)

U. Pen



Pirsa: 13010114 Page 6/23

Gravity reconstruction

- analogous to CMB lensing, 21cm lensing of LSS
- large scale tidal fields aligns smaller scale structures
- use alignment of small filaments to measure large scale tidal field
- perfectly understood for Gaussian fields (Challinor, Lewis, etc)
- optimal quadratic estimators for non-Gaussian fields (Lu and Pen 2008)
- Simple framework for understanding information saturation (White, Meiksen, Rimes, Hamilton)
- equivalent to squeezed bispectrum quadrupole (Creminelli et al 2011)



U. Pen

Heuristic kernel: CMB Lensing

Gradient of the density field convolved by a Gaussian window.

$$W = \exp\left(\frac{-r^2}{2\sigma^2}\right) \tag{1}$$

then we obtain a windowed gradient density field

$$\bar{\delta} = \int W \delta. \tag{2}$$

Take log to Gaussianize: $\delta_g = \log(1+\bar{\delta})$


The tidal shear field is the quadratic outer product

$$T_{ij} = (\partial_i \delta_g)(\partial_j \delta_g). \tag{3}$$

The density field is derived from a convolution:

$$\langle \delta \rangle(x) = \int T_{ij}(x') K_{ij}(x - x') d^3 x' \tag{4}$$

U. Pen





Pirsa: 13010114 Page 9/23



Reconstruction Data

Tidal Shear





$$\gamma_1 = T_{xx} - T_{yy}, \ \gamma_2 = 2T_{xy}$$



Page 10/23


Pirsa: 13010114

U. Pen




Reconstruction Data

Tidal Convergence Reconstruction




 $\kappa = \nabla^{-2}(\gamma_1,_{\mathsf{xx}} - \gamma_1,_{\mathsf{yy}} + 2\gamma_2,_{\mathsf{xy}}), \ \overline{\kappa_b} = \nabla^{-2}(\gamma_2,_{\mathsf{xx}} - \gamma_2,_{\mathsf{yy}} - 2\gamma_1,_{\mathsf{xy}})$


U. Pen



Pirsa: 13010114 Page 12/23



Pirsa: 13010114 Page 13/23



Pirsa: 13010114 Page 14/23

Reconstruction Data

Real Data: SDSS




U. Pen

Beating Cosmic Variance with Cosmic Tides


Pirsa: 13010114 Page 15/23

Reconstruction Data

Cosmic Convergence



Pirsa: 13010114 Page 16/23



Pirsa: 13010114 Page 17/23

Reconstruction Data

Cosmic Power dilemma

- ightharpoonup Power spectrum theoretically cleanest at low k
- ▶ statistics best at high *k*
- ► Cosmic variance limit?

U. Pen

Beating Cosmic Variance with Cosmic Tides

Pirsa: 13010114 Page 18/23

Future

Overcoming Cosmic Variance?

- ightharpoonup measure something better than $1/\sqrt{N}$
- c.f. Mcdonald and Seljak (2009)
- Goal is to measure transfer function
- ▶ could be measured better than P(k) if both density and velocity are measured on the same mode
- RSD measure different modes
- Cosmic tides vs redshift space measure growth factor on same modes!
- can use large wavelength linear modes.



U. Pen


Future

What the future may bring

- ▶ Use full SDSS catalog: 10x more galaxies
- other surveys: 6dF, 2dF, wiggleZ, etc
- ► Future surveys: BOSS, CHIME, etc
- Measure scale and time evolution of transfer function: neutrino masses, modified gravity?
- change observing strategies? Smaller volumes, closer by, all-sky, deeper?



U. Pen



Pirsa: 13010114 Page 21/23

Heuristic kernel: CMB Lensing

Gradient of the density field convolved by a Gaussian window.

$$W = \exp\left(\frac{-r^2}{2\sigma^2}\right) \tag{1}$$

then we obtain a windowed gradient density field

$$\bar{\delta} = \int W \delta. \tag{2}$$

Take log to Gaussianize: $\delta_g = \log(1+\bar{\delta})$

The tidal shear field is the quadratic outer product

$$T_{ij} = (\partial_i \delta_g)(\partial_j \delta_g). \tag{3}$$

The density field is derived from a convolution:

$$\langle \delta \rangle(x) = \int T_{ij}(x') K_{ij}(x - x') d^3 x'$$
 (4)

U. Pen

Heuristic kernel: CMB Lensing

Gradient of the density field convolved by a Gaussian window.

$$W = \exp\left(\frac{-r^2}{2\sigma^2}\right) \tag{1}$$

then we obtain a windowed gradient density field

$$\bar{\delta} = \int W \delta. \tag{2}$$

Take log to Gaussianize: $\delta_g = \log(1+\bar{\delta})$

The tidal shear field is the quadratic outer product

$$T_{ij} = (\partial_i \delta_g)(\partial_j \delta_g). \tag{3}$$

The density field is derived from a convolution:

$$\langle \delta \rangle(x) = \int T_{ij}(x') K_{ij}(x - x') d^3 x' \tag{4}$$

U. Pen