

Title: Quoins versus Coins

Date: Mar 27, 2013 02:00 PM

URL: <http://pirsa.org/13030000>


Abstract: Human monkeys are used to thinking about the problem of choosing from a set of objects according to some desired, biased, probability distribution. Just think about how you chose your partner(s). Even when it is easy for you to do such a sampling, it can be difficult to do a quantum sampling (Q-Sampling) of the same distribution. By Q-Sampling I mean the creation of a coherent superposition of states of such objects whose amplitudes are the (square roots of) of the specified distribution. In this talk I will discuss what we do know about this problem, why it is interesting, and will discuss how it can let us achieve tasks with quantum information that are provably impossible classically.

?


$$\rho = \sum_i p_i |i\rangle\langle i|$$

vs

$$|\psi\rangle = \sum_i \sqrt{p_i} |i\rangle$$



$\text{size}(\lambda) \ll \text{size}(|\psi\rangle)?$



$\text{size}(\lambda) \ll \text{size}(|\psi\rangle)?$

$(H) (T) (T) (H) \longrightarrow n \text{ bits}$


$\uparrow \swarrow \searrow \downarrow \longrightarrow n \text{ bits}$

$\exists$ HVT such that

$$\text{size}(\lambda) = O(n)$$

$$\Pr(\Pi | \psi) = C_0 + C_1 \langle \psi | \Pi | \psi \rangle$$

$$\left. \begin{array}{ccc} \textcircled{T} & \textcircled{H} & \textcircled{T} \\ \textcircled{H} & \textcircled{T} & \textcircled{T} \end{array} \right\} \Pr(H) = p$$




$\textcircled{H}$

$$\Pr(H) = f(p)$$

BERNOULLI FACTORY

"Exact monte-carlo" community

$$\left. \begin{array}{ccc} \textcircled{T} & \textcircled{H} & \textcircled{T} \\ \textcircled{H} & \textcircled{T} & \textcircled{T} \end{array} \right\} \Pr(H) = p$$




$$\textcircled{H}$$

$$\Pr(H) = f(p)$$


BERNOULLI FACTORY

"Exact monte-carlo" community



BERNOULLI FACTORY

"Exact monte-carlo" community



E.g. $f(p) = \sum_{k=0}^n C_{nk} p^k (1-p)^{n-k}$

$\uparrow$
integer $\leq \binom{n}{k}$

E.g. $f(p) = \frac{p^2}{p^2 + (1-p)^2}$

E.g. $f(p) = \frac{p^2}{p^2 + (1-p)^2}$


E.g. $f(p) = \frac{1}{2}$

E.g. $f(p) = \frac{1}{2}$


E.g. $f(p) = \frac{1}{2}$

$k, n=2$	$A_{2,k}$	$B_{4,k}$	$C_{2,k}$
0			00
1	10	01	
2			11
$k, n=4$	$A_{4,k}$	$B_{4,k}$	$C_{4,k}$
0			0000
1	0010, 1000	0001, 0100	
2	0011, 1010, 1001	0101, 0110, 1100	
3	1110, 1011	1101, 0111	
4			1111


E.g. $f(p) = \frac{1}{\pi} = 0.318309\dots$
 $= 0.01010001011$



E.g. $f(p) = \frac{1}{\pi} = 0.318309\dots$
 $= 0.01010001011$




$$f(p) = 2p \quad p \in (0, \frac{1}{2}]$$



$$f(p) = 2p$$

$$p \in (0, \frac{1}{2}]$$



THEOREM

(Kearney, O'Brien, '94)


$f(p)$ can be 'manufactured'
iff f continuous & either
 f is constant, OR

$\forall p, \exists k$ such that:

$$\min(p, 1-p) \geq \min(f(p)^k, (1-f(p))^k)$$

$$f(p) = 2p$$

$$p \in (0, \frac{1}{2}]$$




THEOREM

(Keane, O'Brien, '94)

$f(p)$ can be 'manufactured'
iff f continuous & either
 f is constant, OR

$\forall p, \exists k$ such that:

$$\min(p, 1-p) \geq \min(f(p)^k, (1-f(p))^k)$$



INTRODUCING QUOINS:

④

$$p|0\rangle\langle 0| + (1-p)|1\rangle\langle 1|$$



$$\sqrt{p}|0\rangle + \sqrt{1-p}|1\rangle$$

INTRODUCING QUOINS:

④

$$p|0\rangle\langle 0| + (1-p)|1\rangle\langle 1|$$



$$\sqrt{p}|0\rangle + \sqrt{1-p}|1\rangle$$

Q-Sampling



$$\{P_i, \Omega_i\}$$

$$\Omega_1 = \text{Hugo} = 01001$$

$$\Omega_2 = \text{David} = 10110$$

$$\Omega_3 = \text{Mike} = 01101$$

$$\sqrt{\frac{1}{2}}|01001\rangle + \sqrt{\frac{1}{6}}|10110\rangle + \sqrt{\frac{1}{3}}|01101\rangle \quad ?$$

Q-Sampling



$$\{P_i, \Omega_i\}$$

$$\Omega_1 = \text{Hugo} = 01001$$

$$\Omega_2 = \text{David} = 10110$$

$$\Omega_3 = \text{Mike} = 01101$$

$$\sqrt{\frac{1}{2}}|01001\rangle + \sqrt{\frac{1}{6}}|10110\rangle + \sqrt{\frac{1}{3}}|01101\rangle \quad ?$$

E.G. PERMUTATIONS OF $1, 2, \dots, n$

$$\Omega_1 = \sigma_1 = 123$$

$$\Omega_2 = \sigma_2 = 132$$

$$\vdots \quad \vdots \quad \vdots$$

$$\left\{ p_i = \frac{1}{n!}, \Omega_i = \sigma_i \right\}$$

Easy classically

ONE METHOD

- Choose $i \in \{1, \dots, n!\}$ uniformly
- Compute σ_i

$$\frac{1}{\sqrt{n!}} \sum_i |i\rangle |\sigma_i\rangle$$

$$\frac{1}{\sqrt{n!}} \sum_i |\sigma_i\rangle$$

E.G. PERMUTATIONS OF $1, 2, \dots, n$

$$\Omega_1 = \sigma_1 = 123$$

$$\Omega_2 = \sigma_2 = 132$$

$$\vdots \quad \vdots \quad \vdots$$

$$\{ p_i = \frac{1}{n!}, \Omega_i = \sigma_i \}$$

Easy classically

ONE METHOD


- Choose $i \in \{1, \dots, n!\}$ uniformly
- Compute σ_i

~~$$\frac{1}{\sqrt{n!}} \sum_i |i\rangle |\sigma_i\rangle$$~~

$$\frac{1}{\sqrt{n!}} \sum_i |\sigma_i\rangle$$

Q-SAMPLE

Q-SAMPLING & GRAPH ISOMORPHISM



$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$\sigma_0(G_1) = 101$$



$$\begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

$$\sigma_0(G_2) = 110$$

Q-SAMPLING & GRAPH ISOMORPHISM



$$\begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$



$$\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$$

$$\sigma_0(G_1) = 101$$

$$\sigma_1(G_1) = 101$$



$$\sigma_2(G_1) = 110$$

$$\sigma_0(G_2) = 110$$

$$\sigma_1(G_2) = 011$$

$$\sigma_2(G_2) = 011$$

← SAME




CLASSICAL
(EASY)

$$\{ p_i = \frac{1}{n!}, \Omega_i = \sigma_i(q) \}$$

POO SAMPLE
(EASY)

$$\frac{1}{\sqrt{n!}} \sum_i |i\rangle |\sigma_i(q)\rangle$$

SWAP TEST



DOES NOT WORK FOR POO-SAMPLES

$$\left(\sum_i \langle i | \langle \sigma_i(q_1) | \right) \left(\sum_j | j \rangle | \sigma_j(q_2) \rangle \right) = 0$$

Even if $G_1 = G_2$

WHAT CAN BE Q-SAMPLED?

ALL BIT STRINGS

$P_0 \dots 00$	$0 \dots 00$	
$P_0 \dots 01$	$0 \dots 01$	
$P_0 \dots 10$	$0 \dots 10$	P_1, Ω_1
$\vdots$	$\vdots$	

WHAT CAN BE Q-SAMPLED?

RECENTLY DISCOVERED POWERFUL CLASSES OF CLASSICAL SAMPLING ALGORITHMS SOLVING HARD PROBLEMS:

- Ferromagnetic Ising (Jerrum, Sinclair)
- Permanents of non-negative matrices (Jerrum, Sinclair and Vigoda)
- Volumes of convex bodies in high dimensions (Dyer, Frieze, Kannan)
- Integration of log-concave functions (Applegate, Kannan)

"Self Reducible"

[arXiv.org](#) > [quant-ph](#) > [arXiv:quant-ph/0208112](#)

Quantum Physics

Creating superpositions that correspond to efficiently integrable probability distributions

[Lov Grover](#), [Terry Rudolph](#)

WHAT CAN BE Q-SAMPLED?

$n=2$
example

$$\sqrt{p_{00}+p_{01}}|0\rangle + \sqrt{p_{10}+p_{11}}|1\rangle$$

$$\sqrt{p_{00}}|00\rangle + \sqrt{p_{01}}|01\rangle + \sqrt{p_{10}}|10\rangle + \sqrt{p_{11}}|11\rangle$$

WHAT CAN BE Q-SAMPLED?

$n=2$
example

$$\sqrt{p_{00}+p_{01}}|0\rangle + \sqrt{p_{10}+p_{11}}|1\rangle$$

$$\rightarrow \sqrt{p_{00}+p_{01}}|0\rangle \left| \frac{p_{00}}{p_{00}+p_{01}} \right\rangle$$

$$+ \sqrt{p_{10}+p_{11}}|1\rangle \left| \frac{p_{10}}{p_{10}+p_{11}} \right\rangle$$

$$\sqrt{p_{00}}|00\rangle + \sqrt{p_{01}}|01\rangle + \sqrt{p_{10}}|10\rangle + \sqrt{p_{11}}|11\rangle$$

WHAT CAN BE Q-SAMPLED?

arXiv.org > quant-ph > arXiv:quant-ph/0301023

Quantum Physics

Adiabatic Quantum State Generation and Statistical Zero Knowledge

[Dorit Aharonov](#), [Amnon Ta-Shma](#)

- (Only?) Results potentially applicable to non-self-reducible problems.


THEOREM

(Kearney, O'Brien, '94)

$f(p)$ can be 'manufactured'
iff f is continuous & either
 f is constant, OR

$\forall p, \exists k$ such that:

$$\min(p, 1-p) \geq \min(f(p)^k, (1-f(p))^k)$$




THEOREM

(Keane, O'Brien, '94)

$f(p)$ can be 'manufactured'
iff f continuous & either
 f is constant, OR
 $\forall p, \exists k$ such that:

$$\min(p, 1-p) \geq \min(f(p)^k, (1-f(p))^k)$$



$$\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \sqrt{p} \\ \sqrt{1-p} \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} \sqrt{p} + \sqrt{1-p} \\ \sqrt{p} - \sqrt{1-p} \end{bmatrix}$$

QUOINS INFINITELY MORE
POWERFUL THAN CANS!!

MUCH MORE FUN TO BE HAD

• What functions can Quoins sample?

MUCH MORE FUN TO BE HAD

- What functions can Quoins sample?
- Quoinage savings for $f=2p$?
- Joint measurements more powerful?
- Doubly-Quantum Bernstein factory?


$$(\sqrt{p}|0\rangle + \sqrt{1-p}|1\rangle)^{\otimes N} \rightarrow \sqrt{f(p)}|0\rangle + \sqrt{1-f(p)}|1\rangle$$

SOME LAST FUN WITH Q-SAMPLING

- Why does computation take time?
circuits, adiabatic, cluster...

SOME LAST FUN WITH Q-SAMPLING

- Why does computation take time?
circuits, adiabatic, cluster...



o Q-Sample all measurement records that
output $| \Omega \rangle$ [NOT $X| \Omega \rangle, Z| \Omega \rangle$ etc]

$$| \mu \rangle = | 000 \dots 0 \rangle + \dots | 001101 \dots \rangle + \dots \quad O(2^n) \text{ terms}$$

$$| \mu \rangle \Rightarrow \dots + \dots | \alpha_+ \beta_+ - \gamma_- \delta_- \dots \rangle + \dots$$


($\Pi_i = | \phi_i \rangle \langle \phi_i |$, $\sum_i | \phi_i \rangle$)

o Q-Sample all measurement records that
output $| \Omega \rangle$ [NOT $X| \Omega \rangle$, $Z| \Omega \rangle$ etc]

$$| \mu \rangle = | 000 \dots 0 \rangle + \dots | 001101 \dots \rangle + \dots \quad O(2^n) \text{ terms}$$

$$| \mu \rangle \Rightarrow \dots + \dots | \alpha + \beta + \gamma - \delta - \dots \rangle + \dots$$

$(\Pi_i = | \phi_i \rangle \langle \phi_i |, \sum_i | \phi_i \rangle)$




o Q-Sample all measurement records that output $| \Omega \rangle$ [NOT $X| \Omega \rangle, Z| \Omega \rangle$ etc]

$$| \mu \rangle = | 000 \dots 0 \rangle + \dots | 001101 \dots \rangle + \dots O(2^n) \text{ terms}$$

$$| \mu \rangle \Rightarrow \dots + \dots | \alpha_+ \beta_+ \gamma_- \delta_- \dots \rangle + \dots$$

$(\Pi_i = | \phi_i \rangle \langle \phi_i |, \sum_i | \phi_i \rangle)$



o Q-Sample all measurement records that output $| \Omega \rangle$ [NOT $X| \Omega \rangle, Z| \Omega \rangle$ etc]

$$| \mu \rangle = | 000 \dots 0 \rangle + \dots | 001101 \dots \rangle + \dots O(2^n) \text{ terms}$$

$$| \mu \rangle \Rightarrow \dots + \dots | \alpha + \beta + \gamma + \delta \dots \rangle + \dots$$

$(\Pi_i = | \phi_i \rangle \langle \phi_i |, \sum_i | \phi_i \rangle)$

