

Title: General Relativity for Cosmology - Lecture 18

Date: Nov 19, 2013 04:00 PM

URL: <http://pirsa.org/13110005>

Abstract:

→ Make symmetry assumptions.

□ Potential problem: (with symmetry assumptions):

(E.g. recall that flatness
in FL spacetimes is unstable)

□ The so-obtained highly symmetric solutions,
e.g. Friedman-Lemaître, may possess properties
that are peculiar to high symmetry.

□ E.g.: When a Friedmann-Lemaître solution,
or a Schwarzschild solution exhibits a
singularity: Is it due to symmetry, or realistic?

◻ Singularity theorems (see later) confirm the robustness under certain conditions (such as strong energy condition).

→ More confidence in significance of the properties of highly symmetric solutions.

Question: How much can we weaken the symmetry assumptions of Friedmann-Lemaitre and still get exact solutions?

Strategy:

◻ Classify cosmological models $(M, g), T_{\mu\nu}$ by the amount and type of symmetry assumed.

Strategy:

- Classify cosmological models $(M, g), T_{\mu\nu}$ by the amount and type of symmetry assumed.
- For each amount and type of symmetry assumed, try to find exact solutions or at least (asymptotic) properties of exact solutions.

Remark: Among the other high symmetry models, some come arbitrarily close to F.L. at finite times!
See, e.g., text by Wainwright & Ellis.

Remark: Among the other high symmetry models, some come arbitrarily close to F.L. at finite times!
See, e.g., text by Wainwright & Ellis.

Recall: Symmetries & Killing vector fields


- Two spacetimes (M, g) , (\tilde{M}, \tilde{g}) are isometric (and therefore of exactly identical shape) if there is a diffeomorphism $\phi: M \rightarrow \tilde{M}$ so that the image of the metric g in \tilde{M} is $\tilde{g}: Tg = \tilde{g}$.

Recall: Symmetries & Killing vector fields

□ Two spacetimes (M, g) , (\tilde{M}, \tilde{g}) are isometric (and therefore of exactly identical shape) if there is a diffeomorphism $\phi: M \rightarrow \tilde{M}$ so that the image of the metric g in \tilde{M} is $\tilde{g}: Tg = \tilde{g}$.

□ A space-time has a symmetry, if we find such a ϕ for $\tilde{M} = M$.

□ Example:




ϕ performs a rotation of M about a symmetry axis, to obtain $\tilde{M} = M$

□ Two spacetimes (M, g) , (\tilde{M}, \tilde{g}) are isometric (and therefore of exactly identical shape) if there is a diffeomorphism $\phi: M \rightarrow \tilde{M}$ so that the image of the metric g in \tilde{M} is $\tilde{g}: Tg = \tilde{g}$.

□ A space-time has a symmetry, if we find such a ϕ for $\tilde{M} = M$.

□ Example:



ϕ performs a rotation of M about a symmetry axis, to obtain $\tilde{M} = M$ with $Tg = \tilde{g}$.

Note: The set of all symmetries of a manifold (M, g) forms a "group":

Definition: A "group" G is a set, with an operation, say " \circ ",
 $\circ : G \times G \rightarrow G$

and a "neutral element", say " e ", $e \in G$, such that

$$(a \circ b) \circ c = a \circ (b \circ c) \quad \forall a, b, c \in G$$

$$a \circ e = e \circ a = a \quad \forall a \in G$$

$$\exists a^{-1}: a^{-1} \circ a = a \circ a^{-1} = e \quad \begin{matrix} \uparrow \text{"there exists"} \\ \forall a \in G \end{matrix} \quad \begin{matrix} \\ \leftarrow \text{"for all"} \end{matrix}$$

Definition: A group G is called a Lie group if G is also a finite-dimensional smooth manifold.

Definition: A group G is called a Lie group if G is also a finite-dimensional smooth manifold.

Example: The sets of rotations in \mathbb{R}^3 forms a 3-dimensional Lie group, $SO(3)$.

The angles



α, β, γ are coordinates for elements $g \in SO(3)$.

- Remarks:
- The symmetries of a manifold (M, g) can be discrete, such as reflections.
 - But often, the symmetry group of a manifold (M, g) is actually a Lie group.

Remarks: □ The symmetries of a manifold (M, g) can be discrete, such as reflections.

□ But often, the symmetry group of a manifold (M, g) is actually a Lie group.

Note: □ Each $g \in G$ yields an isometric diffeomorphism:

$$g : M \rightarrow M, \text{ namely } g : p \rightarrow g(p) \quad \forall p \in M$$

□ Consider the set $O_p \subset M$ defined by: $O_p := \{q \in M \mid \exists g \in G : g(p) = q\}$

Definition: The set O_p is called the Orbit of p under the action of the group G .

Note: If G is a Lie group then each orbit O_p is $\overset{\text{hand}}{\text{a}}$ submanifold of (M, g) .

□ Consider the set $O_p \subset M$ defined by: $O_p := \{g \cdot p \mid g \in G\}$

Definition: The set O_p is called the *Orbit* of p under the action of the group G .

Note: If G is a Lie group then each orbit O_p is p or a submanifold of (M, g) .

Question: What are the *infinitesimal isometric diffeomorphisms*?

And if not a Lie group, what structure do the infinitesimal symmetries form?

□ Recall: The Lie derivative,

$$\begin{aligned} L_{\xi} Q^{a \dots b}_{\quad c \dots d} &= Q^{a \dots b}_{\quad c \dots d; k} \xi^k \\ &\quad - Q^{k \dots b}_{\quad c \dots d} \xi^a_{; k} - \dots - Q^{a \dots k}_{\quad c \dots d} \xi^b_{; k} \end{aligned}$$

◻ Recall: The Lie derivative,

$$\begin{aligned} L_{\xi} Q^{a \dots b}_{c \dots d} &= Q^{a \dots b}_{c \dots d j k} \xi^j \xi^k \\ &\quad - Q^{k \dots b}_{c \dots d} \xi^a_{j k} - \dots - Q^{a \dots k}_{c \dots d} \xi^b_{j k} \\ &\quad + Q^{a \dots b}_{k \dots d} \xi^k_{j c} + \dots + Q^{a \dots b}_{c \dots k} \xi^k_{j d} \end{aligned}$$

yields the rate of change of a tensor Q along the flow of diffeomorphisms ϕ generated by a vector field ξ .

→ Here, can use L_{ξ} to differentiate along symmetry group orbits.

◻ Thus, if $L_{\xi} g_{\mu\nu} = 0$

$$+ Q^{\alpha\beta}_{\kappa\lambda} \xi^\kappa_{;\mu} + \dots + Q^{\alpha\beta}_{\epsilon\mu} \xi^\epsilon_{;\mu}$$

yields the rate of change of a tensor Q along the flow of diffeomorphisms ϕ generated by a vector field ξ .

→ Here, can use L_ξ to differentiate along symmetry group orbits.

□ Thus, if $L_\xi g_{\mu\nu} = 0$

then ξ generates isometries $\phi: M \rightarrow M, g \rightarrow \tilde{g} = g$.

□ But $L_\xi g_{\mu\nu} = \xi^\kappa \underbrace{g_{\mu\nu;\kappa}}_{\text{always } f \in \Gamma, g \text{ compatibility}} + g_{\kappa\nu} \xi^\kappa_{;\mu} + g_{\mu\kappa} \xi^\kappa_{;\nu}$

◻ But $\mathcal{L}_\xi g_{\mu\nu} = \xi^k \underbrace{g_{\mu\nu;jk}}_{\text{always for } \Gamma, g \text{ compatibility}} + g_{k\nu} \xi^k_{;\mu} + g_{\mu k} \xi^k_{;\nu}$

i.e. is itself an infinitesimal symmetry

\Rightarrow A vector field ξ generates a symmetry of spacetime if it is a Killing vector field:

$$\xi_{\mu;\nu} + \xi_{\nu;\mu} = 0 \quad (\text{X})$$

Q: Maximum number, d , of Killing vector fields in n dims?

A: $d = n(n+1)/2$ To see this, note that there are 2 ways to obey Eq. (X):

a) $\xi_{\mu;\nu} = 0$ i.e. $\nabla \xi = 0$

I can have maximally n such index vectors

◻ But $\mathcal{L}_\xi g_{\mu\nu} = \xi^k \underbrace{g_{\mu\nu;jk}}_{\text{always for } \Gamma, g \text{ compatibility}} + g_{k\nu} \xi^k_{;\mu} + g_{\mu k} \xi^k_{;\nu}$

i.e. is itself an infinitesimal symmetry

\Rightarrow A vector field ξ generates a symmetry of spacetime if it is a Killing vector field:

$$\xi_{\mu;\nu} + \xi_{\nu;\mu} = 0 \quad (\text{X})$$

Q: Maximum number, d , of Killing vector fields in n dims?

A: $d = n(n+1)/2$ To see this, note that there are 2 ways to obey Eq. (X):

a) $\xi_{\mu;\nu} = 0$ i.e. $\nabla \xi = 0$

I can have maximally n such index vectors

$$\xi_{\mu;\nu} + \xi_{\nu;\mu} = 0 \quad (\text{X})$$

Q: Maximum number, d , of Killing vector fields in n dims?

A: $d = n(n+1)/2$ To see this, note that there are 2 ways to obey Eq. (X):

a) $\xi_{\mu;\nu} = 0$ i.e. $\nabla g = 0$

(can have maximally n such indep. vectors)

b) $\nabla g \neq 0$, but then $K_{\mu\nu} := \xi_{\mu;\nu}$ is antisymmetric

(can have at most $n(n-1)/2$ indep. such cases.)

$\Rightarrow d = n + \frac{n(n-1)}{2} = n(n+1)/2$

From a symmetry Lie group to a "symmetry Lie algebra":

From a symmetry Lie group to a "symmetry Lie algebra":

General idea:

Normally the points of a manifold cannot be multiplied!

- A Lie group is a smooth manifold with extra structure: the multiplication.
- Notice: Product of group elements close to $1 \in G$ yields a group element close to 1.
- Consider the tangent space $T_1(M)$ to the point $1 \in M$ of the Lie group manifold M .
- $T_1(M)$ is a vector space and it has extra structure, inherited from the group's multiplication.
- Define the Lie algebra of a group M to be $T_1(M)$, equipped with the inherited "multiplication".

Identity element of the group: $p = 1$

- Consider the tangent space $T_1(M)$ to the point 1_M of the Lie group manifold M .
- $T_1(M)$ is a vector space and it has extra structure, inherited from the group's multiplication.
- Define the Lie algebra of a group M to be $\overset{\uparrow}{T_1(M)}$, equipped with the inherited "multiplication".
Identity element of the group: $p = 1$

Crucial fact: From knowledge of only the Lie algebra, i.e., only $T_1(M)$ and its "multiplication", the group M can be constructed!
(though not always uniquely)



- Let us collect the properties that the inherited multiplications of all Lie algebras share.
- Then, let us define Lie algebras as anything with these properties.

- Let us collect the properties that the intended multiplications of all Lie algebras share.
- Then, let us define Lie algebras as anything with these properties:

Definition:

A Lie algebra is a vector space A , with an operation $\{, \}$

$$\{, \} : A \times A \rightarrow A \quad \text{"Lie bracket"}$$

obeying $\{r, s\} = -\{s, r\} \quad \forall r, s \in A$

"Jacobi identity"

and $\{\{r, s\}, t\} + \{\{t, r\}, s\} + \{\{s, t\}, r\} = 0$

Theorem: Every vector space A with a "multiplication" $\{, \}$ that

Definition:

A Lie algebra is a vector space A , with an operation $\{, \}$

$$\{, \} : A \times A \rightarrow A \quad \text{"Lie bracket"}$$

obeying $\{r, s\} = -\{s, r\} \quad \forall r, s \in A$
 "Jacobi identity"

and $\{\{r, s\}, t\} + \{\{t, r\}, s\} + \{\{s, t\}, r\} = 0$

Theorem: Every vector space A with a "multiplication" $\{, \}$ that obeys these axioms is isomorphic to $T_1(M)$ of a Lie group M .

Proposition: The set of Killing vector fields $\xi^{(i)}$ of (M, g) is a Lie algebra.

Exercise: Prove this, i.e., show the following:

Assume $\xi^{(1)}, \xi^{(2)}$ are Killing vector fields of (M, g) and $\alpha, \beta \in \mathbb{R}$.

Then:

$$\alpha \xi^{(1)} + \beta \xi^{(2)}$$

(i.e., they form a vector space)

and $\{\xi^{(1)}, \xi^{(2)}\} := \xi^{(1)}\xi^{(2)} - \xi^{(2)}\xi^{(1)}$

are also Killing vector fields,

Exercise: Prove this, i.e., show the following:

Assume $\xi^{(1)}, \xi^{(2)}$ are Killing vector fields of (M, g) and $\lambda, \mu \in \mathbb{R}$.

Then:

$$\lambda \xi^{(1)} + \mu \xi^{(2)}$$

(i.e., they form a vector space)

and $\{\xi^{(1)}, \xi^{(2)}\} := \xi^{(1)}\xi^{(2)} - \xi^{(2)}\xi^{(1)}$

are also Killing vector fields,

and the $\xi^{(i)}$ obey the Jacobi identity.

Summary of the big picture:

1. The symmetries of any (M, g) form a group:
they can be concatenated associatively, and all
possess an inverse. Some symmetries are differentiable,
parametrized by the flow \Rightarrow the symmetries
form a Lie group.

2. Each Killing vector field is the infinitesimal
generator of a flow of isometric diffeomorphisms,
i.e., of a symmetry.
3. We see here that the Killing vector fields indeed form a Lie algebra.

they can be concatenated associatively, and all possess an inverse. Some symmetries are differentiable, parameterized by the flow \Rightarrow the symmetries form a Lie group.

↓ Recall: there can be discrete symmetries too.

2. Each Killing vector field is the infinitesimal generator of a flow of isometric diffeomorphisms, i.e., of a symmetry.
3. We see here that the Killing vector fields indeed form a Lie algebra.
4. Recall that every Lie algebra generates a Lie group.

Surfaces of homogeneity and the isotropy subgroup:

□ Definition:

Let r be the dimension of the Lie algebra, i.e., also the dimension of the Lie group of symmetries.

□ Recall this definition:

□ Consider the set of points $O(p)$ that a point p can flow to along the killing vector fields.

□ $O(p)$ is called the orbit of $p \in M$ under the action of the symmetries around. We denote the dimension of

□ Clearly:

The dimension of an orbit cannot be larger than the dimension of the symmetry group, i. e.

$$s \leq r,$$

but $s < r$ easily happens.

□ Example:

□ Consider $M := \mathbb{R}^2$ and $p = (0, 0)$.

□ Then $r = r_{\max} = \overbrace{n(n+1)/2}^{n=2} = \underline{\underline{3}}$ is dim. of sym. group.

□ Then $\tau = \tau_{\max} = \sqrt{n(n+1)}/2 = \underline{\underline{3}}$ is dim. of sym. group.

□ \Rightarrow The three-dimensional Lie algebra of Killing vector fields is spanned by three Killing vector fields:

□ Concretely:

$$K^{(1)} := \frac{\partial}{\partial x}, \quad K^{(2)} := \frac{\partial}{\partial y}$$

$$K^{(3)} := y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y}$$

□ Orbit of $p = (0,0)$:

$$\mathcal{O}(p) = \mathbb{R}^2 \text{ because generators } \underline{\underline{2}} \text{ and } \underline{\underline{2}}$$

 Group elements generated by them are $e^{\frac{a^2}{2x} + b^2}{\partial}{\partial}$ and they act as $e^{\frac{a^2}{2x} + b^2}{\partial}{\partial} f(x,y) = f(x+a, y+b)$ by Taylor expansion.

□ Concretely:

$$K^{(1)} := \frac{\partial}{\partial x}, \quad K^{(2)} := \frac{\partial}{\partial y}$$

$$K^{(3)} := y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y}$$

□ Orbit of $p = (0,0)$:

$\left(\begin{array}{l} \text{Group elements generated by them} \\ \text{are } e^{\frac{a}{2}x + \frac{b}{2}y} \text{ and they act as} \\ e^{\frac{a}{2}x + \frac{b}{2}y} f(x,y) = f(x+a, y+b) \\ \text{by Taylor expansion.} \end{array} \right)$

$O(p) = \mathbb{R}^2$ because generators $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ generate flow to every where.

Def: The surface of homogeneity has dimension $s = 2 < \infty$

□ Notice:

↑ generated by the Killing vectors (here: $K^{(1)}, K^{(2)}$) which do not have trivial orbits

Since $n=2$, at any given point p , only

□ Rôle of $K^{(3)}$?

$K^{(3)}$ is the angular momentum
and it of course generates rotations:
 $e^{tK^{(3)}} f(x,y) = f(x \cos t - y \sin t, x \sin t + y \cos t)$

The flow generated by $K^{(3)}$ leaves p fixed
and rotates everything around p .

□ Definition:

We say that those Killing vector fields
which do not generate a homogeneity surface,
i.e., which generate a trivial group orbit for a point
are generating the isotropy subgroup (of the
full symmetry group generated by all Killing vectors).

□ Dimension, d , of the isotropy subgroup?

Classification of cosmological models

□ The classification is with respect to:

□ Dimension of isotropy subgroup d :

(# of conserved angular momenta) →

$d = 0, 1, 2, 3, 4, 5, 6$

↑
at each p.m.
one rotational
symmetry axis

anisotropic case

e.g. full Lorentz symmetry

e.g. spatially,
isotropic case

□ Dimension of homogeneity surfaces s :

(# of conserved momenta) →

$s = 0, 1, 2, 3, 4$

□ A large body of literature exists on most cases of (d, s) :

- Many exact solutions are known !
- Many asymptotic behaviors are known !

□ Comprehensive text:

Wainwright & Ellis, Dyn. systems in cosmology,
Cambridge Univ. Press (1997)

□ Examples: homogeneity isotropy

□

\downarrow
s \downarrow
d



4 3
4 1
4 0

Einstein's static model
Gödel's model
Brans - Dicke model

18

□ Examples: homogeneity ↓ isotropy
 □ S d

4	3	Einstein's static model
4	1	Bödel's model
4	0	Oszváth-Kas models
3	3	Friedmann-Lemaitre models
3	1	spatially hom & locally one rot. sym axis
3	0	Bianchi models
:	:	

Powerful alternative classification approach:

Idea: Classify the possible $T_{\mu\nu}$, then use Einstein equation to obtain classification of curvature.

Powerful alternative classification approach:

Idea: Classify the possible $T_{\mu\nu}$, then use Einstein equation to obtain classification of curvature.

Proposition:



For every physical energy momentum tensor $T_{\mu\nu}$ there exists a unique timelike vector field u so that $T_{\mu\nu}$ takes this standard form:

$$T_{ab} = \mu u_a u_b + q_a u_b + q_b u_a + p(g_{ab} + u_a u_b) + \Pi_{ab}$$

scalar
vector
scalar
tensor

now. using / the previous expression we can obtain
equation to obtain classification of curvature.

Proposition:

For every physical energy momentum tensor $T_{\mu\nu}$ there exists a unique timelike vector field u so that $T_{\mu\nu}$ takes this standard form:

$$T_{ab} = \mu u_a u_b + q_a u_b + q_b u_a + p(g_{ab} + u_a u_b) + \pi_{ab}$$

where q and π are a vector field and a tensor field obeying:

$$q_a u^a = 0, \quad \pi_{ab} u^b = 0, \quad \pi_a^a = 0, \quad \pi_{ab} = \pi_{ba}$$

Definition: u is called the "fundamental 4-velocity field"

Note:


E.g., for a perfect fluid this is the fluid velocity:

$$T_{ab} = \mu u_a u_b + p(g_{ab} + u_a u_b), \quad u_a u^a = -1$$

Recall: equation of state is

$$p = \tilde{\mu} (\tilde{\epsilon} - 1) \mu$$

$$\tilde{\epsilon} = \begin{cases} 1 & \text{dust} \\ 4/3 & \text{radiation} \\ 0 & \text{cosmological constant} \end{cases}$$



□ Definition:

If (M, g) possesses spacelike $s=3$ homogeneity

but the fundamental 4-velocity is not orthogonal to

□ Definition:

If (M, g) possesses spacelike $s=3$ homogeneity but the fundamental velocity is not orthogonal to the homogeneity surfaces, then we say that this cosmology is "tilted".

Segré classification:

- A systematic classification of $T_{\mu\nu}$ can be performed, by the analysis of its eigenvalues / eigenvectors. Nontrivial because:
 - $T_{\mu\nu}$ is symmetric.

Segré classification:

□ A systematic classification of $T_{\mu\nu}$ can be performed, by the analysis of its eigenvalues / eigenvectors. Nontrivial because:

□ $T_{\mu\nu}$ is symmetric.

But, the inner product in the vector space is $g_{\mu\nu} \Rightarrow T_{\mu\nu}$ is generally not hermitian!

□ T'_{ν} is in a space with the inner product $g'^{\nu}{}_{\nu} = \delta'^{\nu}_{\nu}$, but T'^{ν}_{ν} is generally not symmetric!

Use Jordan normal form:

Recall strategy:

The classification of possible $T_{\mu\nu}$ should, via the Einstein eqns, yield a classification of possible curvatures.

Indeed: In 3+1 dimensions the Einstein equation also reads:

$$R_{\mu\nu} = 8\pi G \left(T_{\mu\nu} - \frac{1}{2} g_{\mu\nu} T \right)$$

Exercise: Prove this and notice the dimension-dependence

⇒ The 10 degrees of freedom of $T_{\mu\nu}$ has a constraint

⇒ The Segré classification of possible $T_{\mu\nu}$ yields, via the Einstein equation also a classification of possible Ricci tensors $R_{\mu\nu}$.

Q: Does this yield also a classification of the possible Riemann tensors $R^{\mu\nu\rho\sigma}$?

A: No! The Ricci tensor contains only 10 of the 20 degrees of freedom of the Riemann tensor! (In 3+1 dim)



Prop.: The information in $R^{\mu\nu\rho\sigma}$ is shared among the Ricci tensor $R_{\mu\nu}$ and the so-called Weyl tensor, $C^{\mu\nu\rho\sigma}$

A: No! The Ricci tensor contains only 10 of the 20 degrees of freedom of the Riemann tensor! (In 3+1 dim)

Prop.: The information in $R^{\mu}_{\nu\alpha\beta}$ is shared among the Ricci tensor $R_{\mu\nu}$ and the so-called Weyl tensor, $C^{\mu}_{\nu\alpha\beta}$.

⇒ It remains to classify the possible Weyl tensors!

The Weyl tensor, C^{am}_{sq} :

$$C^{am}_{sq} := R^{am}_{sq} - \frac{1}{2} (g_s^a R^m_{\quad q} + g^m_{\quad q} R^a_{\quad s} - g^m_s R^a_{\quad i} - g^a_i R^m_{\quad s}) + \frac{1}{6} (g^a_s g^m_q - g^a_q g^m_s) R$$

The Weyl tensor, C^{am}_{sq} :

$$C^{am}_{sq} := R^{am}_{sq} - \frac{1}{2} (g_s^a R_q^m + g_q^m R_s^a - g_s^m R_a^q - g_a^q R_s^m) + \frac{1}{6} (g_s^a g_q^m - g_a^q g_s^m) R$$

Notice: If R^a_b and C^{am}_{sq} are given, they determine R^{am}_{sq} fully:

$$R^{am}_{sq} = C^{am}_{sq} + \frac{1}{2} (g_s^a R_q^m + g_q^m R_s^a - g_s^m R_a^q - g_a^q R_s^m) - \frac{1}{6} (g_s^a g_q^m - g_a^q g_s^m) R$$


→ R^{am}_{sq} is expressed through C^{am}_{sq} and R^{ab}

↑ 20 indep. components

↑ 10 indep. comp.

↑ 10 indep. comp.

$\Rightarrow C^m_{sq}$ contains all that curvature information which is not determined via the Einstein equation by $T_{\mu\nu}$.



C^m_{sq} describes all that curvature which can exist even where there is no matter! (e.g. gravity waves)

also e.g. sun's gravity away from the sun in empty space

Proposition

□ Assume (M, g) is a 3+1 dimensional Lorentzian manifold.

□ Choose any smooth positive scalar function ϕ on M .

□ Define (M, \tilde{g}) with the new metric \tilde{g} obtained through the "conformal transformation":

Intuition:

Weil curvature didn't change

Proposition

↳ also e.g. sun's gravity away from the sun in empty space

□ Assume (M, g) is a 3+1 dimensional Lorentzian manifold.

□ Choose any smooth positive scalar function ϕ on M .

□ Define (M, \tilde{g}) with the new metric \tilde{g} obtained through the "conformal transformation":

$$g_{\mu\nu}(x) \rightarrow \tilde{g}_{\mu\nu}(x) := \phi(x) g_{\mu\nu}(x)$$

Then: $\tilde{C}^{\nu\sigma\rho}(x) = C^{\nu\sigma\rho}(x) \quad \forall x \in M$ (Exercise: what would be a proof strategy?)

Intuition:

Weyl curvature distorts (0 0 0)
but only Ricci curvature
shrinks or expands overall: (+0 0)

Historical remark

□ Consider the equivalence class of spacetimes

Historical remark

- Consider the equivalence class of spacetimes (M, \tilde{g}) that are conformally equivalent to Minkowski space:

$$\tilde{g}_{\mu\nu}(x) = \phi^2(x) g_{\mu\nu}$$

- Einstein and Föhrer initially considered a theory in which the metric possesses only this conformal degree of freedom ϕ (to play role of Newton's gravitational potential):

Newton gravity
does come out
correctly as a
limiting case!
Newton gravity
does come out
correctly as a
limiting case!

Then, $S = \int_M \sqrt{-g} d^4x + \int_{\text{matter}} \sqrt{g} d^4x$ and $\frac{\delta S}{\delta g} = 0$
yielded:

$$R \equiv 8\pi G T_F$$

\rightarrow in electromagnetism $T^{(EM)}_{\mu\nu} = 0$
i.e. EM fields would not gravitate

26

No gravity waves

- Equivalence principle ok.

Recall: via the Einstein equation the Segré classification implies a classification of properties of the Ricci tensor $R_{\mu\nu}$.

It remains to classify the Weyl tensor:

Petrov classification:

This is a classification of the Weyl tensor $C^{\mu\nu}_{\gamma\delta}$, which possesses the 10 remaining degrees of freedom of $R^\lambda_{\mu\nu\delta}$.

- $C^{\mu\nu}_{\gamma\delta}$, just like the Riemann tensor, is antisymmetric in $\mu \leftrightarrow \nu$ and in $\gamma \leftrightarrow \delta$, and symmetric in $\mu \nu \leftrightarrow \gamma \delta$.

□ Thus $C^{\mu\nu} g_{\mu\nu}$ can locally be viewed as a symmetric map from the antisymmetric part $A_p(M)^2$ of $T_p(M)^2$ (so called bi-vectors) into itself:

$$C : A_p(M)^2 \rightarrow A_p(M)^2$$

□ But, the inner product in $A_p(M)^2$ is not positive definite!

$\Rightarrow C$ is generally not hermitian.
Therefore, use Jordan normal form again:

Result : 6 main Petrov classes for Weyl curvature:
according to eigenvalues/eigen-vector decompositions

Result :

6 main Petrov classes for Weyl curvature:
according to eigenvalue/eigenvector decomposition.

Type O: Weyl curvature vanishes

Type D: "Static" Weyl curvature, e.g., in
vicinity of a star.

Type N: Transverse gravitational waves, the
type LIGO aims to detect. Like light,
their strength decays $\sim \frac{1}{r}$ from the source.

Type I: Longitudinal gravitational waves

Type O: Weyl curvature vanishes

Type D: "Static" Weyl curvature, e.g., in vicinity of a star.

Type N: Transverse gravitational waves, the type LIGO aims to detect. Like light, their strength decays $\sim \frac{1}{r}$ from the source.

Type I: Longitudinal gravitational waves

These waves cause a shear effect.

However, they decay fast: $\sim \frac{1}{r^2}$

Type N: Transverse gravitational waves, the type LIGO aims to detect. Like light, their strength decays $\sim \frac{1}{r}$ from the source.

Type I: Longitudinal gravitational waves

These waves cause a shear effect.

However, they decay fast: $\sim \frac{1}{r^2}$

Why? Gravitational waves, when small enough, travel with speed of light. Like light, they then cannot oscillate longitudinally.

Types II, III: Mixtures of the above.