

Title: Quantum Turbulence

Date: May 28, 2014 10:30 AM

URL: <http://pirsa.org/14050104>

Abstract: This hour will be devoted to a description of quantum turbulence, that is turbulence in superfluids. The first talk (~20 minutes) will be given by Russell Donnelly. He will describe briefly the problem of classical turbulence and how turbulence in superfluids is different. The second talk will be given by Carlo Barenghi who will discuss progress in the simulation of quantum turbulence which is capable of suggesting insights so far inaccessible to experiment.


Quantum Turbulence




Quantum Turbulence

Russell J. Donnelly
Department of Physics
University of Oregon






Classical and Quantum Turbulence




Water flowing past a cylinder → turbulent wake

Flow past an aerofoil



Water flowing through a grid → homogeneous turbulence



- The flow of **water** is governed by **classical mechanics**.
- How is turbulence modified if the water is replaced by a **superfluid**?
(Remember flow of a **superfluid** is strongly influenced by **quantum effects**).
- How does quantum turbulence differ from classical turbulence?



UNIVERSITY OF OREGON

Turbulence is the chaotic motion of a fluid that occurs at sufficiently high flow rates. A common example is the flow of water from a tap: turn the tap on a little and the water runs out in a regular fashion called laminar flow. Turn the tap on fully and the flow becomes an irregular torrent. Such turbulent flow is characterized by regions of circulating fluid called eddies or vortices. If these features are ordered, they can lead to large scale structures such as whirlpools or tornados. Usually the eddies are highly irregular and interact in a complex and unpredictable manner.



Classical turbulence consists of eddies (Leonardo 1452-1519)



UNIVERSITY OF OREGON

Classical turbulence consists of eddies (Leonardo 1452-1519)



UNIVERSITY OF OREGON

Classical Turbulence

- Based on the **Navier-Stokes equation**.

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{1}{\rho} \nabla p + \nu \nabla^2 \mathbf{v}$$

↑
 Non-linear
inertial term

↑
 Dissipative term

- In turbulent flows the non-linear (inertial) term of this equation becomes important relative to the dissipative term.
- The ratio $\frac{\text{non-linear inertial term}}{\text{viscous term}}$ is the Reynolds number, Re.

$$\text{Re} = \frac{LU}{\nu}$$

At high Re dissipation can be neglected.

- Behaviour of turbulence at very high Re when **fully developed**: Richardson cascade and Kolmogorov energy spectrum).



Classical Turbulence

- Based on the **Navier-Stokes equation**.

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{1}{\rho} \nabla p + \nu \nabla^2 \mathbf{v}$$

Non-linear
inertial term

Dissipative term

- In turbulent flows the non-linear (inertial) term of this equation becomes important relative to the dissipative term.
- The ratio $\frac{\text{non-linear inertial term}}{\text{viscous term}}$ is the Reynolds number, Re.

$$\text{Re} = \frac{LU}{\nu}$$

At high Re dissipation can be neglected.

- Behaviour of turbulence at very high Re when **fully developed**: Richardson cascade and Kolmogorov energy spectrum).




UNIVERSITY OF OREGON

Richardson cascade and the Kolmogorov spectrum

- Non-linearities cause coupling of eddies of different sizes.
- This coupling causes turbulent energy to flow (at a rate ϵ per unit mass) from large eddies to small eddies in a cascade.

$$u'l/\nu \gg 1$$

ϵ




UNIVERSITY OF OREGON

Richardson cascade and the Kolmogorov spectrum

- Non-linearities cause coupling of eddies of different sizes.
- This coupling causes turbulent energy to flow (at a rate ϵ per unit mass) from large eddies to small eddies in a cascade.

$$u'l/\nu \gg 1$$

ϵ



UNIVERSITY OF OREGON

Richardson cascade and the Kolmogorov spectrum

- Non-linearities cause coupling of eddies of different sizes.
- This coupling causes turbulent energy to flow (at a rate ϵ per unit mass) from large eddies to small eddies in a cascade.

- Let u' be the characteristic velocity in an eddy of size l .


- If $u'l/\nu \gg 1$ the flow of energy in the cascade is unaffected by viscosity and conserves energy. We call this an **inertial flow of energy**.

- When $u'l/\nu \sim 1$ **viscous dissipation** takes place and the cascade is terminated.

- How is energy distributed over eddy size in the steady state?

- According to Kolmogorov, for the inertial range: $u_r^2 = \epsilon^{2/3} l^{2/3}$

- Cut-off due to viscosity is at $\eta = (\nu^3/\epsilon)^{1/4}$



Kolmogorov spectrum



UNIVERSITY OF OREGON

Homogeneous Isotropic Turbulence

The energy spectrum E is independent of energy production processes for all wavenumbers large compared to those at which the production occurs. Then E depends only on wavenumber, dissipation and viscosity

$$E = E(k, \varepsilon, \nu).$$


If the cascade is long enough, there may be an intermediate range (the inertial sub-range) in which the action of viscosity has not yet come in, that is $E = E(k, \varepsilon)$. Then dimensional analysis gives

$$E = A \varepsilon^{2/3} k^{-5/3}$$

which is the famous Kolmogorov result. Here A is a constant.



Energy spectrum on Kolmogorov ideas



UNIVERSITY OF OREGON

Search for the $5/3$ spectrum
The difference in tidal phase drives a steady current
through the tidal channel at about $Re \sim 4 \times 10^7$



UNIVERSITY OF OREGON




UNIVERSITY OF OREGON

Turbulence probe



UNIVERSITY OF OREGON




UNIVERSITY OF OREGON

What is Quantum Turbulence and Why Study It?


Helium II and $^3\text{He-B}$ are regarded as a mixture of a normal viscous fluid and a totally inviscid superfluid. In principle either or both fluids could become turbulent



Phase diagram for liquid ^4He



Fountain Pressure




$$\Delta P / \Delta T = \rho S$$




UNIVERSITY OF OREGON

Viscosity Paradox in He II



$\eta < 10^{-11}$ poise

(a)



$\eta \sim 10^{-5}$ poise

(b)



UNIVERSITY OF OREGON

These experiments suggest there are two different fluid present: a "normal fluid" which damps the oscillating disc, and a "superfluid" which can flow through the smallest channels.

$$\rho = \rho_n + \rho_s$$

The two fluids can also have their own velocity fields

$$V_n, V_s$$



Tisza Two Fluid Equations


$$\rho_s \frac{D\mathbf{v}_s}{Dt} = - \frac{\rho_s}{\rho} \nabla P + \rho_s S \nabla T$$

$$\rho_n \frac{D\mathbf{v}_n}{Dt} = - \frac{\rho_n}{\rho} \nabla P - \rho_s S \nabla T + \eta \nabla^2 \mathbf{v}_n$$




These equations result in two forms of sound, called “first” and “second” sound

$$u_1^2 \approx dp / d\rho, \quad u_2^2 \approx TS^2 \rho_s / C \rho_n$$



First sound




Second sound



UNIVERSITY OF OREGON



Side view



Top view




UNIVERSITY OF OREGON

Shortly thereafter Schwarz and Donnelly built an apparatus that could fire vortex rings at an array of quantized vortex lines such as is shown in Figure 5 [9]. The results show that the cross section for scattering of rings from lines is the order of the ring diameter, which indicates that the line and ring need to be very nearly in contact to have a strong interaction. This result is in good agreement with Feynman's speculations as quoted in the discussions by Feynman of Figure 4. Schwarz then began his careful numerical simulations of vortex line turbulence continuing over a period of years. For an interesting example of Schwarz's work see Figure 7




In turbulent counterflows the vortex lines are considered to be a tangle of quantized vortex lines (a suggestion of Feynman).



Much can be learned about quantized vortex lines by computer simulations pioneered by Schwarz and continued by Barenghi, Tsubota and others.



UNIVERSITY OF OREGON




UNIVERSITY OF OREGON

Experimental results on superfluid ^4He

- **Pressure fluctuations:** experiment of Maurer and Tabeling on swirling flow generated by two counter-rotating blades.

(a) 2.3 K
(b) 2.08 K
(c) 1.4 K





Experimental results on superfluid ^4He

- Pressure fluctuations - experiment of Maurer and Tabeling on swirling flow generated by two counter-rotating blades

- (a) 2.3 K
- (b) 2.08 K
- (c) 1.4 K

No change as we go through the superfluid phase transition!
Hence the same type of cascade, with the same Kolmogorov spectrum, above and below the transition temperature!







Vortex lines

$$\Psi = |\Psi|e^{i\phi} \Rightarrow \text{Velocity } \mathbf{v} = (\hbar/m)\nabla\phi$$

$$\oint_C \mathbf{v} \cdot d\mathbf{r} = \frac{h}{m} = \kappa \quad \text{if } C \text{ encloses a vortex line}$$



A vortex is a hole (radius $\xi \approx 10^{-8}\text{cm}$ in ^4He) around which the phase changes by 2π , and $v = \kappa/(2\pi r)$



Phase



Vortex in a homogeneous condensate (e.g. liquid helium)

Vortex lines



Vortex in an inhomogeneous condensate (e.g. trapped atomic gas)

Vortex lines




- Vortex rings, Kelvin waves, vortex knots:
studied since the times of Helmholtz, Lord Kelvin and Tait
- Vortex lines = stable topological defects in a perfect fluid =
= attractive models of elementary particles

W. Rankine *Molecular vortices* 1849


W. Thomson (Lord Kelvin) *On vortex atoms* 1867

J.J. Thomson *A treatise on the motion of vortex rings* 1883


Visualization of individual vortex lines




Vortex lattice in a rotating superfluid



(Maryland)



(MIT)





(Berkeley)




Vortex tangle

Quantum turbulence is a **tangle of vortex lines**, generated

- with propellers, grids, forks, wires, heat flows, etc in liquid helium
- with laser stirrers, shaking the trap, etc in atomic condensates





Within a turbulent tangle, we recognize processes such as **Kelvin waves, links and knots, reconnections**



Vortex tangle


Quantum turbulence is a **tangle of vortex lines**, generated

- with propellers, grids, forks, wires, heat flows, etc in liquid helium
- with laser stirrers, shaking the trap, etc in atomic condensates




Within a turbulent tangle, we recognize processes such as **Kelvin waves, links and knots, reconnections**









MPlayer

ncfb@lario: ~/conf/per2014

ncfb@lario:~\$ cd conf/per2014

per2.pdf - Adobe Reader

File Edit View Document Tools Window Help

per2.pdf x

124%

Find

Kelvin waves

t = 40 s

t = 20 s

t = 0 s

Playing crabchase.avi.
 libavformat version 53.
 Mismatching header vers
 AVI file format detecte
 [aviheader] Video strea
 AVI: No audio stream fou
 VIDEO: [MP42] 512x384 24bpp 30.000 fps 912.0 kbps (111.3 kbyte/s)
 Clip info:
 Software: Lavf52.16.0
 Load subtitles in ./
 Failed to open VDPAU backend libvdpau_nvidia.so: cannot open shared object file:
 No such file or directory
 [vdpau] Error when calling vdp_device_create_x11: 1
 =====
 Opening video decoder: [ffmpeg] FFmpeg's libavcodec codec family
 libavcodec version 53.35.0 (external)
 Mismatching header version 53.32.2
 Selected video codec: [ffmp42] vfm: ffmpeg (FFmpeg MSMPEG-4 v2)
 =====
 Audio: no sound
 Starting playback...
 Movie-Aspect is 1.33:1 - prescaling to correct movie aspect.
 VO: [xv] 512x384 => 512x384 Planar YV12
 V: 3.2 95/ 95 3% 0% 0.0% 0 0

MPlayer

z = 4R

t = 4.8s

Adobe Reader 9

ncfb@lario: ~/conf/per2014

ncfb@lario:~\$ cd conf/per2014

MPlayer

per2.pdf - Adobe Reader

File Edit View Document Tools Window Help

per2.pdf [x]

124%

Find

Kelvin waves

Dispersion relation of Kelvin waves was determined in 1880, but it took more than a century to notice that a wavy vortex ring moves slower

Kiknadze & Mamaladze, JLTP 2002
 Hänninen, Tsubota & CFB, PRE 2006
 Helm, CFB & al, PRA 2011

Carlo F. Barenghi Quantum turbulence

V0: [XV] 512X384 => 512X384 Planar YV1Z
 calc V: 1.3 40/ 40 4% 0% 0.0% 0 0
 ===== PAUSE =====

object file:
 =====
 =====

```
ncfb@lario: ~/conf/per2014
ncfb@lario:~$ cd conf/per2014
```

```
MPlayer
```

per2.pdf - Adobe Reader


File Edit View Document Tools Window Help

per2.pdf x


124% Find

Reconnections

- Bewley, Paoletti, Sreenivasan, & Lathrop 2008
Direct observation of vortex reconnections in superfluid helium
(vortex lines visualised by trapped hydrogen particles)



$\delta(t) \sim (t_0 - t)^{1/2}$ before
 $\delta(t) \sim (t - t_0)^{1/2}$ after



Carlo F. Barenghi Quantum turbulence

VO: [XV] 512X384 => 512X384 Planar YV12
 calc V: 1.3 40/ 40 4% 0% 0.0% 0 0
 ===== PAUSE =====

MPlayer

ncfb@lario: ~/conf/per2014

ncfb@lario:~\$ cd conf/per2014

per2.pdf - Adobe Reader

File Edit View Document Tools Window

per2.pdf x

124%

Reconnections

- Bewley, Paoletti, ...
- Direct ob ...
- (vortex li ...

Clip info

major_br ...

minor_ve ...

compatib ...

creation ...

Load subt ...

Failed to open VDPAU backend libvdpau_nvidia.so: cannot open shared object file: No such file or directory

[vdpau] Error when calling vdp_device_create_x11: 1

=====

Opening video decoder: [ffmpeg] FFmpeg's libavcodec codec family

libavcodec version 53.35.0 (external)

Mismatching header version 53.32.2

Selected video codec: [ffh264] vfm: ffmpeg (FFmpeg H.264)

=====

Audio: no sound

Starting playback...

Unsupported PixelFormat 61

Unsupported PixelFormat 53

Unsupported PixelFormat 81

Movie-Aspect is 1.61:1 - prescaling to correct movie aspect.

VO: [xv] 640x360 => 640x396 Planar YV12

V: 0.5 0/ 0 15% 2% 0.0% 0 0

M: 14.3 0/ 0 12% 0% 0.0% 0 0

calc

$t < t_0$

$\delta(t) \sim$

$\delta(t) \sim$

MPlayer

Adobe Reader 9

ncfb@lario: ~/conf/per2014

```
ncfb@lario:~$ cd conf/per2014
```

per2.pdf - Adobe Reader

File Edit View Document Tools Window Help

per2.pdf [x]

124%


Find

Reconnections

- In classical fluids reconnections are associated with **losses of kinetic energy** arising from **viscous** or **resistive** effects
- In quantum fluids near absolute zero such losses are **acoustic** (sound = sink of kinetic energy)

Sound waves emitted by:

- Kelvin waves
Vinen, Kozik & Svistunov, Sonin, L'vov & Nazarenko, Krstulovic, Baggaley & Hänninen, etc
- Reconnections
Zuccher, CFB & al, PoF 2012
Leadbeater, CFB & al, PRL 2001



Carlo F. Barenghi Quantum turbulence

calc Exiting... (End of file)
ncfb@lario:~/conf/per2014\$

MPlayer

ncfb@lario: ~/conf/per2

ncfb@lario:~\$ cd conf/per2

per2.pdf - Adobe Reader

File Edit View Document To

per2.pdf

Reconnect

- In classi of kine
- In quar (sound

Sound wav

- Kelvin wa Vinen, K L'vov & Baggaley
- Reconned Zuccher, Leadbeat

Audio: no sound
Starting playback...
Movie-Aspect is undefined - no prescaling applied.
VO: [xv] 716x608 => 716x608 Planar YV12
M: 0.4 12/ 12 ???% ???% ??,??% 0 0 0.10x

MPlayer

ncfb@lario: ~/conf/per2

ncfb@lario:~\$ cd conf/per2

per2.pdf - Adobe Reader

File Edit View Document To

per2.pdf

Reconnect

- In classi of kine
- In quar (sound

Sound wav

- Kelvin wa Vinen, K L'vov & Baggaley
- Reconned Zuccher, Leadbeat

Audio: no sound
Starting playback...
Movie-Aspect is undefined - no prescaling applied.
VO: [xv] 716x608 => 716x608 Planar YV12
V: 0.2 7/ 7 ???% ???% ??,??% 0 0 0.10x

Adobe Reader 9

ncfb@lario: ~/conf/per2014

```
ncfb@lario:~$ cd conf/per2014
```

per2.pdf - Adobe Reader

File Edit View Document Tools Window Help

per2.pdf [x]

124% Find

Quantum turbulence

There seems to be two forms of quantum turbulence:
(Volovik 2004; Walmsley & Golov 2008)

- **Ultra-quantum:**
featureless, most energy at mesoscales,
Vinen 1957
- **Semi-classical:**
contains coherent structures,
most energy at large scales as in ordinary turbulence
classical Kolmogorov spectrum (Maurer & Tabeling 1998)
classical decay rate (Smith, Donnelly, Goldenfeld & Vinen 1993)

Carlo F. Barenghi Quantum turbulence

Exiting... (Quit)
ncfb@lario:~/conf/per2014\$

Adobe Reader 9

ncfb@lario: ~/conf/per2014

```
ncfb@lario:~$ cd conf/per2014
```

per2.pdf - Adobe Reader

File Edit View Document Tools Window Help

per2.pdf [x]

124% Find

Quantum turbulence

There seems to be two forms of quantum turbulence:
(Volovik 2004; Walmsley & Golov 2008)

- **Ultra-quantum:**
featureless, most energy at mesoscales,
Vinen 1957
- **Semi-classical:**
contains coherent structures,
most energy at large scales as in ordinary turbulence
classical Kolmogorov spectrum (Maurer & Tabeling 1998)
classical decay rate (Smith, Donnelly, Goldenfeld & Vinen 1993)

Carlo F. Barenghi Quantum turbulence

Exiting... (Quit)
ncfb@lario:~/conf/per2014\$

Adobe Reader 9

ncfb@lario: ~/conf/per2014

```
ncfb@lario:~$ cd conf/per2014
```

per2.pdf - Adobe Reader

File Edit View Document Tools Window Help

per2.pdf [x] 124% Find

Quantum turbulence

Total vorticity (left) decomposed into
 coherent (right, top) and
 incoherent part (right, bottom)

Baggaley, Laurie & CFB, PRL 2012

Carlo F. Barenghi Quantum turbulence

calc Exiting... (Quit)
 ncfb@lario:~/conf/per2014\$

ncfb@lario: ~/conf/per2014
ncfb@lario:~\$ cd conf/per2014


per2.pdf - Adobe Reader

File Edit View Document Tools Window Help

per2.pdf

124% Find

Quantum turbulence



Energy spectrum, Roche & al, Grenoble



Double cascade

Possibility of **bottleneck** accumulation of energy between the semi-classical (3D) $k^{-5/3}$ **Kolmogorov cascade** for $k \ll 2\pi/\ell$ and the (1D) **Kelvin wave cascade** for $k \gg 2\pi/\ell$, where ℓ is the average distance between vortex lines

CFB, L'vov and Roche, PNAS 2014

Exiting... (Quit)
ncfb@lario:~/conf/per2014\$

ncfb@lario: ~/conf/per2014

ncfb@lario:~\$ cd conf/per2014

per2.pdf - Adobe Reader


File Edit View Document Tools Window Help

per2.pdf

124%

Find

Quantum turbulence



Energy spectrum, Roche & al, Grenoble



Double cascade

Possibility of **bottleneck** accumulation of energy between the semi-classical (3D) $k^{-5/3}$ **Kolmogorov cascade** for $k \ll 2\pi/\ell$ and the (1D) **Kelvin wave cascade** for $k \gg 2\pi/\ell$, where ℓ is the average distance between vortex lines

CFB, L'vov and Roche, PNAS 2014

Exiting... (Quit)
ncfb@lario:~/conf/per2014\$

Adobe Reader 9

ncfb@lario: ~/conf/per2014

```
ncfb@lario:~$ cd conf/per2014
```

per2.pdf - Adobe Reader

File Edit View Document Tools Window Help

per2.pdf [x]

124% Find

Conclusion

Why quantum turbulence ?

- Interesting physics per se (vortex dynamics, Kibble–Zurek, etc)
- Relation with classical turbulence: Does classical behaviour emerge from a sufficient number of quanta ?
- Topology of turbulent flows
- Challenge: flow visualization, small probes

See recent **PNAS** special issue on quantum turbulence edited by CFB, Skrbek & Sreenivasan (March 2014)

Thank you !

EPSRC

Carlo F. Barenghi Quantum turbulence

Exiting... (Quit)
ncfb@lario:~/conf/per2014\$