

Title: Lattice models for anomalous field theories

Date: May 29, 2014 04:00 PM

URL: <http://pirsa.org/14050110>


Abstract:



# Lattice models for anomalous field theories

John McGreevy, UCSD

based on work with  
S. M. Kravec and Brian Swingle



# This talk is about 'emergeability'

[Senthil]

- Crucial step in EFT: identify
- degrees of freedom
  - realization of symmetry.

# This talk is about 'emergeability'

[Senthil]

- Crucial step in EFT: identify
- degrees of freedom
  - realization of symmetry.

Extra *high-energy* requirement:

Given a microphysical  $\mathcal{H}_{UV}$ , ( e.g.  $\mathcal{H}_{UV} = \otimes_j \mathcal{H}_j$  )  
do these dofs act on (a subspace of)  $\mathcal{H}_{UV}$ ?





Créteur, Alexander Monin,  
& Francesco Riva

effective SSB

/ gravity

$a, b, c \rightarrow \ell, \ell$   
 $u, d, k \rightarrow s, \ell, \ell$   
 $m, v, \rightarrow st.$

## This talk is about 'emergeability'

[Senthil]

Crucial step in EFT: identify

- degrees of freedom
- realization of symmetry.

Extra *high-energy* requirement:

Given a microphysical  $\mathcal{H}_{UV}$ , (e.g.  $\mathcal{H}_{UV} = \otimes_j \mathcal{H}_j$ )  
do these dofs act on (a subspace of)  $\mathcal{H}_{UV}$ ?

Checks: all RG invariants must be the same.

Familiar hep-th avatar: 't Hooft anomaly matching.

A different kind of example: no gauge invariant fermion operators in an EFT for a bosonic system.

# This talk is about 'emergeability'

[Senthil]

- Crucial step in EFT: identify
- degrees of freedom
  - realization of symmetry.

Extra *high-energy* requirement:

Given a microphysical  $\mathcal{H}_{UV}$ , ( e.g.  $\mathcal{H}_{UV} = \otimes_j \mathcal{H}_j$  )  
do these dofs act on (a subspace of)  $\mathcal{H}_{UV}$ ?

Checks: all RG invariants must be the same.

Familiar hep-th avatar: 't Hooft anomaly matching.

A different kind of example: no gauge invariant fermion operators in an EFT for a bosonic system.



acretac, Alexander Monin,  
& Francesco Riva

effective SSB

/ gravity

$a, b, c \rightarrow \text{LL}$   
 $u, d, s \rightarrow \text{LL}$   
 $m, v, \dots \rightarrow \text{st.}$

## This talk is about 'emergeability'

[Senthil]

Crucial step in EFT: identify

- degrees of freedom
- realization of symmetry.

Extra *high-energy* requirement:

Given a microphysical  $\mathcal{H}_{UV}$ , (e.g.  $\mathcal{H}_{UV} = \otimes_j \mathcal{H}_j$ )  
do these dofs act on (a subspace of)  $\mathcal{H}_{UV}$ ?

Checks: all RG invariants must be the same.

Familiar hep-th avatar: 't Hooft anomaly matching.

A different kind of example: no gauge invariant fermion operators in an EFT for a bosonic system.



crétac, Alexander Manin,  
& Francesco Riva

active SSB

/ gravity

$a, b, c \rightarrow \mathbb{C} \mathbb{C}$   
 $u, v, k \rightarrow \mathbb{S} \mathbb{C} \mathbb{C}$   
 $m, v, \rightarrow \mathbb{S} \mathbb{t}$

## This talk is about 'emergeability'

[Senthil]

Crucial step in EFT: identify

- degrees of freedom
- realization of symmetry.

Extra *high-energy* requirement:

Given a microphysical  $\mathcal{H}_{UV}$ , (e.g.  $\mathcal{H}_{UV} = \otimes_j \mathcal{H}_j$ )  
do these dofs act on (a subspace of)  $\mathcal{H}_{UV}$ ?

Checks: all RG invariants must be the same.

Familiar hep-th avatar: 't Hooft anomaly matching.

A different kind of example: no gauge invariant fermion operators in an EFT for a bosonic system.

This might be considered a

High-Energy Problem for Low-Energy Physicists.

Many examples raised so far this week, e.g.:

- can a  $p + ip$  superfluid be coupled to gravity without a spin structure?
- does a CFT always have a stress tensor?



Créteil, Alexander Monin,  
& Francesco Riva

active SSB

/ gravity

$a, b, c \rightarrow c, c$   
 $u, v, t \rightarrow s, c, c$   
 $m, v, \rightarrow st.$

## High-energy physics point of view

**Goal:** Identify obstructions  
to symmetry-preserving regulators of QFT,  
by thinking about certain states of matter in one higher dimension  
which have an energy gap  
(i.e.  $E_1 - E_{gs} > 0$  in thermodynamic limit).

These 'SPT (symmetry-protected topological) states'

[Wen et al; Reviews: Turner-Vishwanath, 1301.0330; Senthil, review of 'physics-based approach' 1405.4015]

are machines for producing such obstructions.

(Their study is also a useful step toward understanding more difficult states.)



# Plan

1. Introduction
2. Ideas about regularizing the Standard Model  
[Wen, J. Wang-Wen, You-BenTov-Xu]
3. Constraints on 3+1d QFT  
from symmetry-protected topological (SPT) states  
In particular, we'll identify constraints on manifest electric-magnetic duality symmetry.  
[Shauna Kravec, JM, 1306.3992, PRL  
work in progress with Brian Swingle]
4. A machine for *explicitly* realizing SPT states  
[Shauna Kravec, JM, Brian Swingle, in progress]

Alexander Morion,

Massimo Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$


$m, v, \rightarrow \text{st.}$

## Realizations of symmetries in QFT and cond-mat

**Basic Q:** What are possible gapped phases of matter?

**Def:** Two gapped states are equivalent if they are adiabatically connected

(varying the parameters in the  $H$  whose ground state they are to get from one to the other, without closing the energy gap).



One important distinguishing feature: how are the symmetries realized?

**Landau distinction:** characterize by *broken* symmetries

e.g. ferromagnet vs paramagnet, insulator vs SC.



Alexander Morion,

Francesco Riva

$$a, b, c \rightarrow \text{c.c.}$$

$$u, v, k \rightarrow \text{s.c.c.}$$


$$m, v, \rightarrow \text{st.}$$

## Realizations of symmetries in QFT and cond-mat

**Basic Q:** What are possible gapped phases of matter?

**Def:** Two gapped states are equivalent if they are adiabatically connected

(varying the parameters in the  $H$  whose ground state they are to get from one to the other, without closing the energy gap).



One important distinguishing feature: how are the symmetries realized?

**Landau distinction:** characterize by *broken* symmetries

e.g. ferromagnet vs paramagnet, insulator vs SC. ✓

**Mod out by Landau:** "What are possible (gapped) phases that don't break symmetries?" How do we distinguish them?

One (far) answer: symmetries can be *fractionalized*.

[Wen]: topological order.

This means emergent deconfined gauge theory, long-range entanglement.



Alexander Morita,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{s.C.L.}$

$m, v, \rightarrow \text{st.}$

## Mod out by Wen, too

"What are possible (gapped) phases that don't break symmetries and don't have topological order?"




Alexander Morion,

Francesco Riva

$a, b, c \rightarrow \text{c.l.}$   
 $u, v, k \rightarrow \text{s.l.}$   
 $m, v, \rightarrow \text{st.}$

## SRE states are characterized by their edge states

**Rough idea:** just like varying the Hamiltonian in time to another phase requires closing the gap  $\mathbf{H} = \mathbf{H}_1 + g(t)\mathbf{H}_2$ , so does varying the Hamiltonian in space  
 $\mathbf{H} = \mathbf{H}_1 + g(x)\mathbf{H}_2$ .



- Important role of SRE assumption: Here we are assuming that the bulk state has short-ranged correlations, so that changes we might make at the surface cannot have effects deep in the bulk.



Alexander Morita,

Massimo Riva

$$a, b, c \rightarrow \text{c.c.}$$

$$u, v, k \rightarrow \text{s.c.c.}$$

$$m, v \rightarrow \text{s.t.}$$

## SPT states

**Def:** An *SPT state* (symmetry-protected topological state), protected by a symmetry group  $G$  is:  
a SRE state, which is not adiabatically connected to a product state by local hamiltonians preserving  $G$ .

e.g.: free fermion topological insulators in 3+1d, protected by  $U(1)$  and  $\mathcal{T}$ , have an odd number of Dirac cones on the surface.

One reason to care: if you gauge  $H \subset G$ , you get a state with topological order.

## SPT states

**Def:** An *SPT state* (symmetry-protected topological state), protected by a symmetry group  $G$  is:

a SRE state, which is not adiabatically connected to a product state by local hamiltonians preserving  $G$ .

**e.g.:** free fermion topological insulators in 3+1d, protected by  $U(1)$  and  $\mathcal{T}$ , have an odd number of Dirac cones on the surface.

One reason to care: if you gauge  $H \subset G$ , you get a state with topological order.



Alexander Morion,

Massimo Riva

$$a, b, c \rightarrow \text{C.L.}$$

$$u, v, k \rightarrow \text{s.c.}$$

$$m, v, \rightarrow \text{st.}$$

## SPT states

**Def:** An *SPT state* (symmetry-protected topological state), protected by a symmetry group  $G$  is:  
a SRE state, which is not adiabatically connected to a product state by local hamiltonians preserving  $G$ .

e.g.: free fermion topological insulators in 3+1d, protected by  $U(1)$  and  $\mathcal{T}$ ,  
have an odd number of Dirac cones on the surface.

One reason to care: if you gauge  $H \subset G$ , you get a state with topological order.



# SPT states

**Def:** An *SPT state* (symmetry-protected topological state), protected by a symmetry group  $G$  is:


a SRE state, which is not adiabatically connected to a product state by local hamiltonians preserving  $G$ .

e.g.: free fermion topological insulators in 3+1d, protected by  $U(1)$  and  $\mathcal{T}$ , have an odd number of Dirac cones on the surface.

One reason to care: if you gauge  $H \subset G$ , you get a state with topological order.

- ▶ Free fermion TIs classified [Kitaev: homotopy theory; Schneider et al: edge]

Interactions can affect the connectivity of the phase diagram in both directions:



- ▶ There are states which are adiabatically connected only via interacting Hamiltonians [Fidkowski-Kitaev, 0904.2197, Qi, Yao-Ryu, Wang-Senthil, You-BenTov-Xu].
- ▶ There are states whose existence requires interactions:  
e.g. Bosonic SPT states – w/o interactions, superfluid.

Alexander Morion,

Massimo Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{s.t.}$

## SPT states


**Def:** An *SPT state* (symmetry-protected topological state), protected by a symmetry group  $G$  is:  
a SRE state, which is not adiabatically connected to a product state by local hamiltonians preserving  $G$ .

e.g.: free fermion topological insulators in 3+1d, protected by  $U(1)$  and  $\mathcal{T}$ , have an odd number of Dirac cones on the surface.

One reason to care: if you gauge  $H \subset G$ , you get a state with topological order.

- Free fermion TIs classified [Kitaev: homotopy theory; Schneider et al: edge]

Interactions can affect the connectivity of the phase diagram in both directions:



- There are states which are adiabatically connected only via interacting Hamiltonians [Fidkowski-Kitaev, 0904.2197, Qi, Yao-Ryu, Wang-Senthil, You-BenTov-Xu].
- There are states whose existence requires interactions:  
e.g. Bosonic SPT states – w/o interactions, superfluid.



## SPT states

**Def:** An *SPT state* (symmetry-protected topological state), protected by a symmetry group  $G$  is:


a SRE state, which is not adiabatically connected to a product state by local hamiltonians preserving  $G$ .

e.g.: free fermion topological insulators in  $3+1d$ , protected by  $U(1)$  and  $\mathcal{T}$ , have an odd number of Dirac cones on the surface.

One reason to care: if you gauge  $H \subset G$ , you get a state with topological order.

- ▶ Free fermion TIs classified [Kitaev: homotopy theory; Schneider et al: edge]

Interactions can affect the connectivity of the phase diagram in both directions:



- ▶ There are states which are adiabatically connected only via interacting Hamiltonians [Fidkowski-Kitaev, 0904.2197, Qi, Yao-Ryu, Wang-Senthil, You-BenTov-Xu].
- ▶ There are states whose existence requires interactions:  
e.g. Bosonic SPT states – w/o interactions, superfluid.

Alexander Morion,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

## SPT states


**Def:** An *SPT state* (symmetry-protected topological state), protected by a symmetry group  $G$  is:  
a SRE state, which is not adiabatically connected to a product state by local hamiltonians preserving  $G$ .

e.g.: free fermion topological insulators in 3+1d, protected by  $U(1)$  and  $\mathcal{T}$ , have an odd number of Dirac cones on the surface.

One reason to care: if you gauge  $H \subset G$ , you get a state with topological order.

- Free fermion TIs classified [Kitaev: homotopy theory; Schneider et al: edge]

Interactions can affect the connectivity of the phase diagram in both directions:



- There are states which are adiabatically connected only via interacting Hamiltonians [Fidkowski-Kitaev, 0904.2197, Qi, Yao-Ryu, Wang-Senthil, You-BenTov-Xu].
- There are states whose existence requires interactions:  
e.g. Bosonic SPT states – w/o interactions, superfluid.



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, v, \rightarrow \text{st.}$

## Group structure of SPT states



Simplifying feature:

SPT states (for given G) form a group:





Alexander Morion,

cesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

## Group structure of SPT states

Simplifying feature:

SPT states (for given  $G$ ) form a group:




$A$



$A + B$

$-A$  : is the mirror image.



$A - A$



Alexander Morita,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, n, l \rightarrow \text{st.}$


## Group structure of SPT states



Simplifying feature:

SPT states (for given  $G$ ) form a group:

$-A$  : is the mirror image.



- With bulk topological order, bulk quasiparticles still nontrivial. Not a group.
  - There can be many realizations of the edge of  $A$ , same 'SPT-ness'.
  - The edge of  $A$  can be symmetric and gapped but topologically ordered.
- Inverse of  $A$  cancels the SPT-ness of  $A$ 's edge.



Alexander Morion,

cesco Riva

$a, b, c \rightarrow \mathbb{C}, \mathbb{C}$   
 $u, v, k \rightarrow s, \mathbb{C}, \mathbb{C}$   
 $m, v, \rightarrow st.$


## Group structure of SPT states



Simplifying feature:

SPT states (for given G) form a group:

$-A$  : is the mirror image.



- With bulk topological order, bulk quasiparticles still nontrivial. Not a group.
  - There can be many realizations of the edge of A, same 'SPT-ness'.
  - The edge of A can be symmetric and gapped but topologically ordered.
- Inverse of A cancels the SPT-ness of A's edge.



Alexander Morion,

cesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$


## Group structure of SPT states



Simplifying feature:

SPT states (for given G) form a group:

$-A$  : is the mirror image.



- With bulk topological order, bulk quasiparticles still nontrivial. Not a group.
  - There can be many realizations of the edge of A, same 'SPT-ness'.
  - The edge of A can be symmetric and gapped but topologically ordered.
- Inverse of A cancels the SPT-ness of A's edge.



Alexander Morin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.L.}$   
 $m, v, \rightarrow \text{st.}$


## Group structure of SPT states

Simplifying feature:

SPT states (for given  $G$ ) form a group:



$-A$  : is the mirror image.



- With bulk topological order, bulk quasiparticles still nontrivial. Not a group.
  - There can be many realizations of the edge of  $A$ , same 'SPT-ness'.
  - The edge of  $A$  can be symmetric and gapped but topologically ordered.
- Inverse of  $A$  cancels the SPT-ness of  $A$ 's edge.



Alexander Morita,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{S.L.}$

$m, v, \rightarrow \text{st.}$


## Group structure of SPT states



Simplifying feature:


SPT states (for given G) form a group:

$-A$  : is the mirror image.



- With bulk topological order, bulk quasiparticles still nontrivial. Not a group.
  - There can be many realizations of the edge of A, same 'SPT-ness'.
  - The edge of A can be symmetric and gapped but topologically ordered.
- Inverse of A cancels the SPT-ness of A's edge.


## Group structure of SPT states



Simplifying feature:

SPT states (for given  $G$ ) form a group:

$-A$  : is the mirror image.



- With bulk topological order, bulk quasiparticles still nontrivial. Not a group.
- There can be many realizations of the edge of  $A$ , same 'SPT-ness'.
- The edge of  $A$  can be symmetric and gapped but topologically ordered.

Inverse of  $A$  cancels the SPT-ness of  $A$ 's edge.



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{S.L.}$

$m, v \rightarrow \text{st.}$

## Surface-only models

### Counterfactual:

Suppose the edge theory of an SPT state were realized *otherwise*  
– intrinsically in  $D$  dimensions, with a local hamiltonian.

Then we could paint that the conjugate local theory on the surface without changing anything about the bulk state.





Alexander Morita,

Vincenzo Riva


$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, v_1 \rightarrow \text{st.}$

## Surface-only models

### Counterfactual:

Suppose the edge theory of an SPT state were realized *otherwise*  
– intrinsically in  $D$  dimensions, with a local hamiltonian.

Then we could paint that the conjugate local theory on the surface without changing anything about the bulk state.  
And then small deformations of the surface Hamiltonian, localized on the surface, consistent with symmetries, can pair up the edge states.



But this contradicts the claim that we could characterize the  $D + 1$ -dimensional SPT state by its edge theory.

**Conclusion:** Edge theories of  $\text{SPT}_G$  states cannot be regularized intrinsically in  $D$  dims, *exactly* preserving on-site  $G$  –  
“surface-only models” or “not edgeable”.

[Wang-Senthil, 1302.6234 – general idea, concrete surprising examples of 2+1 surface-only states  
Wen, 1303.1803 – attempt to characterize the underlying mathematical structure, classify *all* such obstructions  
Metlitski-Kane-Fisher, 1302.6535; Burnell-Chen-Fidkowski-Vishwanath, 1302.7072]



Alexander Monin,

Francesco Riva

$a, b, c_i \rightarrow \text{L.L.}$

$u, v, k \rightarrow \text{s.L.L.}$

$m, v_i \rightarrow \text{st.}$

## Ideas about regularizing the Standard Model



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, v, \rightarrow \text{st.}$

## Recasting the NN result as a statement about SPT states

Consider free massive relativistic fermions in  
4+1 dimensions (with conserved  $U(1)$ ):

$$S = \int d^{4+1}x \bar{\Psi} (\not{\partial} + m) \Psi$$

$\pm m$  label distinct Lorentz-invariant  
( $P$ -broken) phases.



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, v \rightarrow \text{st.}$

## Recasting the NN result as a statement about SPT states

Consider free massive relativistic fermions in  
 4+1 dimensions (with conserved  $U(1)$ ):

$$S = \int d^{4+1}x \bar{\Psi} (\not{\partial} + m) \Psi$$

$\pm m$  label distinct Lorentz-invariant  
 ( $P$ -broken) phases.

Domain wall between them  
 hosts (exponentially-localized)

3+1 chiral fermions: [Jackiw-Rebbi,


Callan-Harvey, Kaplan...]

One proof of this:

Couple to external gauge field

$$\Delta S = \int d^5x A^\mu \bar{\Psi} \gamma_\mu \Psi.$$

$$\log \int [D\Psi] e^{iS_{4+1}[\Psi, A]} \propto \frac{m}{|m|} \int A \wedge F \wedge F$$



Galling fact: if we want the extra dimension to be finite, there's another

domain wall with the antichiral fermions.

And if we put it too far away, the KK gauge bosons are too light...



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, v, \rightarrow \text{st.}$

## Recasting the NN result as a statement about SPT states

Consider free massive relativistic fermions in  
 4+1 dimensions (with conserved  $U(1)$ ):

$$S = \int d^{4+1}x \bar{\Psi} (\not{\partial} + m) \Psi$$

$\pm m$  label distinct Lorentz-invariant  
 ( $P$ -broken) phases.

Domain wall between them  
 hosts (exponentially-localized)

3+1 chiral fermions: [Jackiw-Rebbi,


Callan-Harvey, Kaplan...]

One proof of this:

Couple to external gauge field

$$\Delta S = \int d^5x A^\mu \bar{\Psi} \gamma_\mu \Psi.$$

$$\log \int [D\Psi] e^{iS_{4+1}[\Psi, A]} \propto \frac{m}{|m|} \int A \wedge F \wedge F$$



Galling fact: if we want the extra dimension to be finite, there's another  
 domain wall with the antichiral fermions.

If we put it too far away, the KK gauge bosons are too light...



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{L.L.}$

$u, d, k \rightarrow \text{s.L.L.}$

$\mu, \nu \rightarrow \text{st.}$

## Loophole in NN theorem

But the SM gauge group is *not* anomalous, shouldn't need extra dimensions.

Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{L.L.}$

$l, j, k \rightarrow \text{s.L.L.}$

$\mu, \nu \rightarrow \text{s.t.}$

## Loophole in NN theorem

But the SM gauge group is *not* anomalous, shouldn't need extra dimensions.



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow L-L$   
 $u, d, k \rightarrow S-L-L$   
 $u, V, \rightarrow S-L$

## Loophole in NN theorem

But the SM gauge group is *not* anomalous, shouldn't need extra dimensions.

**Loophole:** Interactions between fermions!


**Old idea:** add four-fermion interactions (or couplings to other fields) which gap the mirror fermions, but not the SM, and preserve the SM gauge group  $G$ .

These interactions should explicitly break all anomalous symmetries.

This requires a right-handed neutrino. [Preskill-Eichten 1986]:  
SU(5) and SO(10) lattice GUTs. [Many other papers ... recent work: Wen, J. Wang]

[Preskill-Eichten 1986]: Evidence for mirror-fermion mass generation without symmetry-breaking ... coupling expansion.

[Geidt-Chen-Poppitz]: null ... for ... of a related proposal in 1+1d.





Alexander Morion,

cesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, d, k \rightarrow \text{s.c.c.}$   
 $m, v, \rightarrow \text{st.}$

## New evidence for a special role of $n_F = 16 \cdot n$

Collapse of free-fermion classification:


Dimensional recursion strategy [Wang-Senthil, Qi, Ryu-Yao, Wen, You-BenTov-Xu]:

1. Consider neighboring phase where  $G$  is spontaneously broken  $\langle \phi \rangle \neq 0$ .
2. Proliferate defects of  $\phi$  to reach paramagnetic phase.
3. Must  $\phi$ -defects carry quantum numbers which make the paramagnet nontrivial?

Initial step: [Fidkowski-Kitaev]

edge of  $8 \times$  majorana chain is symmetrically gappable.

same reformation as shows equivalence of GS and RNS superstrings SO(8) triality.



[You-BenTov-Xu]:  $4+1d$ , with many  $G$ , the collapse again happens at  $n_F = 8 \approx 0$  ( $\rightarrow 16$  Weyl fermions per domain wall.)



Alexander Morion,

cesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, d, k \rightarrow \text{s.c.c.}$   
 $m, v \rightarrow \text{st.}$

## New evidence for a special role of $n_F = 16 \cdot n$

Collapse of free-fermion classification:


Dimensional recursion strategy [Wang-Senthil, Qi, Ryu-Yao, Wen, You-BenTov-Xu]:

1. Consider neighboring phase where  $G$  is spontaneously broken  $\langle \phi \rangle \neq 0$ .
2. Proliferate defects of  $\phi$  to reach paramagnetic phase.
3. Must  $\phi$ -defects carry quantum numbers which make the paramagnet nontrivial?

Initial step: [Fidkowski-Kitaev]

edge of  $8 \times$  majorana chain is symmetrically gappable.

same refermionization as shows equivalence of GS and RNS superstrings,  $SO(8)$  triality.



[You-BenTov-Xu]: In  $4+1d$ , with many  $G$ , the collapse again happens at  $k = 8 \simeq 0$  ( $\rightarrow 16$  Weyl fermions per domain wall.)



Alexander Morin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, d, k \rightarrow \text{s.c.c.}$   
 $m, v \rightarrow \text{st.}$

## New evidence for a special role of $n_F = 16 \cdot n$

Collapse of free-fermion classification:


Dimensional recursion strategy [Wang-Senthil, Qi, Ryu-Yao, Wen, You-BenTov-Xu]:

1. Consider neighboring phase where  $G$  is spontaneously broken  $\langle \phi \rangle \neq 0$ .
2. Proliferate defects of  $\phi$  to reach paramagnetic phase.
3. Must  $\phi$ -defects carry quantum numbers which make the paramagnet nontrivial?

Initial step: [Fidkowski-Kitaev]

edge of  $8 \times$  majorana chain is symmetrically gappable.

same refermionization as shows equivalence of GS and RNS superstrings,  $SO(8)$  triality.



[You-BenTov-Xu]: In  $4+1d$ , with many  $G$ , the collapse again happens at  $k = 8 \simeq 0$  ( $\rightarrow 16$  Weyl fermions per domain wall.)  $G \supset \mathbb{Z}_2^T$  awkward:

Conclusion: This novel strategy for identifying obstructions to gapping the mirror fermions shows none when  $n_F = 16n$ .



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{s.c.L.}$

$\mu, \nu \rightarrow \text{st.}$

## Surface-only electrodynamics, by example



Alexander Monin,

Francesco Riva


$a, b, c \rightarrow L-L$   
 $l, j, k \rightarrow s, L-L$   
 $n, v, \rightarrow s, t$

## Loophole in NN theorem

But the SM gauge group is *not* anomalous, shouldn't need extra dimensions.

**Loophole:** Interactions between fermions!

**Old idea:** add four-fermion interactions (or couplings to other fields) which gap the mirror fermions, but not the SM, and preserve the SM gauge group  $G$ .



These interactions should explicitly break all anomalous symmetries.

This requires a right-handed neutrino. [Preskill-Eichten 1986]:  
SU(5) and SO(10) lattice GUTs. [Many other papers ... recent work: Wen, J. Wang]

[Preskill-Eichten 1986]: Evidence for mirror-fermion mass generation without symmetry-breaking via eucl. strong coupling expansion.

[Geidt-Chen-Poppitz]: numerical evidence for troubles of a related proposal in 1+1d.



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{S.L.}$

$\mu, \nu \rightarrow \text{st.}$

## Surface-only electrodynamics, by example



Alexander Monin,

Massimo Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{S.L.}$

$m, v, \rightarrow \text{st.}$

## Strategy

Study a simple (unitary) gapped or topological field theory in 4+1 dimensions without topological order, with symmetry  $G$ .

Consider the model on the disk with  $G$ -inv't boundary conditions.

The resulting edge theory is

a "surface-only theory with respect to  $G$ "

– it cannot be regulated by a local 3 + 1-dim'l model while preserving  $G$ .



Alexander Monin,

Massimo Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{s.c.}$

$m, v, \rightarrow \text{st.}$

## Strategy

Study a simple (unitary) gapped or topological field theory in 4+1 dimensions without topological order, with symmetry  $G$ .

Consider the model on the disk with  $G$ -inv't boundary conditions.

The resulting edge theory is

a "surface-only theory with respect to  $G$ "

– it cannot be regulated by a local 3 + 1-dim'l model while preserving  $G$ .



Alexander Morion,

cesco Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{S.L.}$

$m, v, \rightarrow \text{st.}$

## What does it mean to be a surface-only state?

These theories are perfectly consistent and unitary – they *can* be realized as the edge theory of some gapped bulk. They just can't be regularized in a local way consistent with the symmetries – without the bulk.

1. It (probably) means these QFTs will not be found as low-energy EFTs of solids or in cold atom lattice simulations.



Alexander Morion,

cesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, v_i \rightarrow \text{st.}$

## Some known examples of surface-only states

The first three examples are in  $D = 2 + 1$ , realized on the surface of  $D = 3 + 1$  boson SPTs protected by time-reversal:

- ▶ Non-linear sigma model on  $S^2$  with Hopf term at  $\theta = \pi$

$$Z = \sum_{\text{instanton number, } n} (-1)^n Z_n$$

[model of high- $T_c$ : Dzyaloshinskii-Polyakov-Wiegmann 88, surface only: Vishwanath-Senthil 12]

- ▶ "Algebraic vortex liquid": an insulating state of bosons (or a paramagnet) with massless fermionic vortices

[proposed by Fisher et al 06, surface only: Wang-Senthil 13]

- ▶ "All-fermion toric code": a version of  $\mathbb{Z}_2$  gauge theory where  $e, m, \epsilon \equiv em$  are all fermions.

[surface only: Burnell-Chen-Fidkowski-Vishwanath, Wang-Senthil]



Alexander Morion,

cesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, v, \rightarrow \text{st.}$

## Some known examples of surface-only states

The first three examples are in  $D = 2 + 1$ , realized on the surface of  $D = 3 + 1$  boson SPTs protected by time-reversal:

- ▶ Non-linear sigma model on  $S^2$  with Hopf term at  $\theta = \pi$

$$Z = \sum_{\text{instanton number, } n} (-1)^n Z_n$$

[model of high- $T_c$ : Dzyaloshinskii-Polyakov-Wiegmann 88, surface only: Vishwanath-Senthil 12]

- ▶ “Algebraic vortex liquid”: an insulating state of bosons (or a paramagnet) with massless fermionic vortices

[proposed by Fisher et al 06, surface only: Wang-Senthil 13]

- ▶ “All-fermion toric code”: a version of  $\mathbb{Z}_2$  gauge theory where  $e, m, \epsilon \equiv em$  are all fermions.

[surface only: Burnell-Chen-Fidkowski-Vishwanath, Wang-Senthil]

- ▶ “All-fermion electrodynamics”: a version of Maxwell theory where  $e, m, \epsilon \equiv em$  are all fermions.

In  $D = 3 + 1$ , with  $G = \mathbb{1}$  [surface only: Wang-Potter-Senthil 13]



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, n \rightarrow \text{st.}$

## A simple topological field theory in 4+1 dimensions

Consider 2-forms  $B_{MN}$  in 4 + 1 dimensions, with action

$$S[B] = \frac{K_{IJ}}{2\pi} \int_{\mathbb{R} \times \Sigma} B^I \wedge dB^J$$

In  $4\ell + 1$  dims,  $K$  is a skew-symmetric integer  $2N_B \times 2N_B$  matrix.


Note:  $B \wedge dB = \frac{1}{2} d(B \wedge B)$ .

Independent of choice of metric on  $\mathbb{R} \times \Sigma_{2p}$ .

Related models studied in: [Horowitz 1989, Blau et al 1989, Witten 1998,

Maldacena-Moore-Seiberg 2001, Belov-Moore 2001, Hartnoll 2006]

[Horowitz-Srednicki]: coupling to string sources  $\Delta S = \int_{\Gamma'} B^I$   
computes linking # of conjugate species of worldsheets  $\Gamma'$ .





Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$\mu, \nu \rightarrow \text{st.}$

## A simple topological field theory in 4+1 dimensions

Consider 2-forms  $B_{MN}$  in 4 + 1 dimensions, with action

$$S[B] = \frac{K_{IJ}}{2\pi} \int_{\mathbb{R} \times \Sigma} B^I \wedge dB^J$$

In  $4\ell + 1$  dims,  $K$  is a skew-symmetric integer  $2N_B \times 2N_B$  matrix.


Note:  $B \wedge dB = \frac{1}{2} d(B \wedge B)$ .

Independent of choice of metric on  $\mathbb{R} \times \Sigma_{2p}$ .

Related models studied in: [Horowitz 1989, Blau et al. 1989, Witten 1998,

Maldacena-Moore-Seiberg 2001, Belov-Moore 2005, Mottola 2006]

[Horowitz-Srednicki]: coupling to string sources  $\Delta S = \int_{\Gamma'} B^I$   
computes linking # of conjugate species of worldsheets  $\Gamma'$ .





Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

## A simple topological field theory in 4+1 dimensions

Consider 2-forms  $B_{MN}$  in 4 + 1 dimensions, with action

$$S[B] = \frac{K_{IJ}}{2\pi} \int_{\mathbb{R} \times \Sigma} B^I \wedge dB^J$$

In  $4\ell + 1$  dims,  $K$  is a skew-symmetric integer  $2N_B \times 2N_B$  matrix.


Note:  $B \wedge dB = \frac{1}{2} d(B \wedge B)$ .

Independent of choice of metric on  $\mathbb{R} \times \Sigma_{2p}$ .

Related models studied in: [Horowitz 1989, Blau et al 1989, Witten 1998,

Maldacena-Moore-Seiberg 2001, Belov-Moore 2005, Hartnoll 2006]

[Horowitz-Srednicki]: coupling to string sources  $\Delta S = \int_{\Gamma'} B^I$   
computes linking # of conjugate species of worldsheets  $\Gamma'$ .





Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, n \rightarrow \text{st.}$

## A simple topological field theory in 4+1 dimensions

Consider 2-forms  $B_{MN}$  in 4 + 1 dimensions, with action

$$S[B] = \frac{K_{IJ}}{2\pi} \int_{\mathbb{R} \times \Sigma} B^I \wedge dB^J$$

In  $4\ell + 1$  dims,  $K$  is a skew-symmetric integer  $2N_B \times 2N_B$  matrix.


Note:  $B \wedge dB = \frac{1}{2} d(B \wedge B)$ .

Independent of choice of metric on  $\mathbb{R} \times \Sigma_{2p}$ .

Related models studied in: [Horowitz 1989, Blau et al 1989, Witten 1998,

Maldacena-Moore-Seiberg 2001, Belov-Moore 2005, Hartnoll 2006]

[Horowitz-Srednicki]: coupling to string sources  $\Delta S = \int_{\Gamma'} B^I$   
computes linking # of conjugate species of worldsheets  $\Gamma'$ .





Alexander Monin,

Massimo Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

## A simple topological field theory in 4+1 dimensions

Consider 2-forms  $B_{MN}$  in 4 + 1 dimensions, with action

$$S[B] = \frac{K_{IJ}}{2\pi} \int_{\mathbb{R} \times \Sigma} B^I \wedge dB^J$$

In  $4\ell + 1$  dims,  $K$  is a skew-symmetric integer  $2N_B \times 2N_B$  matrix.


Note:  $B \wedge dB = \frac{1}{2} d(B \wedge B)$ .

Independent of choice of metric on  $\mathbb{R} \times \Sigma_{2p}$ .

Related models studied in: [Horowitz 1989, Blau et al 1989, Witten 1998,

Maldacena-Moore-Seiberg 2001, Belov-Moore 2005, Hartnoll 2006]

[Horowitz-Srednicki]: coupling to string sources  $\Delta S = \int_{\Gamma'} B^I$   
computes linking # of conjugate species of worldsheets  $\Gamma'$ .





Alexander Monin,

cesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, n \rightarrow \text{st.}$

## A simple topological field theory in 4+1 dimensions

Consider 2-forms  $B_{MN}$  in 4 + 1 dimensions, with action

$$S[B] = \frac{K_{IJ}}{2\pi} \int_{\mathbb{R} \times \Sigma} B^I \wedge dB^J$$

In  $4\ell + 1$  dims,  $K$  is a skew-symmetric integer  $2N_B \times 2N_B$  matrix.


Note:  $B \wedge dB = \frac{1}{2} d(B \wedge B)$ .

Independent of choice of metric on  $\mathbb{R} \times \Sigma_{2p}$ .

Related models studied in: [Horowitz 1989, Blau et al 1989, Witten 1998,

Maldacena-Moore-Seiberg 2001, Belov-Moore 2005, Hartnoll 2006]

[Horowitz-Srednicki]: coupling to string sources  $\Delta S = \int_{\Gamma'} B^I$   
 computes linking # of conjugate species of worldsheets  $\Gamma'$ .



Simplest case ( $N_B = 1$ ) is realized in IIB strings on  $AdS_5 \times S^5$ ,

$B \equiv B_{NSNS}$ ,  $C \equiv C_{RR}$ :

$$S = \frac{1}{2\pi} \int_{AdS_5 \times S^5} F_{RR}^{(5)} \wedge B \wedge dC = \frac{N}{2\pi} \int_{\mathbb{R} \times \Sigma} B \wedge dC$$

Crucially, B S-duality acts by  $B \rightarrow B + dC$



Alexander Morion,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, n \rightarrow \text{st.}$

## Strategy

1. Solve the model – when is it an EFT for an SPT state?
2. Identify the edge states, and the symmetry  $G$  protecting them.  
(Whatever we get is surface-only with respect to  $G$ .)



Alexander Morita,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$\mu, \nu \rightarrow \text{st.}$

## Strategy

1. Solve the model – when is it an EFT for an SPT state?

Answer: when  $\text{Pfaff} K = 1$ .

2. Identify the edge states, and the symmetry  $G$  protecting them.

(Whatever we get is surface-only with respect to  $G$ .)

Answer: in the simplest realization, the edge theory is ordinary Maxwell theory, but with manifest electric-magnetic duality  $(\vec{E}, \vec{B}) \rightarrow (\vec{B}, -\vec{E})$ .



Alexander Morion,

cesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, d, k \rightarrow \text{s.c.c.}$

$\mu, \nu \rightarrow \text{s.t.}$

## Strategy

1. Solve the model – when is it an EFT for an SPT state?

Answer: when  $\text{Pfaff}K = 1$ .

2. Identify the edge states, and the symmetry  $G$  protecting them.

(Whatever we get is surface-only with respect to  $G$ .)

Answer: in the simplest realization, the edge theory is ordinary Maxwell theory, but with manifest electric-magnetic duality  $(\vec{E}, \vec{B}) \rightarrow (\vec{B}, -\vec{E})$ .



Alexander Monin,

Vincenzo Riva

$$a, b, c \rightarrow \text{C.L.}$$

$$u, v, k \rightarrow \text{S.L.}$$

$$m, v, \rightarrow \text{st.}$$

## All-fermion electrodynamics

So far, we've discussed 'pure' U(1) gauge theory (free).

A stronger obstruction can be found by adding matter

(Ends of strings are electric and magnetic charges)

Manifest duality symmetry

$\Rightarrow e, m$  and dyon  $\epsilon \equiv em$

must have the same statistics.

$$\begin{pmatrix} q_e \\ q_m \end{pmatrix} \rightarrow \underbrace{\begin{pmatrix} a & b \\ c & d \end{pmatrix}}_{\in \text{SL}(2, \mathbb{Z})} \begin{pmatrix} q_e \\ q_m \end{pmatrix}$$



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, v_i \rightarrow \text{st.}$

## All-fermion electrodynamics

So far, we've discussed 'pure' U(1) gauge theory (free).

A stronger obstruction can be found by adding matter

(Ends of strings are electric and magnetic charges)

Manifest duality symmetry  
 $\Rightarrow e, m$  and dyon  $\epsilon \equiv em$   
must have the same statistics.

$$\begin{pmatrix} q_e \\ q_m \end{pmatrix} \rightarrow \underbrace{\begin{pmatrix} a & b \\ c & d \end{pmatrix}}_{\in \text{SL}(2, \mathbb{Z})} \begin{pmatrix} q_e \\ q_m \end{pmatrix}$$



Alexander Monin,  
 Francesco Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, n \rightarrow \text{st.}$

## All-fermion electrodynamics

So far, we've discussed 'pure' U(1) gauge theory (free).

A stronger obstruction can be found by adding matter

(Ends of strings are electric and magnetic charges)

Manifest duality symmetry

$\Rightarrow e, m$  and dyon  $\epsilon \equiv em$

must have the same statistics.


- If  $e, m$  are bosons,  $\epsilon$  is a fermion!

'spin from isospin': [Jackiw-Rebbi]

$$\psi(x_1, x_2) = e^{i\varphi} \psi(x_2, x_1)$$

$$\varphi = e \int_0^\pi d\varphi \underbrace{\mathcal{A}_\varphi(\theta = \frac{\pi}{2}, \varphi)}_{\text{Dirac monopole field}} \stackrel{\text{Dirac}}{=} \pi$$

$$\begin{pmatrix} q_e \\ q_m \end{pmatrix} \rightarrow \underbrace{\begin{pmatrix} a & b \\ c & d \end{pmatrix}}_{\in \text{SL}(2, \mathbb{Z})} \begin{pmatrix} q_e \\ q_m \end{pmatrix}$$





Alexander Monin,

cesco Riva

$a, b, c \rightarrow \text{c.l.}$

$u, v, k \rightarrow \text{s.l.}$

$m, v, \rightarrow \text{st.}$

$da$

## All-fermion electrodynamics

So far, we've discussed 'pure' U(1) gauge theory (free).

A stronger obstruction can be found by adding matter

(Ends of strings are electric and magnetic charges)

Manifest duality symmetry

$\Rightarrow e, m$  and dyon  $\epsilon \equiv em$

must have the same statistics.


- If  $e, m$  are bosons,  $\epsilon$  is a fermion!

'spin from isospin': [Jackiw-Rebbi]

$$\psi(x_1, x_2) = e^{i\varphi} \psi(x_2, x_1)$$

$$\varphi = e \int_0^\pi d\varphi \underbrace{\mathcal{A}_\varphi(\theta = \frac{\pi}{2}, \varphi)}_{\text{Dirac monopole field}} \stackrel{\text{Dirac}}{=} \pi$$

$$\begin{pmatrix} q_e \\ q_m \end{pmatrix} \rightarrow \underbrace{\begin{pmatrix} a & b \\ c & d \end{pmatrix}}_{\in \text{SL}(2, \mathbb{Z})} \begin{pmatrix} q_e \\ q_m \end{pmatrix}$$





Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, n \rightarrow \text{S.L.}$

## All-fermion electrodynamics

So far, we've discussed 'pure' U(1) gauge theory (free).

A stronger obstruction can be found by adding matter

(Ends of strings are electric and magnetic charges)

Manifest duality symmetry

$\Rightarrow e, m$  and dyon  $\epsilon \equiv em$

must have the same statistics.


- If  $e, m$  are bosons,  $\epsilon$  is a fermion!

'spin from isospin': [Jackiw-Rebbi]

$$\psi(x_1, x_2) = e^{i\varphi} \psi(x_2, x_1)$$

$$\varphi = e \int_0^\pi d\varphi \underbrace{\mathcal{A}_\varphi(\theta = \frac{\pi}{2}, \varphi)}_{\text{Dirac monopole field}} \stackrel{\text{Dirac}}{=} \pi$$

$$\begin{pmatrix} q_e \\ q_m \end{pmatrix} \rightarrow \underbrace{\begin{pmatrix} a & b \\ c & d \end{pmatrix}}_{\in \text{SL}(2, \mathbb{Z})} \begin{pmatrix} q_e \\ q_m \end{pmatrix}$$





Alexander Morion,

Francesco Riva


$a, b, c \rightarrow \text{c.l.}$   
 $u, v, k \rightarrow \text{s.c.l.}$   
 $m, v, \rightarrow \text{st.}$

## All-fermion electrodynamics

This is evidence for a 4+1d SPT  
*with no symmetry* (analogous to Kitaev's  $E_8$  state).

[work in progress:] Coupled-layer construction

following [Wang-Senthil] for all-fermion TC  
 produces trivial bosonic bulk, correct edge.



Idea: 'dyon string condensation' (like [Metlitski-Kane-Fisher])

- Each layer is ordinary electrodynamics with bosonic charges.
- $b_i \equiv \epsilon_i^\dagger m_{i+1} \epsilon_{i+2}$  are mutually-local bosons.
- Condensing  $b_i$  (obliquely) confines  $a_{i+1}$ ,  $i+1 = 2 \dots N-1$ .
- At top layer:  $m_1 \epsilon_2, \epsilon_1^\dagger m_1 \epsilon_2, \epsilon_1^\dagger$  survive, are fermions, are electron, monopole & dyon of  $U(1)_{\text{odd}}$ .

$\exists$  a 4+1d local lattice model which realizes this construction.

In the bulk, continuous the BdC theory with gapped string matter.



Alexander Morin,

Francesco Riva


$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, n \rightarrow \text{st.}$

## All-fermion electrodynamics

This is evidence for a 4+1d SPT  
*with no symmetry* (analogous to Kitaev's  $E_8$  state).

[work in progress] Coupled-layer construction

following [Wang-Senthil] for all-fermion TC  
 produces trivial bosonic bulk, correct edge.



Idea: 'dyon string condensation' (like [Metlitski-Kane-Fisher])

- Each layer is ordinary electrodynamics with bosonic charges.
- $b_i \equiv e_i^\dagger m_{i+1} \epsilon_{i+2}$  are mutually-local bosons.
- Condensing  $b_i$  (obliquely) confines  $a_{i+1}$ ,  $i+1 = 2 \dots N-1$ .
- At top layer:  $m_1 \epsilon_2, e_1^\dagger m_1 \epsilon_2, e_1^\dagger$  survive, are fermions, are electron, monopole & dyon of  $U(1)_{\text{odd}}$ .

$\exists$  a 4+1d local lattice model which realizes this construction.

In the bulk, content is the BdC theory with gapped string matter.



Alexander Morion,  
 Francesco Riva


$a, b, c \rightarrow \text{c.l.}$   
 $u, v, k \rightarrow \text{s.l.}$   
 $m, v, \rightarrow \text{st.}$

## All-fermion electrodynamics

This is evidence for a 4+1d SPT  
 with *no symmetry* (analogous to Kitaev's  $E_8$  state).

[work in progress:] Coupled-layer construction

following [Wang-Senthil] for all-fermion TC  
 produces trivial bosonic bulk, correct edge.



Idea: 'dyon string condensation' (like [Metlitski-Kane-Fisher])

- Each layer is ordinary electrodynamics with bosonic charges.
- $b_i \equiv \epsilon_i^\dagger m_{i+1} \epsilon_{i+2}$  are mutually-local bosons.
- Condensing  $b_i$  (obliquely) confines  $a_{i+1}$ ,  $i+1 = 2 \dots N-1$ .
- At top layer:  $m_1 \epsilon_2, \epsilon_1^\dagger m_1 \epsilon_2, \epsilon_1^\dagger$  survive, are fermions, are electron, monopole & dyon of  $U(1)_{\text{odd}}$ .

$\exists$  a 4+1d local lattice model which realizes this construction.

In the bulk, continuum: this is the P/C theory with gapped string matter.



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $\mu, \nu \rightarrow \text{st.}$

## Dyon string condensation

If we did this:  $\rightarrow$

$$\Delta H = V \sum_i (|\tilde{b}_i|^2 - v)^2, \quad \tilde{b}_i \equiv e_i^\dagger e_{i+2} = v e^{ia_{i,i+2}}$$

would higgs  $\prod_i U(1)_i \rightarrow U(1)_{\text{even}} \times U(1)_{\text{odd}}$ .

4 + 1d Maxwell theory with  $G = U(1)_{\text{even}} \times U(1)_{\text{odd}}$ , bulk photons.

Can dualize to 2-form potentials:  $f^{o/e} = da^{o/e} = \star dC^{o/e}$

$$S = \sum_{\alpha=o,e} \int_{5d} \left( \frac{1}{g_\alpha^2} dC^\alpha \wedge \star dC^\alpha + C^\alpha \wedge \star J_m^\alpha \right).$$

Magnetic flux tubes of broken  $U(1)$ s collimate monopoles into *monopole strings*.

If instead we do *this*:  $\rightarrow$



$b_i \equiv \epsilon_i^\dagger m_{i+1} \epsilon_{i+2}$  are mutually-local bosons.

$\Delta H = V \sum_i (|b_i|^2 - v)^2$ . Condensing  $b_i$  (obliquely) *confines*  $a_{i+1}, i+1 = 2 \dots N-1$ .

Binds monopole strings of  $a^{e/o}$  to electric flux lines of  $a^{o/e}$ .

This is the effect of the additional term

$$\Delta S = \int \frac{1}{4\pi} C^e \wedge C^o.$$






Francesco Riva

$a, b, c \rightarrow C.L.$   
 $l, i, k \rightarrow S.L.L$   
 $\mu, v \rightarrow st.$

 $da$ 

## A solvable coupled-island construction of SPT states in $2+1$ dimensions


$$1 - \frac{-6 \ln(2x)}{4} \cdot \frac{1}{2x} + \dots$$



Alexander Morion,

Massimo Riva


$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{S.L.}$

$m, v, \rightarrow \text{st.}$

da

# A solvable coupled-island construction of SPT states in $2 + 1$ dimensions



$$\left( \frac{1}{2} \partial_\mu \phi \partial^\mu \phi \right) - \left( \frac{1}{4} \epsilon_{\mu\nu\lambda} \partial^\mu \phi \partial^\nu \phi \partial^\lambda \phi + \dots \right)$$



Alexander Morita,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, v, \rightarrow \text{st.}$

## An SPT machine

A lot of effort has been put into classifying SPT states.

Fewer explicit constructions exist.

Useful e.g. for understanding the phase transitions between them, and the topologically-ordered states that result upon gauging (subgroups of)  $G$ .



Alexander Monin,

Massimo Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, n, l \rightarrow \text{st.}$

## An SPT machine

A lot of effort has been put into classifying SPT states.

Fewer explicit constructions exist.

Useful e.g. for understanding the phase transitions between them, and the topologically-ordered states that result upon gauging (subgroups of)  $G$ .



Alexander Morion,

Vincenzo Riva

$$\begin{aligned} a, b, c &\rightarrow \text{c.c.} \\ u, v, k &\rightarrow \text{s.c.c.} \\ m, v, &\rightarrow \text{st.} \end{aligned}$$

## An SPT machine

A lot of effort has been put into classifying SPT states.

Fewer explicit constructions exist.

Useful e.g. for understanding the phase transitions between them, and the topologically-ordered states that result upon gauging (subgroups of)  $G$ .

Here: 2+1d  $\mathbb{Z}_N$  paramagnets.

Virtues of our construction:

- ▶ Translation invariance not required. (Often translation invariance can protect an otherwise unprotected edge.)
- ▶ Uniform construction of domain wall operators.  
→ [Levin-Gu] braiding statistics proof of nontriviality
- ▶ Illuminates connections between the few existing examples:  
[Levin-Gu ( $\mathbb{Z}_2$ ), Chen-Liu-Wen ( $\mathbb{Z}_2$ ), Chen-Gu-Liu-Wen (mysterious general formula?)]
- ▶ The 'duality' method of [Levin-Gu] was not available: gauging the bulk symmetry provides a (simpler?) construction of recent 'generalized string-net models' [Lin-Levin].



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{S.L.}$

$m, v \rightarrow \text{st.}$


da

## Non-onsite symmetry

An anomalous symmetry can be realized in a lattice model if it is *not on-site*: its action on one site depends on others.

This means you can't gauge it just by coupling to link variables (without coarse-graining first).

(Like chiral symmetry with staggered fermions.)





Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$

$u, v, k \rightarrow \text{S.L.}$

$m, v, \rightarrow \text{st.}$


da

## Non-onsite symmetry

An anomalous symmetry can be realized in a lattice model if it is *not on-site*: its action on one site depends on others.

This means you can't gauge it just by coupling to link variables (without coarse-graining first).

(Like chiral symmetry with staggered fermions.)





Alexander Morion,

Massimo Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, v, \rightarrow \text{st.}$

## Non-onsite symmetry

An anomalous symmetry can be realized in a lattice model if it is *not on-site*: its action on one site depends on others.

This means you can't gauge it just by coupling to link variables (without coarse-graining first).

For example, at the edge of the Levin-Gu  $\mathbb{Z}_2$  paramagnet,


$$S = \prod_J x_J \prod_J i^{\frac{1}{2}(1-Z_J Z_{J+1})} = \prod_J x_J \cdot i^{\text{number of domain walls}}$$

A non-onsite symmetry  $S$  is nontrivial if  $S \neq U \prod_J s_J U^\dagger$  with  $U$  a *local symmetric unitary* (unitary evolution by a symmetric  $H$ ).  
 How to tell?? We will find a practical criterion below.

Focus on  $\mathbb{Z}_N$  spins at each site

$$P_n(Z) = \prod_J (Z_J - \omega^n), \quad \omega = e^{\frac{2\pi i}{N}}, \quad Z|n\rangle = \omega^n |n\rangle, \quad n = 0..N-1 \pmod{N}$$

(Like chiral symmetry with staggered fermions.)





Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, v, \rightarrow \text{st.}$

## Non-onsite symmetry

An anomalous symmetry can be realized in a lattice model if it is *not on-site*: its action on one site depends on others.

This means you can't gauge it just by coupling to link variables (without coarse-graining first).

For example, at the edge of the Levin-Gu  $\mathbb{Z}_2$  paramagnet,

$$S = \prod_J \mathbf{x}_J \prod_J i^{\frac{1}{2}(1-Z_J Z_{J+1})} = \prod_J \mathbf{x}_J \cdot i^{\text{number of domain walls}}$$


A non-onsite symmetry  $S$  is nontrivial if  $S \neq U \prod_J s_J U^\dagger$  with  $U$  a *local symmetric unitary* (unitary evolution by a symmetric  $H$ ).  
 How to tell?? We will find a practical criterion below.

Focus on  $\mathbb{Z}_N$  spins at each site:

$$\mathbf{XZ} = \omega \mathbf{ZX}, \quad \omega \equiv e^{\frac{2\pi i}{N}} \quad \mathbf{Z}|n\rangle = \omega^n |n\rangle, .$$

$$\mathbf{P}_n(\mathbf{Z}) \equiv |n\rangle\langle n|, \quad \mathbf{X}|n\rangle = |n-1\rangle, \quad n = 0..N-1 \pmod{N}$$

(Like chiral symmetry with staggered fermions.)





Alexander Monin,

Francesco Riva


$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

## An SPT machine

Given desired action of  
non-onsite symmetry on edge:






Alexander Morin,

Francesco Riva


$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, v, \rightarrow \text{st.}$

## An SPT machine

Given desired action of  
non-onsite symmetry on edge:



Couple together 'bags':  
(Inspired by CZX model for  $\mathbb{Z}_2$  [Chen-Liu-Wen, Swingle].)



Think of each bag as a site.



Alexander Morion,


cesco Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, v, \rightarrow \text{st.}$


da

## An SPT machine

Given desired action of  
non-onsite symmetry on edge:



Couple together 'bags':  
(Inspired by CZX model for  $\mathbb{Z}_2$  [Chen-Liu-Wen, Swingle].)



Think of each bag as a site.



Alexander Morion,


Francesco Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, v_i \rightarrow \text{st.}$


da

## An SPT machine

Given desired action of  
non-onsite symmetry on edge:



Couple together 'bags':  
(Inspired by CZX model for  $\mathbb{Z}_2$  [Chen-Liu-Wen, Swingle].)



Think of each bag as a site.



Alexander Monin,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$


$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

da

## An SPT machine

Given desired action of  
non-onsite symmetry on edge:




Couple together 'bags':

(Inspired by CZX model for  $\mathbb{Z}_2$  [Chen-Liu-Wen, Swingle].)


$$H_{\text{CZX}} = - \sum_{\square} b_{\square} + h.c.$$

$$b_{\square} \equiv \text{XXXX} \mathcal{P}. \quad [b_{\square}, b_{\square'}] = 0$$

$$|\psi\rangle = \prod_{\square} \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} |nnnn\rangle$$



Think of each bag as a site.





Alexander Morion,


cesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, v_i \rightarrow \text{st.}$

da

## An SPT machine

Given desired action of  
non-onsite symmetry on edge:




Couple together 'bags':

(Inspired by CZX model for  $\mathbb{Z}_2$  [Chen-Liu-Wen, Swingle].)


$$H_{\text{CZX}} = - \sum_{\square} b_{\square} + h.c.$$

$$b_{\square} \equiv \text{XXXXP}. \quad [b_{\square}, b_{\square'}] = 0$$

$$|gs\rangle = \prod_{\square} \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1}$$



Think of each bag as a site.





Alexander Morion,


cesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, n \rightarrow \text{st.}$

## Symmetry-protected edge states


Note:  $\mathbf{H}$  doesn't know about  $p$ ;  $\mathbf{S}$  does: ( $[\mathbf{S}, \mathbf{H}_{\text{CZx}}] = 0$ )

$$\mathbf{S} = \prod_j \mathbf{x}_j \left( \prod_{\text{bags}} \prod_j C_p(\mathbf{z}_j, \mathbf{z}_{j+1}) \right)$$



Not onsite on edge:  $\rightarrow$

Rough edge realizes the  
desired  $\mathbf{S}$  on edge modes:





Alexander Morion,


Francesco Riva

$a, b, c \rightarrow \text{C.L.}$   
 $u, v, k \rightarrow \text{S.L.}$   
 $m, n \rightarrow \text{st.}$

## Symmetry-protected edge states


Note:  $\mathbf{H}$  doesn't know about  $p$ ;  $\mathbf{S}$  does:  $([\mathbf{S}, \mathbf{H}_{\text{CZx}}] = 0)$

$$\mathbf{S} = \prod_j \mathbf{x}_j \left( \prod_{\text{bags}} \prod_j C_p(\mathbf{z}_j, \mathbf{z}_{j+1}) \right)$$



Not onsite on edge:  $\rightarrow$

Rough edge realizes the desired  $\mathbf{S}$  on edge modes:



Claim of robustness: perturbing  $\mathbf{H}_{\text{CZx}}$  by terms respecting  $\mathbf{S}$ , you cannot remove this edge stuff.

i.e. no local, symmetric unitary can make  $|gs\rangle_{\text{edge}}$  a product state. (Gapless or symmetry-breaking degeneracy.)

Proving (?): requires bipartite graph of connections between bags.

No draw lattices for simplicity of drawing, but translation invariance is not necessary.



## Bag link phases

$$S = \prod_j x_j \prod_{\text{bags}} \prod_j C_p(z_j, z_{j+1})$$

Wanted: A unitary operator  $C(1, 2) \equiv C(z_1, z_2)$  on two  $\mathbb{Z}_N$ -valued variables which satisfies the following three simple-looking conditions:

- group law :  $S^N = \prod_j (z_j^\dagger z_{j+1})^p$

only for closed loops:  $= 1$



Alexander Morion,

Massimo Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

## Bag link phases

$$S = \prod_j x_j \prod_{\text{bags}} \prod_j C_p(z_j, z_{j+1})$$

Wanted: A unitary operator  $C(1, 2) \equiv C(z_1, z_2)$  on two  $\mathbb{Z}_N$ -valued variables which satisfies the following three simple-looking conditions:

- group law :  $S^N = \prod_j (z_j^\dagger z_{j+1})^p$

only for closed loops:  $= 1$



Alexander Morion,  
 Francesco Riva


$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, n \rightarrow \text{st.}$

## Bag link phases

$$S = \prod_j x_j \prod_{\text{bags}} \prod_j C_p(z_j, z_{j+1})$$

Wanted: A unitary operator  $C(1, 2) \equiv C(Z_1, Z_2)$  on two  $\mathbb{Z}_N$ -valued variables which satisfies the following three simple-looking conditions:

- group law :  $S^N = \prod_j (z_j^\dagger z_{j+1})^p$   
 only for closed loops:  $= 1$
- gappability :  $C_p(1, 2) C_{-p}(1, 2) = 1$
- non-triviality (flux braiding) :



$$\frac{X_1^{-k_2} X_3^{k_1} C_{-p}^{k_2}(23) X_3^{-k_1}}{C_{-p}^{k_2}(23)} \stackrel{?}{=} e^{2\pi i \frac{2p}{N} k_1 k_2}.$$



Alexander Morion,


Francesco Riva

$a, b, c \rightarrow \text{L.L.}$   
 $u, v, k \rightarrow \text{S.L.L.}$   
 $m, v, \rightarrow \text{st.}$

## A uniform construction of domain-wall operators

$$W(R) \equiv \prod_{j \in R \setminus \text{last col}} \mathbf{x}_j \prod_{\text{bags in } R} \mathcal{P} \cdot \prod_{j \in \text{bag}} C_p(\mathbf{z}_j, \mathbf{z}_{j+1})$$

Acts like **S** in the interior of  $R$ .  
 Threads  $2\pi/N$ -flux along its boundary.  
 Becomes the string operator in the  
 topologically-ordered model with  $G$  gauged  
 and deconfined.





Alexander Morion,  
 Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $n, v, \rightarrow \text{st.}$

## Braiding of flux insertions

[Levin-Gu]

$W_R$  generate symmetries:

$$[H, W_R] = 0.$$

If domain walls intersect *once*:

$$W_{R_1}^{k_1} W_{R_2}^{k_2} = e^{2\pi i k_1 k_2 \frac{2p}{N^2}} W_{R_2}^{k_2} W_{R_1}^{k_1}$$



(annoying fine print:

this formula works for  $k_1 k_2 \propto N$ )

This must be represented on  
 groundstates

(in fact, the whole spectrum)

$\Rightarrow$  nontrivial edge spectrum.





Alexander Morion,

Massimo Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, v, \rightarrow \text{st.}$

da

## Braiding of flux insertions

[Levin-Gu]

$W_R$  generate symmetries:

$$[H, W_R] = 0.$$

If domain walls intersect *once*:

$$W_{R_1}^{k_1} W_{R_2}^{k_2} = e^{2\pi i k_1 k_2 \frac{2p}{N^2}} W_{R_2}^{k_2} W_{R_1}^{k_1}$$



(annoying fine print:

this formula works for  $k_1 k_2 \propto N$ )

This must be represented on  
groundstates

(in fact, the whole spectrum)

$\Rightarrow$  nontrivial edge spectrum.





Alexander Morion,

Francesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

## Braiding of flux insertions

[Levin-Gu]

$W_R$  generate symmetries:

$$[H, W_R] = 0.$$

If domain walls intersect *once*:

$$W_{R_1}^{k_1} W_{R_2}^{k_2} = e^{2\pi i k_1 k_2 \frac{2p}{N^2}} W_{R_2}^{k_2} W_{R_1}^{k_1}$$



(annoying fine print:

this formula works for  $k_1 k_2 \propto N$ )

This must be represented on  
groundstates

(in fact, the whole spectrum)

$\Rightarrow$  nontrivial edge spectrum.





Alexander Morion,

cesco Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

da

## Braiding of flux insertions

$W_R$  generate symmetries:

$$[H, W_R] = 0.$$

If domain walls intersect *once*:

$$W_{R_1}^{k_1} W_{R_2}^{k_2} = e^{2\pi i k_1 k_2 \frac{2p}{N^2}} W_{R_2}^{k_2} W_{R_1}^{k_1}$$



(annoying fine print:

this formula works for  $k_1 k_2 \propto N$ )

This must be represented on  
groundstates

(in fact, the whole spectrum)

$\Rightarrow$  nontrivial edge spectrum.





Alexander Morion,  
 Francesco Riva

$a, b, c \rightarrow \text{c.c.}$   
 $u, v, k \rightarrow \text{s.c.c.}$   
 $m, v, \rightarrow \text{st.}$

## Coarse-graining transformation


step 1: change basis to on-site symmetry:

$$\mathbf{U} \mathbf{S}_1 \mathbf{U}^\dagger = \prod_j \mathbf{X}_j \quad (\mathbf{U} \text{ known but, so far, ugly.})$$

Weirdness of  $\mathbf{U}$ : it's a local unitary, but not continuously connected to  $\mathbb{1}$  by local symmetric unitaries.

In this basis, easy to gauge. Alternative[Swingle]: diagonalize action on bags.







Alexander Morion,

Francesco Riva

$a, b, c \rightarrow c.c.$

$u, v, k \rightarrow s.c.c.$

$m, v, \rightarrow s.t.$

da

## Non-onsite edge symmetry revisited

Chern-Simons description [Lu-Vishwanath]:

Edge is non-chiral boson, only the left-mover carries the  $\mathbb{Z}_N$ .



Alexander Morion,

Massimo Riva

$a, b, c \rightarrow \text{c.c.}$

$u, v, k \rightarrow \text{s.c.c.}$

$m, v, \rightarrow \text{st.}$

## Questions

- ▶ For  $G = U(1)$ , gauging the symmetry produces a model with  $c_L \neq c_R$ , but the model should still be solvable.  
Some tension with theorems of [Kitaev, honeycomb paper; Lin-Levin].
- ▶ We have not yet made precise the connection to group cohomology.  
The condition on the link phases that the DW commutator is a c-number should be the cocycle condition. Is it?
- ▶ Origin of bipartite restriction?!?  
In the continuum, there is no difference between  $p \rightarrow -p$  and orientation reversal.
- ▶ Non-abelian  $G$ ?
- ▶ 3d?
- ▶ What is anomalous about the all-fermion electrodynamics?



