

Title: Are the recent IceCube events hinting at O(100) TeV decaying dark matter?

Date: May 22, 2014 01:00 PM

URL: <http://pirsa.org/14050118>

Abstract: The IceCube detector has recently reported the observation of 28 events at previously unexplored energies. While the statistics of the observed events are still low, these events hint at the existence of a neutrino flux over and above the atmospheric neutrino background. We investigate the possibility that a significant component of the additional neutrino flux originates due to the decay of a very heavy dark matter (VHDM) particle via several possible channels into standard model particles. We show that a combination of a power law astrophysical neutrino spectrum and the neutrino flux from the decay of a DM species of mass in the range 150-400 TeV improves the fit to the observed neutrino events than that obtained from a best-fit astrophysical flux alone. Assuming the existence of an astrophysical background described by the IC best-fit, we also show that, for the decay of even heavier DM particles ($m_{\text{DM}} \sim 1$ PeV), the same observations impose significant constraints on the decay lifetimes. Allowing the astrophysical flux normalization to vary leads to modifications of these limits; however, there is still a range of dark matter mass and lifetime that is excluded by the IC results.

**ARE THE RECENT ICECUBE EVENTS HINTING AT $O(100)$ TeV DECAYING
DARK MATTER?**


Atri Bhattacharya

Talk at Perimeter Institute


22 May 2014




THE NEUTRINO SKY... TO THE HIGHEST ENERGIES



THE NEUTRINO SKY... TO THE HIGHEST ENERGIES



THE NEUTRINO SKY... TO THE HIGHEST ENERGIES




THE NEUTRINO SKY... TO THE HIGHEST ENERGIES

Probe highest energy neutrino production mechanisms

Existence of tiny non-std physical effects (LV, etc.)

Hunting for astrophysical point objects (AGN, etc.) using neutrinos



Probe neutrino oscillation at highest energies

DM annihilation at the galactic centre


Indirect search for very heavy DM decay

THE NEUTRINO SKY... TO THE HIGHEST ENERGIES

Probe highest energy neutrino production mechanisms

Existence of tiny non-std physical effects (LV, etc.)

Hunting for astrophysical point objects (AGN, etc.) using neutrinos



Probe neutrino oscillation at highest energies

DM annihilation at the galactic centre


Indirect search for very heavy DM decay

THE NEUTRINO SKY... TO THE HIGHEST ENERGIES

Probe highest energy neutrino production mechanisms

Existence of tiny non-std physical effects (LV, etc.)

Hunting for astrophysical point objects (AGN, etc.) using neutrinos




Probe neutrino oscillation at highest energies

DM annihilation at the galactic centre

Indirect search for very heavy DM decay

NEUTRINOS @ HIGHEST ENERGIES: HOW CATCH 'EM



Main issues with detection


- Extremely **low incident fluxes**
- **Huge incident energies** – reconstruction requires voluminous detectors
- **Flavour** discrimination?

Solution?
Km³ Detectors

Km³ detectors

- Trap **high fraction of incident neutrino fluxes**
- Proper **energy and direction (for tracks) reconstruction** of large event signature tracks
 - Big enough to **contain hadronic/em cascades**
 - Possibility of detection of **double-bang signatures from incident ν_τ 's**

NEUTRINOS @ HIGHEST ENERGIES: HOW CATCH 'EM



Main issues with detection


- Extremely **low incident fluxes**
- **Huge incident energies** – reconstruction requires voluminous detectors
- **Flavour** discrimination?

Solution?
Km³ Detectors

Km³ detectors

- Trap **high fraction of incident neutrino fluxes**
- Proper **energy and direction (for tracks) reconstruction** of large event signature tracks
 - Big enough to **contain hadronic/em cascades**
 - Possibility of detection of **double-bang signatures from incident ν_{τ} 's**


PRESENT SETUP FOR UHE ν DETECTION



IceCube

- Operational since 2010
 - Full exposure since Dec. 2011
- Capable of flavour discrimination
 - Limited to detection of three distinct event signatures
- Excellent energy reconstruction
 - < 10% for contained cascades
 - ~ 30% for tracks with contained vertices
- Good direction reconstruction
 - Up to 1° for tracks
 - $\sim 30^\circ$ for cascades
- Designed to run (minimal op. cost) for 10+ yrs
- Collected 28 UHE events in 662 days of run-time
 - 2 events at PeV+ energies
 - Recently reported 9 more events, making total event number 37 over 988 days

PRESENT SETUP FOR UHE ν DETECTION




IceCube

- Operational since 2010
 - Full exposure since Dec. 2011
- Capable of flavour discrimination
 - Limited to detection of three distinct event signatures
- Excellent energy reconstruction
 - < 10% for contained cascades
 - ~ 30% for tracks with contained vertices
- Good direction reconstruction
 - Up to 1° for tracks
 - ~30° for cascades
- Designed to run (minimal op. cost) for 10+ yrs
- Collected 28 UHE events in 662 days of run-time
 - 2 events at PeV+ energies
 - Recently reported 9 more events, making total event number 37 over 988 days

FLAVOUR @ IC

Muon Track


$\nu_{\mu} N CC$




$\nu_{e,\tau} N CC$
 $\nu N NC$

$\nu_{\tau} N CC (\geq 1 \text{ PeV})$


Cascades



Double Bang



RECONSTRUCTING EVENTS @ ICECUBE




Muon Tracks

Charged current interaction
of the muon-neutrino

Clear tracks and excellent
direction reconstruction

Energy reconstruction is
indirect – energy loss along
track




Cascades

Charged current interaction
of the electron-neutrino
and tau-neutrino

Neutral current
interactions of all flavours

Excellent energy but poorer
direction reconstruction

RECONSTRUCTING EVENTS @ ICECUBE




Muon Tracks

Charged current interaction
of the muon-neutrino

Clear tracks and excellent
direction reconstruction

Energy reconstruction is
indirect – energy loss along
track



Cascades

Charged current interaction
of the electron-neutrino
and tau-neutrino

Neutral current
interactions of all flavours

Excellent energy but poorer
direction reconstruction

INCIDENT FLUXES FROM STD. THEORY

- Diffuse flux from all-sky astrophysical sources

- Expected to follow a power-law spectrum


- Fermi 1st order shocks $\rightarrow \alpha = 2.0$

- Normalisation fixed by observational best-fits

$$\Phi_\nu \propto E^{-\alpha}$$

- Neutrinos in sources predominantly from pion decays

- Std. oscillation \rightarrow incident flavour 1:1:1 at earth



INCIDENT FLUXES FROM STD. THEORY

- Diffuse flux from all-sky astrophysical sources

- Expected to follow a power-law spectrum


- Fermi 1st order shocks $\rightarrow \alpha = 2.0$

- Normalisation fixed by observational best-fits


$$\Phi_\nu \propto E^{-\alpha}$$

- Neutrinos in sources predominantly from pion decays

- Std. oscillation \rightarrow incident flavour 1:1:1 at earth




OBSERVATIONS @ IC [662 DAYS]



27 total events

- Two PeV+ cascades
 - Highest energy neutrino events ever observed
- Additional 18 lower energy cascades
- 7 track events
- Events from all sky
- No event from 300 TeV – 1 PeV


OBSERVATIONS @ IC [662 DAYS]



27 total events

- Two PeV+ cascades
 - Highest energy neutrino events ever observed
- Additional 18 lower energy cascades
- 7 track events
- Events from all sky
- No event from 300 TeV – 1 PeV


OBSERVATIONS @ IC [662 DAYS]



27 total events

- Two PeV+ cascades
 - Highest energy neutrino events ever observed
- Additional 18 lower energy cascades
- 7 track events
- Events from all sky
- No event from 300 TeV – 1 PeV

OBSERVATIONS @ IC [662 DAYS]



At least 3.6 σ signal over atmospheric neutrino background with 90% c.l. charm estimates


Best-fit largely consistent with E⁻² power flux up to 1.1 PeV...

$$E^2\Phi = 1.2_{0.8}^{1.6} \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$

...BUT

- Unexplained sharp drop above 1 PeV
- Lack of events within 300 TeV – 1 PeV
- Sub-100 TeV energy event numbers consistently higher than prediction from E⁻² flux

OBSERVATIONS @ IC [662 DAYS]



At least 3.6σ signal over atmospheric neutrino background with 90% c.l. charm estimates

Best-fit largely consistent with E^{-2} power flux up to 1.1 PeV...

$$E^2\Phi = 1.2_{0.8}^{1.6} \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$

...BUT


- Unexplained sharp drop above 1 PeV
- Lack of events within 300 TeV – 1 PeV
- Sub-100 TeV energy event numbers consistently higher than prediction from E^{-2} flux

THE NEUTRINO SKY... TO THE HIGHEST ENERGIES

Probe highest energy neutrino production mechanisms

Existence of tiny non-std physical effects (LV, etc.)

Hunting for astrophysical point objects (AGN, etc.) using neutrinos




Probe neutrino oscillation at highest energies

DM annihilation at the galactic centre

Indirect search for very heavy DM decay

OBSERVATIONS @ IC [662 DAYS]



At least 3.6 σ signal over atmospheric neutrino background with 90% c.l. charm estimates


Best-fit largely consistent with E⁻² power flux up to 1.1 PeV...

$$E^2\Phi = 1.2_{0.8}^{1.6} \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$

...BUT

- Unexplained sharp drop above 1 PeV
- Lack of events within 300 TeV – 1 PeV
- Sub-100 TeV energy event numbers consistently higher than prediction from E⁻² flux

OBSERVATIONS @ IC [662 DAYS]



At least 3.6σ signal over atmospheric neutrino background with 90% c.l. charm estimates

Best-fit largely consistent with E^{-2} power flux up to 1.1 PeV...

$$E^2\Phi = 1.2_{0.8}^{1.6} \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$

...BUT

- Unexplained sharp drop above 1 PeV
- Lack of events within 300 TeV – 1 PeV
- Sub-100 TeV energy event numbers consistently higher than prediction from E^{-2} flux

ADDITIONAL NEUTRINO FLUX COMPONENT?

- **An additional flux of neutrinos at 10 TeV – 100 TeV, over & above the astrophysical background?**
 - Normalisation to diffuse astrophysical flux fixed by PeV events ()
 - Lower normalisation allows for better match with lack of events in the high energy “well” (300 TeV – 1 PeV)
 - *Lower energy events, then, could be accounted for by a neutrino flux from an additional source*

AB, Sarcevic, Reno
arXiv:1403.1862

ADDITIONAL NEUTRINO FLUX COMPONENT?

- **An additional flux of neutrinos at 10 TeV – 100 TeV, over & above the astrophysical background?**
 - Normalisation to diffuse astrophysical flux fixed by PeV events ()
 - Lower normalisation allows for better match with lack of events in the high energy “well” (300 TeV – 1 PeV)
 - *Lower energy events, then, could be accounted for by a neutrino flux from an additional source*

AB, Sarcevic, Reno
arXiv:1403.1862

ADDITIONAL NEUTRINO FLUX COMPONENT?

Possible sources

- **NOT**
 - Atmospheric
 - Cosmogenic
- Heavy DM – $m_{\text{DM}} = O(100)$ TeV – annihilation?
 - Predominance of signal from galactic centre
 - Slight excess towards GC seen
 - For WIMP-like $\langle\sigma v\rangle^\dagger$, expected IC signal too low
 - 1 event in 50 yrs
- Heavy DM decay...

[†] For WIMP's, $\langle\sigma v\rangle \approx 10^{-27} \text{ cm}^3 \text{ s}^{-1}$

INCIDENT FLUXES FROM STD. THEORY

- Diffuse flux from all-sky astrophysical sources

- Expected to follow a power-law spectrum

- Fermi 1st order shocks $\rightarrow \alpha = 2.0$

- Normalisation fixed by observational best-fits


$$\left. \begin{array}{l} \text{Expected to follow a power-law spectrum} \\ \text{Fermi 1}^{\text{st}} \text{ order shocks } \rightarrow \alpha = 2.0 \\ \text{Normalisation fixed by observational best-fits} \end{array} \right\} \Phi_{\nu} \propto E^{-\alpha}$$

- Neutrinos in sources predominantly from pion decays

- Std. oscillation \rightarrow incident flavour 1:1:1 at earth

- Cosmogenic neutrinos ($E \geq 100$ PeV)

- Cosmic rays interacting with CMBR photons



ADDITIONAL NEUTRINO FLUX COMPONENT?

Possible sources

- **NOT**
 - Atmospheric
 - Cosmogenic
- Heavy DM – $m_{\text{DM}} = O(100)$ TeV – annihilation?
 - Predominance of signal from galactic centre
 - Slight excess towards GC seen
 - For WIMP-like $\langle\sigma v\rangle^\dagger$, expected IC signal too low
 - 1 event in 50 yrs
- Heavy DM decay...

[†] For WIMP's, $\langle\sigma v\rangle \approx 10^{-27} \text{ cm}^3 \text{ s}^{-1}$

ADDITIONAL NEUTRINO FLUX COMPONENT?

Possible sources


- **NOT**
 - Atmospheric
 - Cosmogenic
- Heavy DM – $m_{\text{DM}} = O(100)$ TeV – annihilation?
 - Predominance of signal from galactic centre
 - Slight excess towards GC seen
 - For WIMP-like $\langle\sigma v\rangle^\dagger$, expected IC signal too low
 - 1 event in 50 yrs
- Heavy DM decay...

[†] For WIMP's, $\langle\sigma v\rangle \approx 10^{-27} \text{ cm}^3 \text{ s}^{-1}$

HEAVY DM DECAY: FEATURES

- Same OoM contribution from galactic and extra-galactic source
 - Diffuse flux from 4π sky
 - Observable event rates predicted (comparable to IC b.f.)
 - Contrary to DM annihilation scenarios
 - Reasonable lifetimes for DM relic particles: $O(10^{27})$ s
 - Explains relic abundance, 27% of universe's energy content
 - Possible decay channels: ZZ , WW , $\tau+\tau^-$, $\mu+\mu^-$, $\nu\bar{\nu}$, $q\bar{q}$...

OBSERVATIONS @ IC [662 DAYS]



At least 3.6σ signal over atmospheric neutrino background with 90% c.l. charm estimates

Best-fit largely consistent with E^{-2} power flux up to 1.1 PeV...

$$E^2\Phi = 1.2_{0.8}^{1.6} \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$

...BUT

- Unexplained sharp drop above 1 PeV
- Lack of events within 300 TeV – 1 PeV
- Sub-100 TeV energy event numbers consistently higher than prediction from E^{-2} flux

DM DECAY: CONSIDERATIONS

- Simplicity:
 - Single DM species decaying with lifetimes $\sim O(10^{27})s$
 - Limit to study of scalar DM
 - Limit to two-body decays, single decay channel
 - Decays to std. model particle pairs
 - Z^0Z^0
 - W^+W^-
 - $\tau^+\tau^-$
 - $\mu^+\mu^-$


DM DECAY: CONSIDERATIONS

- Simplicity:
 - Single DM species decaying with lifetimes $\sim O(10^{27})s$
 - Limit to study of scalar DM
 - Limit to two-body decays, single decay channel
 - Decays to std. model particle pairs
 - Z^0Z^0
 - W^+W^-
 - $\tau^+\tau^-$
 - $\mu^+\mu^-$

DM DECAY: CONSIDERATIONS

- Simplicity:
 - Single DM species decaying with lifetimes $\sim O(10^{27})s$
 - Limit to study of scalar DM
 - Limit to two-body decays, single decay channel
 - Decays to std. model particle pairs
 - Z^0Z^0
 - W^+W^-
 - $\tau^+\tau^-$
 - $\mu^+\mu^-$

DM DECAY: SPECTRUM



All flavour neutrino flux at source from different DM decay/annihilation channels

FLUXES FROM HEAVY DM DECAY

Total Flux = Galactic Flux + Extra-Galactic Flux

- Galactic

$$- \frac{d\Phi^G}{dE_\nu} = \frac{1}{4\pi m_{\text{DM}} \tau_{\text{DM}}} \frac{dN_\nu}{dE_\nu} \int_0^\infty \rho(r(s, l, b)) ds$$


- Extragalactic

$$- \frac{d\Phi^{\text{EG}}}{dE} = \frac{\Omega_{\text{DM}} \rho_c}{4\pi m_{\text{DM}} \tau_{\text{DM}}} \int_0^\infty \frac{1}{H(z)} \frac{dN_\nu}{dE'_\nu} [(1+z)E'_\nu] dz$$

- Comparable contributions from G and EG fluxes, flux obtained from 4π sky


- High energy neutrinos attenuated by earth => more downgoing neutrinos than up-going

“HIDDEN” COMPONENT: ν FROM DM DECAY?




- Presence of Astrophysical power-law flux alongwith ν 's from heavy DM decay
- DM of mass $O(100)$ TeV could decay to SM particles
 - W^+W^- , Z^0Z^0 , $\tau^+\tau^-$, $\mu^+\mu^-$
 - Secondary neutrinos @ 10 TeV $\sim m_{DM}/2$
- DM decay augments lower energy event rates
- Allows for reduced power-law flux normalisation
 - Consistency with lack of events at 300 TeV – 1 PeV, and 1 PeV+ energies

“HIDDEN” COMPONENT: ν FROM DM DECAY?




- Presence of Astrophysical power-law flux alongwith ν 's from heavy DM decay
- DM of mass $O(100)$ TeV could decay to SM particles
 - W^+W^- , Z^0Z^0 , $\tau^+\tau^-$, $\mu^+\mu^-$
 - Secondary neutrinos @ 10 TeV $\sim m_{DM}/2$
- DM decay augments lower energy event rates
- Allows for reduced power-law flux normalisation
 - Consistency with lack of events at 300 TeV – 1 PeV, and 1 PeV+ energies

CONSTRAINTS ON DM PARAMETER SPACE




Astro flux at IC b.f.




Astro flux
normalisation as free
parameter

CONSTRAINTS ON DM PARAMETER SPACE



Astro flux at IC b.f.



Astro flux
normalisation as free
parameter

IC EVENTS AS DM + ASTRO

- **MORE DATA REQUIRED**
- Within the purview of limited statistics, reduced astrophysical flux + low energy neutrinos from DM decay fits observed data significantly better than the IC best-fit with a power-law astrophysical flux alone
 - Consistency with lack of events in the “well”
 - Better match to the sub-100 TeV events
- Event spectrum favours TeV scale DM, and is strongly constraining for PeV scale DM, esp. with astrophysical flux at the IC b.f.

IC EVENTS AS DM + ASTRO


- **MORE DATA REQUIRED**
- Within the purview of limited statistics, reduced astrophysical flux + low energy neutrinos from DM decay fits observed data significantly better than the IC best-fit with a power-law astrophysical flux alone
 - Consistency with lack of events in the “well”
 - Better match to the sub-100 TeV events
- Event spectrum favours TeV scale DM, and is strongly constraining for PeV scale DM, esp. with astrophysical flux at the IC b.f.

REMARKS

- Other efforts
 - Esmaili, Serpico JCAP 1311 (2013) 054
 - No astrophysical neutrinos, only DM decay
 - $DM \rightarrow \nu_e \nu_e$ (12%), $q\bar{q}$ (88%)
 - Also explains lack of events in the 300 TeV – 1 PeV bins
 - In some tension with lower energy events
- Updated background estimates for the neutrino fluxes from charm decay in the low TeV energies will help
 - Present calculations (Enberg et al, 2008) based on outdated pdf's

“HIDDEN” COMPONENT: ν FROM DM DECAY?

$\chi^2 = 4.209$
 $p\text{-value} = 0.061$



$\chi^2 = 4.209$
 $p\text{-value} = 0.061$

$\chi^2 = 4.188$
 $p\text{-value} = 0.060$

$\chi^2 = 4.445$
 $p\text{-value} = 0.072$

Compare with IC best-fit
 $\chi^2_{IC} = 10.7$

ADDENDUM: MORE RECENT EVENTS @ IC

- 9 more events since Dec. 2013, 37 in total
- One event at 370 TeV, none from 400 TeV – 1 PeV
- One more PeV+ event at ~ 2.1 PeV
 - Total of three PeV+ events
- E^{-2} best fit now at $0.95e-8 \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$

Talks by Halzen and Sullivan
<http://icecube.wisc.edu/meetings/neutrinos-beyond-icecube>
Apr. 24, Arlington

ADDENDUM: MORE RECENT EVENTS @ IC

- 9 more events since Dec. 2013, 37 in total
- One event at 370 TeV, none from 400 TeV – 1 PeV
- One more PeV+ event at ~ 2.1 PeV
 - Total of three PeV+ events
- E^{-2} best fit now at $0.95e-8 \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$

Talks by Halzen and Sullivan
<http://icecube.wisc.edu/meetings/neutrinos-beyond-icecube>
Apr. 24, Arlington

ADDITIONAL NEUTRINO FLUX COMPONENT?

Possible sources

- **NOT**
 - Atmospheric
 - Cosmogenic
- Heavy DM – $m_{\text{DM}} = O(100)$ TeV – annihilation?
 - Predominance of signal from galactic centre
 - Slight excess towards GC seen
 - For WIMP-like $\langle\sigma v\rangle^\dagger$, expected IC signal too low
 - 1 event in 50 yrs
- Heavy DM decay...

[†] For WIMP's, $\langle\sigma v\rangle \approx 10^{-27} \text{ cm}^3 \text{ s}^{-1}$