Title: 14/15 PSI - Relativity- Lecture 4

Date: Sep 11, 2014 09:00 AM

URL: http://pirsa.org/14090010

Abstract:


Correction to statement of Noether's theorem:
For every continuous symmetry of the action (up to surface terms)


Correction to statement of Noether's theorem:
$S' = S$
action (up to su
erms)
For every continuous symmetry of the
action principle,

Correction to statement of Noether's theorem.

For every continuous symmetry of the ~~equations of motion derived from an action principle,~~ action (up to surface terms), which holds without using the equations of motion,

// off-shell symmetries give on-shell conservation laws" (using the eq's of motion)

Example: free relativistic particle

$$S = -m_0 c^2 \int dt \sqrt{1 - \left(\frac{dx}{dt}\right)^2 \frac{1}{c^2}} \approx$$




continuous symmetry of the ~~equations of motion, derived from an action principle,~~
"off-shell symmetries give on-shell conservation laws" using the eq's of motion
Example: free relativistic particle
$S = -m_0 c^2 \int dt \sqrt{1 - \left(\frac{dx}{dt}\right)^2 \frac{1}{c^2}} \approx -m_0 c^2 \int dt + \int dt \, \frac{1}{2} m_0 \left(\frac{dx}{dt}\right)^2 + \cdots$
⇒
consider symmetry under spatial translations $\underline{x}(t) \to \underline{x}'(t) = \underline{x}(t) + \underline{\epsilon}$


symmetry of the equations of motion, derived from an action principle,
"off-shell symmetries give on-shell conservation laws"
using the eq's of motion
Example: free relativistic particle
$-m_0c^2\int dt\sqrt{1-\left(\frac{dx}{dt}\right)^2\frac{1}{c^2}} \approx -m_0c^2\int dt + \int dt\,\frac{1}{2}m_0\left(\frac{dx}{dt}\right)^2 + \cdots$
consider symmetry under spatial translations $\underline{x}(t) \to \underline{x}'(t) = \underline{x}(t) + \underline{\epsilon}$, a symmetry if $\underline{\epsilon}$ is constant (indept of t).
$\delta\left(\frac{dx}{dt}\right) = \frac{dx'}{dt} - \frac{dx}{dt} = \frac{d\epsilon}{dt}$


symmetry of the equations of motion, derived from an action principle,
"off-shell symmetries give on-shell conservation laws"
using the eq's of motion
Example: free relativistic particle
$S = -m_0c^2\int dt\sqrt{1-\quad} \approx -m_0c^2\int dt + \int dt\,\frac{1}{2}m_0\left(\frac{dx}{dt}\right)^2 + \cdots$
⇒ conserved quantities: consider symmetry unde translations $\underline{x}(t) \to \underline{x}'(t) = \underline{x}(t) + \underline{\epsilon}$, a symmetry if $\underline{\epsilon}$ is constant (indept of t).
Noether: let ϵ be a function of t
$\frac{dx}{dt} = \frac{dx}{dt} = \frac{d\epsilon}{dt}$
$\delta S = +m_0c^2\int dt\ \frac{\frac{dx}{dt}\frac{d\delta x}{dt}\frac{1}{c^2}}{\sqrt{1-\left(\frac{ds}{dt}\right)^2\frac{1}{c^2}}}$

$$\delta S = +m_0 c^2 \int dt \, \frac{\frac{dx}{dt}\frac{d\delta x}{dt}\frac{1}{c^2}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2\frac{1}{c^2}}} = m_0 \int dt \, \frac{\frac{dx}{dt}\frac{d\epsilon}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2\frac{1}{c^2}}}$$

by Noether argument, $\delta S = 0$ on shell (i.e. using the eq^{n}s of motion) for arbitrary $\epsilon(t)$

$$= - m_0 \int dt \, \frac{d}{dt}\left(\frac{\frac{dx}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2\frac{1}{c^2}}}\right)\epsilon(t) \qquad \int dt \left(\frac{\frac{dx}{dt}}{\sqrt{\quad}}\right) = 0$$

ted quantities: consider symmetry under spatial translations $\underline{x}(t) \to \underline{x}'(t) = \underline{x}(t) + \underline{\epsilon}$, a symmetry if $\underline{\epsilon}$ is constant (indept of t).

Noether: let ϵ be a function of t $\quad \delta\left(\frac{dx}{dt}\right) = \frac{dx'}{dt} - \frac{dx}{dt} = \frac{d\epsilon}{dt}$

$$\delta S = +m_0 c^2 \int dt \, \frac{\frac{dx}{dt}\frac{d\delta x}{dt}\frac{1}{c^2}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2\frac{1}{c^2}}} = m_0 \int dt \, \frac{\frac{dx}{dt}\frac{d\epsilon}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2\frac{1}{c^2}}}$$

by Noether argument, $\delta S = 0$ on shell (i.e. using the eq^{n}s of motion) for arbitrary $\underline{\epsilon}(t)$

$$= - m_0 \int dt \, \frac{d}{dt}\left(\frac{\frac{dx}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2\frac{1}{c^2}}}\right)\epsilon(t) \quad \Rightarrow \frac{d}{dt}\left(\frac{m_0 \frac{dx}{dt}}{\sqrt{\quad}}\right) = 0 \quad \Rightarrow \underline{p} = \frac{m_0 \frac{dx}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2\frac{1}{c^2}}}$$ is conserved

⇒ Conserved quantities: Consider symmetry under spatial translations $\underline{x}(t) \to \underline{x}'(t) = \underline{x}(t) + \underline{\epsilon}$, a symmetry if $\underline{\epsilon}$ is const

Noether: let ϵ be a function of t : $\delta\left(\frac{d\underline{x}}{dt}\right) = \frac{d\underline{x}'}{dt} - \frac{d\underline{x}}{dt} = \frac{d\underline{\epsilon}}{dt}$

$$\delta S' = -m_0 c^2 \int dt \frac{\frac{d\underline{x}}{dt}\cdot\frac{d\underline{\epsilon}}{dt}\frac{1}{c^2}}{\sqrt{1-\left(\frac{d\underline{x}}{dt}\right)^2\frac{1}{c^2}}} = m_0 \int dt \frac{\frac{d\underline{x}}{dt}\cdot\frac{d\underline{\epsilon}}{dt}}{\sqrt{1-\left(\frac{d\underline{x}}{dt}\right)^2\frac{1}{c^2}}}$$

by Noether argument, $\delta S' = 0$ on shell (i.e. using the eq^{n}s of motion) for arbitrary $\underline{\epsilon}(t)$

$$= - m_0 \int dt \frac{d}{dt}\left(\frac{\frac{d\underline{x}}{dt}}{\sqrt{1-\left(\frac{d\underline{x}}{dt}\right)^2\frac{1}{c^2}}}\right)\epsilon(t) \Rightarrow \frac{d}{dt}\left(\frac{m_0\frac{d\underline{x}}{dt}}{\sqrt{\quad}}\right) = 0 \Rightarrow \underline{p} = \frac{m_0\frac{d\underline{x}}{dt}}{\sqrt{1-(}}$$

Now consider symmetry under time translations: $\underline{x}(t) \to \underline{x}'(t) = \underline{x}(t) + \epsilon\frac{d\underline{x}}{dt}$: $\delta\underline{x}(t) = \epsilon\frac{d\underline{x}}{dt}$: $\delta\frac{d\underline{x}}{dt} =$

$$\delta S' = m_0 c^2 \int \frac{dt \frac{1}{c^2}\frac{d\underline{x}}{dt}\frac{d}{dt}\overbrace{\left(\epsilon\frac{d\underline{x}}{dt}\right)}^{\delta\underline{x}}}{\sqrt{1-\left(\frac{d\underline{x}}{dt}\right)^2\frac{1}{c^2}}}$$


Action principle for a particle, in a certain sense, gives e nagnetism and gravity for free!
$S = \int(-mc\sqrt{-\eta_{\mu\nu}dx^\mu dx^\nu} + q dx^\mu A$
$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$


Action principle for a particle, in a cent nse, gives electromagnetism and gravity for free!
$x^\mu(\lambda)$
$x^\mu \to x^\mu + \delta x^\mu$
$S = \int (-mc\sqrt{-} + q dx^\mu A_\mu(x))$
$x^\mu(\lambda)$
$\delta S = \int \frac{mc dx^\mu}{\sqrt{}} A_\mu + q\, dx^\mu A_{\mu,\nu}$
$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$
A_μ is the electromagnetic vector potential


integrate
by parts


$+ q\frac{d\delta x^\mu}{d\tau} A_\mu + q\frac{dx^\mu}{d\tau} A_{\mu,\nu}\delta x^\nu \Big) d\tau$
$mc\frac{dx_\mu}{d\tau}\frac{d\delta x^\mu}{d\tau}$
$\Delta M = \frac{E}{c^2}$
$\delta x^\mu \, mc\frac{d^2x_\mu}{d\tau^2} - q\delta x^\mu \partial_\nu A_\mu \frac{dx^\nu}{d\tau} + q\delta x^\mu A_{\nu,\mu}\frac{dx^\nu}{d\tau}\Big) = 0 \quad \forall \delta x^\mu(\tau)$


integrate by parts
x(τ)


A_μ is the electromagnetic vector potential
$(M-m) = 2\frac{E_r}{c^2}$
$\Delta M = \frac{E_{emitted}}{c^2}$
$\forall \delta x^\mu(\tau)$
QM
Lorentz force.
phase $S/\hbar$


Action principle for a particle, in a certain sense, gives electromagnetism and gravity for free!
$S = \int \left(-mc\sqrt{-\eta_{\mu\nu}dx^\mu dx^\nu} + q dx^\mu A_\mu \right)$
$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$
A_μ is the electromagnetic vector potential
$\delta S = \int \left(\frac{mc\, dx^\mu d\delta x^\mu}{d\tau} + q\frac{d\delta x^\mu}{d\tau} A_\mu + q\frac{dx^\mu}{d\tau} A_{\mu,\nu}\delta x^\nu \right) d\tau$
$\frac{dx^\mu}{d\tau} d\delta x^\nu \eta_{\mu\nu} = \frac{dx^\mu}{d\tau} d\delta x_\mu$
$d\tau$ $mc\frac{dx_\mu}{d\tau}\frac{d\delta x^\mu}{d\tau}$
integrate by parts
$= \int \left(-\delta x_\mu mc\frac{d^2x^\mu}{d\tau^2} - q\,\delta x^\mu\, \partial_\nu A_\mu \frac{dx^\nu}{d\tau} + q\,\delta x^\mu\, A_{\nu,\mu}\frac{dx^\nu}{d\tau} \right) = 0 \quad \forall \delta x^\mu(\tau)$
$\frac{d}{d\tau}\left(A_\mu(x(\tau))\right) = \frac{\partial A_\mu}{\partial x^\nu}$
$\Rightarrow mc\frac{d^2x^\mu}{d\tau^2} = q\left(\partial_\mu A_\nu - \partial_\nu A_\mu\right)\frac{dx^\nu}{d\tau} = q F_{\mu\nu}\frac{dx^\nu}{d\tau}$
Lorentz force
QM
$\int \sim \int e^{\frac{i}{\hbar}S'}$
phase $\frac{S}{\hbar} = \frac{q}{\hbar}\int A_\mu dx^\mu$
Aharonov-Bohm
$E \cdot |p| = E_{rest}$
$(M - m) = \frac{2E_x}{c^2}$
$\Delta M = \frac{E_{emitted}}{c^2}$


QM
$= \frac{\partial}{\partial x^\mu} A_\nu \cdot \frac{dx^\nu}{d\tau}$
$\Rightarrow m \frac{d^2 x^\mu}{d\tau^2} = q(\partial_\mu A_\nu - \partial_\nu A_\mu)\frac{dx^\nu}{d\tau} = q F_{\mu\nu}\frac{dx^\nu}{d\tau}$
Lorentz force.
$\Psi \sim \int e^{\frac{i}{\hbar}S'}$
phase $\frac{S'}{\hbar} = \frac{q}{\hbar}\int A_\mu dx^\mu$
In fact, action for electromagnetic fields is
$\int d^4x \left(-\frac{1}{4\mu_0} F_{\mu\nu}F^{\mu\nu} + J_\mu A^\mu\right)$
$\propto \epsilon_{\mu\nu\rho\sigma} dx^\mu dx^\nu dx^\rho dx^\sigma$
particles
: Can check that, for $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$,
vary S' wrt. A_μ $\Rightarrow \partial_\mu F^{\mu\nu} = -\mu_0 J^\nu$
(Aharonov-Bohm.
for a particle $J_\mu \propto q\frac{dx_\mu}{d\tau}$
charged
another symmetry we have used is the symmetry under
translations in space: $x^\mu \to x'^\mu = x^\mu + a^\mu$
$\int -mc\sqrt{-\eta_{\mu\nu}dx^\mu dx^\nu}$
$\downarrow$
$g_{\mu\nu}(x)$
$\longleftrightarrow g_{\mu\nu}(x)$
Lorentz symmetry generalises to $x^\mu \to x'^\mu(x)$
any coordinate transformation
LIs
$\Lambda^\alpha_\gamma \Lambda^\beta_\delta$
x^γ
$\Rightarrow$ Coordinate transformation matrix $\frac{\partial x'^\alpha}{\partial x^\beta} = \Lambda^\alpha_\beta$

$\frac{dx^\nu}{d\tau} \Rightarrow m\frac{d^2x^\mu}{d\tau^2} = q(\partial_\mu A_\nu - \partial_\nu A_\mu)\frac{dx^\nu}{d\tau} = qF_{\mu\nu}\frac{dx^\nu}{d\tau}$ Lorentz force. $\Psi \sim e^{\frac{i}{\hbar}S}$

phase $\frac{S}{\hbar} = \frac{q}{\hbar}\int A_\mu dx^\mu$

, action for electromagnetic fields is $\int d^4x \left(-\frac{1}{4\mu_0}F_{\mu\nu}F^{\mu\nu} + J_\mu A^\mu\right)$: Can check that, for $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, vary S w.r.t. A_μ $\Rightarrow \partial_\mu F^{\mu\nu} = -\mu_0 J^\nu$ (Aharonov-Bohm.

$\propto \epsilon_{\mu\nu\rho\sigma}dx^\mu dx^\nu dx^\rho dx^\sigma$

for a charged particle $J_\mu \propto q\frac{dx_\mu}{d\tau}$

J_μ: particles

$\int -mc\sqrt{-\eta_{\mu\nu}dx^\mu dx^\nu}$

$\downarrow$

$g_{\mu\nu}(x)$

another symmetry we have used is the symmetry under translations in spacetime, $x^\mu \to x'^\mu = x^\mu + a^\mu$

Together, Lorentz transformations and spacetime translations form the Poincaré group.

(x) Lorentz symmetry generalises to $x^\mu \to x'^\mu(x)$ any coordinate transformation

coordinate transformation matrix $\frac{\partial x'^\mu}{\partial x^\beta} = \Lambda^\mu{}_\beta$

$\frac{dx^\nu}{d\tau} \Rightarrow m\frac{d^2x^\mu}{d\tau^2} = q(\partial_\mu A_\nu - \partial_\nu A_\mu)\frac{dx^\nu}{d\tau} = qF_{\mu\nu}\frac{dx^\nu}{d\tau}$ Lorentz force. $\Psi \sim e^{\frac{i}{\hbar}S}$

phase $\frac{S}{\hbar} = \frac{q}{\hbar}\int A_\mu dx^\mu$

, action for electromagnetic fields is $\int d^4x \left(-\frac{1}{4\mu_0}F_{\mu\nu}F^{\mu\nu} + J_\mu A^\mu\right)$: Can check that, for $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, vary S w.r.t. A_μ $\Rightarrow \partial_\mu F^{\mu\nu} = -\mu_0 J^\nu$ (Aharonov-Bohm.

$\propto \epsilon_{\mu\nu\rho\sigma}dx^\mu dx^\nu dx^\rho dx^\sigma$

for a particle (charged) $J_\mu \propto q\frac{dx_\mu}{d\tau}$

particles

$\int -mc\sqrt{-\eta_{\mu\nu}dx^\mu dx^\nu}$

$\downarrow$

$g_{\mu\nu}(x)$

(x) Lorentz symmetry generalises to $x^\mu \to x'^\mu(x)$ any coordinate transformation

coordinate transformation matrix $\frac{\partial x'^\alpha}{\partial x^\beta} = \Lambda^\alpha{}_\beta$

another symmetry we have used is the symmetry under transl... n spacetime, $x^\mu \to x'^\mu = x^\mu + a^\mu$

Toget... z transformations and spacetime translations form the Po... group (a 10-parameter group).

In fact, ... the most general continuous transformations $x^\alpha \to x'^\alpha$ ... preserve $\eta_{\mu\nu}$, ie. for which

... $= \eta_{\gamma\beta}\frac{\partial x'^\gamma}{\partial x^\delta}\frac{\partial x'^\delta}{\partial x^\alpha}$


Noether: Let ϵ be a function of t
δ(dx/dt) = dx'/dt − dx/dt = dϵ/dt
δS' = +m₀c² ∫dt (dx/dt · d(δx)/dt) / √(1 − (dx/dt)²/c²) = m₀ ∫dt (dx/dt · dϵ/dt) / √(1 − (dx/dt)²/c²)
Proof: ∂/∂xᵉ
0 = η_αβ ∂²(x'^β)/∂x^δ


Noether: Let ϵ be a function of t
$\delta\left(\frac{dx}{dt}\right) = \frac{dx'}{dt} - \frac{dx}{dt} = \frac{d\epsilon}{dt}$
$\delta S = +m_0 c^2 \int dt \frac{\frac{dx}{dt}\frac{d\delta x}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2}} = m_0 \int dt \frac{\frac{dx}{dt}\frac{d\epsilon}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2}}$
Proof: $\frac{\partial}{\partial x^\epsilon}$
$0 = \eta_{\alpha\beta}$
$+ \eta$


Noether: Let ϵ be a function of t
$\delta\left(\frac{dx}{dt}\right) = \frac{dx'}{dt} - \frac{dx}{dt} = \frac{d\epsilon}{dt}$
$\delta S = +m_0 c^2 \int dt \frac{\frac{dx}{dt}\frac{d\delta x}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2}} = m_0 \int dt \frac{\frac{dx}{dt}\frac{d\epsilon}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2}}$
Proof: $\frac{\partial}{\partial x^\epsilon}$
$0 = \eta_{\alpha\beta}\frac{\partial^2 x'^\alpha}{\partial x^\gamma \partial x^\epsilon}\frac{\partial x'^\beta}{\partial x^\delta} + \eta_{\alpha\beta}\frac{\partial x'^\alpha}{\partial x^\gamma}\frac{\partial^2 x'^\beta}{\partial x^\delta \partial x^\epsilon}$
$+(\gamma \leftrightarrow \epsilon)$
$+ \frac{x'}{\partial x^\delta \partial x^\epsilon}\frac{\partial x'^\beta}{\partial x^\delta} + \eta_{\alpha\beta}\frac{\partial x'^\alpha}{\partial x^\epsilon}\frac{\partial^2 x'^\beta}{\partial x^\delta \partial x^\gamma}$
$-(\epsilon \leftrightarrow \delta)$

Noether: Let ϵ be a function of t : $\delta\left(\frac{dx}{dt}\right) = \frac{dx'}{dt} - \frac{dx}{dt} = \frac{d\epsilon}{dt}$

$$\delta S' = +m_0 c^2 \int dt \frac{\frac{dx}{dt}\frac{d\delta x}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2}} = m_0 \int dt \frac{\frac{dx}{dt}\frac{d\epsilon}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2}}$$

Proof: $\frac{\partial}{\partial x^\epsilon}$

$$0 = \eta_{\alpha\beta}\frac{\partial^2 x'^\alpha}{\partial x^\delta \partial x^\epsilon}\frac{\partial x'^\beta}{\partial x^\delta} + \eta_{\alpha\beta}\frac{\partial x'^\alpha}{\partial x^\delta}\frac{\partial^2 x'^\beta}{\partial x^\delta \partial x^\epsilon}$$

$$+(\delta \leftrightarrow \epsilon) \quad +\eta_{\alpha\beta}\frac{\partial^2 x'}{\partial x^\delta \partial x^\epsilon}\frac{\partial x'^\beta}{\partial x^\delta} + \eta_{\alpha\beta}\frac{\partial x'^\alpha}{\partial x^\epsilon}\frac{\partial^2 x'^\beta}{\partial x^\delta \partial x^\delta}$$

$$-(\epsilon \leftrightarrow \delta) \quad -\eta_{\alpha\beta}\frac{\partial^2 x'^\alpha}{\partial x^\delta \partial x^\delta}\frac{\partial x'^\beta}{\partial x^\epsilon} - \eta_{\alpha\beta}\frac{\partial x'^\alpha}{\partial x^\delta}\frac{\partial^2 x'^\beta}{\partial x^\delta \partial x^\epsilon}$$



Noether: Let ϵ be a function of t : $\delta\left(\frac{dx}{dt}\right) = \frac{dx'}{dt} - \frac{dx}{dt} = \frac{d\epsilon}{dt}$

$$\delta S' = +m_0 c^2 \int dt \, \frac{\frac{dx}{dt}\frac{d\delta x}{dt}}{\sqrt{1-\left(\frac{dx}{dt}\right)^2}} = m_0 \int dt \, \frac{\frac{dx}{dt}\frac{d\epsilon}{dt}}{\sqrt{1-(dx)^2}}$$

Proof: $\frac{\partial}{\partial x^\epsilon}$

$$0 = \eta_{\alpha\beta}\frac{\partial^2 x'^\alpha}{\partial x^\gamma \partial x^\epsilon}\frac{\partial x'^\beta}{\partial x^\delta} + \eta_{\alpha\beta}\frac{\partial x'^\alpha}{\partial x^\gamma}\frac{\partial^2 x'^\beta}{\partial x^\delta \partial x^\epsilon}$$

$$+(\gamma \leftrightarrow \epsilon) \quad +\eta_{\alpha\beta}\frac{\partial^2 x'}{\partial x^\delta \partial x^\epsilon}\frac{\partial x'^\beta}{\partial x^\delta} + \eta_{\alpha\beta}\frac{\partial x'^\alpha}{\partial x^\epsilon}\frac{\partial^2 x'^\beta}{\partial x^\delta \partial x^\gamma}$$

$$-(\epsilon \leftrightarrow \delta) \quad -\eta_{\alpha\beta}\frac{\partial^2 x'^\alpha}{\partial x^\gamma \partial x^\delta}\frac{\partial x'^\beta}{\partial x^\epsilon} - \eta_{\alpha\beta}\frac{\partial x'^\alpha}{\partial x^\delta}\frac{\partial^2 x'^\beta}{\partial x^\gamma \partial x^\epsilon}$$

$\Rightarrow \eta_{\alpha\beta}$


function of t
Invertible
invertible