

Title: Growth of supermassive black holes and their relationships to their host galaxies

Date: Nov 10, 2014 09:30 AM

URL: <http://pirsa.org/14110062>

Abstract: Black holes are the engines that power quasars and active galactic nuclei throughout cosmic time. The masses of black holes in nearby galaxies define clear correlations with the properties of their host galaxies. These results suggest that black holes, while a thousand times lighter than the galaxy, grow alongside their hosts during its cosmic evolution. I will discuss the growth of black holes, and the establishment of the connection between galaxies and black holes.

Massive Black Holes and their host galaxies


Marta Volonteri

Institut d'Astrophysique de Paris

BLACK 
ERC Research Grant

MBHG 

Quasars and AGN




Point-like, or star-like, radio sources which varied rapidly: 'quasi-stellar' radio sources or quasars.

The quasar 3C273 is 640 Mpc ~ 2.6 billion light years away.

The luminosity of 3C273 is more than 100 times the luminosity of our entire galaxy.

Quiescent MBHs


Many MBHs are quiescent.
We have an example is in the
center of the Milky Way.



Quiescent MBHs

Many MBHs are quiescent.
We have an example is in the
center of the Milky Way.


The typical luminosity
coming from the very
center, Sgr A*, is $\sim 10^{34}$ erg/s.




Massive black holes out there....



Hubble Deep Field
visible light – starlight
all galaxies
“black” black holes?



Chandra Deep Field
X-ray – high energy processes
quasars and active
galactic nuclei (AGN)




Galaxies

mass: 10^9 - 10^{12} solar masses

$$R_{\text{halo}} \sim GM_{\text{halo}} / \sigma^2 \quad \text{MEGAPARSEC}$$

$$R_{\text{bulge}} \sim GM_{\text{bulge}} / \sigma^2 \quad \text{KILOPARSEC}$$

1 parsec = 3.26 light years = 3×10^{18} cm
 $\sigma \sim 50$ - 300 km/s for most galaxies



Galaxies

mass: 10^9 - 10^{12} solar masses

$$R_{\text{halo}} \sim GM_{\text{halo}}/\sigma^2 \quad \text{MEGAPARSEC}$$

$$R_{\text{bulge}} \sim GM_{\text{bulge}}/\sigma^2 \quad \text{KILOPARSEC}$$

1 parsec = 3.26 light years = 3×10^{18} cm
 $\sigma \sim 50$ - 300 km/s for most galaxies

Massive Black Holes


mass: 10^5 - 10^9 solar masses

$$R_{\text{bondi}} \sim GM_{\text{BH}}/c_s^2 \quad \text{PARSEC}$$

$$R_{\text{inf}} \sim GM_{\text{BH}}/\sigma^2 \quad \text{PARSEC}$$

$$R_{\text{sch}} = 2GM_{\text{BH}}/c^2 \quad \text{MICROPARSEC}$$

$c_s \sim 10$ - 100 km/s for most galaxies
 $c = 3 \times 10^5$ km/s



Galaxies

mass: 10^9 - 10^{12} solar masses

$$R_{\text{halo}} \sim GM_{\text{halo}}/\sigma^2 \quad \text{MEGAPARSEC}$$

$$R_{\text{bulge}} \sim GM_{\text{bulge}}/\sigma^2 \quad \text{KILOPARSEC}$$

1 parsec = 3.26 light years = 3×10^{18} cm
 $\sigma \sim 50$ - 300 km/s for most galaxies

Massive Black Holes

mass: 10^5 - 10^9 solar masses

$$R_{\text{bondi}} \sim GM_{\text{BH}}/c_s^2 \quad \text{PARSEC}$$


$$R_{\text{inf}} \sim GM_{\text{BH}}/\sigma^2 \quad \text{PARSEC}$$

$$R_{\text{sch}} = 2GM_{\text{BH}}/c^2 \quad \text{MICROPARSEC}$$

$c_s \sim 10$ - 100 km/s for most galaxies
 $c = 3 \times 10^5$ km/s

MBH- host relations

Black hole masses correlate with galaxy properties. This may mean their growth/evolution are intimately connected.




- $M_{\text{BH}} \sim 10^{-3} M_{\text{bulge}}$

MV using data from Kormendy & Ho 2013

MBH- host relations

Black hole masses correlate with galaxy properties. This may mean their growth/evolution are intimately connected.



- $M_{\text{BH}} \sim 10^{-3} M_{\text{bulge}}$

- $M_{\text{BH}} \sim \sigma^4 - \sigma^5$

σ : velocity dispersion

- Through Faber-Jackson: $L \sim \sigma^4$
- IR light traces mass

MV using data from Kormendy & Ho 2013

M_{BH} - host relations: **how** are they established?

Is the correlation regulated by the galaxy or by the MBH?

Feeding: the galaxy sets the MBH mass by regulating the amount of gas that trickles to the MBH

M_{BH} - host relations: **how** are they established?

Is the correlation regulated by the galaxy or by the MBH?

Feeding: the galaxy sets the MBH mass by regulating the amount of gas that trickles to the MBH

Feedback: The MBH regulates the process: when it reaches a limiting mass and luminosity it drives outflows that sweep away the surrounding gas, thus halting both its own growth and star formation in the galaxy.

M_{BH} - host relations: **how** are they established?


Is the correlation regulated by the galaxy or by the MBH?

Feeding: the galaxy sets the MBH mass by regulating the amount of gas that trickles to the MBH

Feedback: The MBH regulates the process: when it reaches a limiting mass and luminosity it drives outflows that sweep away the surrounding gas, thus halting both its own growth and star formation in the galaxy.


Casuality: a large number of mergers will average out the extreme values of $M_{\text{BH}}/M_{\text{bulge}}$ towards the ensemble average

Feeding



Courtesy of
P. Capelo

Galaxy mergers trigger nuclear inflows of low angular
momentum gas



Feeding

Streams of gas plunge into halos and feed the galaxy with large amounts of fresh material

Bellovary, MV et al. 2013

Feedback

The mass of the MBH is $\sim M_{\text{BH}} \sim 10^{-3} M_{\text{gal}}$.

The energy released by the MBH growth is $E_{\text{BH}} = 0.1 M_{\text{BH}} c^2$

Fabian 2012

Feedback

The mass of the MBH is $\sim M_{\text{BH}} \sim 10^{-3} M_{\text{gal}}$.

The energy released by the MBH growth is $E_{\text{BH}} = 0.1 M_{\text{BH}} c^2$

The binding energy of the galaxy is $E_{\text{gal}} \sim M_{\text{gal}} \sigma^2$, with $\sigma < 400 \text{ km s}^{-1}$

$$\Rightarrow E_{\text{BH}}/E_{\text{gal}} \sim 10^{-4} (c/\sigma)^2 > 80$$

Feedback

The mass of the MBH is $\sim M_{\text{BH}} \sim 10^{-3} M_{\text{gal}}$.

The energy released by the MBH growth is $E_{\text{BH}} = 0.1 M_{\text{BH}} c^2$


The binding energy of the galaxy is $E_{\text{gal}} \sim M_{\text{gal}} \sigma^2$, with $\sigma < 400 \text{ km s}^{-1}$

$$\Rightarrow E_{\text{BH}}/E_{\text{gal}} \sim 10^{-4} (c/\sigma)^2 > 80$$

If even a small fraction of the energy can be transferred to the gas, then an AGN can have a profound effect on the evolution of its host galaxy

Fabian 2012


The growth of MBHs in galaxies




How do MBHs grow to become supermassive?

How do MBH seeds grow to become supermassive?

MBH-MBH mergers vs gas accretion




Mergers: Total mass density in MBHs is constant in time: just reshuffle the mass function



Accretion: Total mass density in MBHs grows with time

How do MBH seeds grow to become supermassive?




Mergers: total mass density in MBHs is constant in time: just reshuffle the distribution of masses

Accretion: adds external matter => total mass density in MBHs grows with time

Soltan's argument:
mass density increases by > one order of magnitude in the last ~10 Gyr: **accretion** leads

How do MBH seeds grow to become supermassive?




Mergers: total mass density in MBHs is constant in time: just reshuffle the distribution of masses

Accretion: adds external matter => total mass density in MBHs grows with time

Soltan's argument:
mass density increases by > one order of magnitude in the last ~10 Gyr: **accretion** leads

Do MBH-MBH mergers ever count?




Fraction of mass gained through MBH-MBH mergers:

$$f_{\text{merge}} = \Delta M_{\text{merge}} / M_{\text{BH}}$$

ΔM_{merge} is the sum of the masses of all merged MBHs and does not account for gas accretion on these MBHs

Dubois, Volonteri & Silk 2013

Do MBH-MBH mergers ever count?



High-mass
MBHs!

SgrA* and M87
had very different
growth histories

Fraction of mass gained through MBH-MBH mergers:


$$f_{\text{merge}} = \Delta M_{\text{merge}} / M_{\text{BH}}$$

ΔM_{merge} is the sum of the masses of all merged MBHs and does not account for gas accretion on these MBHs


Dubois, Volonteri & Silk 2013

What is the typical MBH spin?


✓ mergers can spin MBHs either up or down, but typical mergers yield spin ~ 0.7



✓ angular momentum is transferred from disk to hole, spin up or down depends on disk/hole mass ratio and alignment




Spins from accretion



Coherent: material has
~ constant direction of
angular momentum
vector \Rightarrow high spin


Chaotic: droplets of
material with random
direction of the angular
momentum vector \Rightarrow
low spin

Spins from accretion: local environment



Dubois, MV, Silk et al. 2014

Spins from accretion: local environment



Cold, dense gas tends to reside in a rotationally supported disc => Eddington-limited accretion episodes are synchronous with **coherent** accretion ($J_{\text{acc}} \sim \text{const}$) that dominates the mass budget

In phases characterized by low accretion rate the gas is hot and diffuse => **chaotic** accretion (J_{acc} fluctuates) dominates, but neither mass nor spin change much

Dubois, MV, Silk et al. 2014

Sgr A* & M87 in context

Mass: they're somewhat at the two extremes of the mass range known today

Accretion rate: both very sub-Eddington => must have grown in the past

M87: have MBH-MBH mergers played a role?

Milky Way: when has the MBH grown last?

Sgr A* & M87 in context

Spin: conjectures

M87: if MBH-MBH mergers, expect spin ~ 0.7
(statistically!) cf Doeleman+2012: $a > 0.2$

SgrA*: if most of the growth driven by rapid accretion: initially high-spin, but spinning down because of slow accretion of hot gas
cf Broderick, $a \sim 0$ (< 0.7); Moscibrodzka, Drappeau, $a > 0.5$
Orientation? Psaltis+2014