


Title: Detecting gravitational waves from supermassive binary black holes with pulsar timing

Date: Nov 12, 2014 09:50 AM


URL: <http://pirsa.org/14110089>

Abstract: I will discuss the current status of the NANOGrav pulsar timing array, and the prospects for a detection of the stochastic background produced by the mergers of supermassive black holes.

The North American Nanohertz Observatory for Gravitational Waves: about 50 students and scientists in the US and Canada working to characterize the gravitational wave universe at low frequencies using pulsar timing. Part of a world-wide effort including European and Australian partners.




Gravitational waves (GWs)



Gravitational waves are ripples in spacetime that propagate at the speed of light. They are a key prediction of general relativity.


They are produced by massive objects moving rapidly, and they change the distance between freely falling objects.

Their strength is measured by “strain”.




Gravitational wave observations will provide an entirely new means to study the universe.

Evidence for GWs




- Hulse and Taylor observed a rotating neutron star in a binary for 25 years

- Found orbit decay consistent with gravitational wave emission (Nobel prize 1993)




Now we're trying to detect GWs directly and build a telescope

Gravitational wave physics experiments



credit: NANOGrav Collaboration

Analogy with electromagnetic observations



Observations in different parts of the EM spectrum have provided us with invaluable complementary insights and knowledge about our universe.


When a new part of phase space is revealed there are always new discoveries.

Our goal is to inaugurate the era of low-frequency GWs.




Gravitational wave sources

The most promising sources are supermassive binary black holes (SMBBHs):



Other sources at nanohertz frequencies include cosmic strings, inflation, and phase transitions in the early universe.





Pulsars

Pulsars are a type of neutron star which have strong magnetic fields, spin rapidly, and emit beams of radio waves along their magnetic axes.

Masses of about a solar mass, radii of about 10km.

Nuclear density—aside from black holes the most extreme objects in the universe.



Rotate with periods of milliseconds to seconds



Celestial lighthouses/clocks

Pulsars

Pulsars are a type of neutron star which have strong magnetic fields, spin rapidly, and emit beams of radio waves along their magnetic axes.

Masses of about a solar mass, radii of about 10km.

Nuclear density—aside from black holes the most extreme objects in the universe.



Rotate with periods of milliseconds to seconds

Celestial lighthouses/clocks

Pulsar timing


Millisecond pulsars (MSPs) have periods < 20 ms and rotate rapidly with a long term stability that rivals atomic clocks. Millisecond pulsars are Nature's celestial clocks:

E.g.: Rotational period of PSR J1713+0747 on Tuesday Oct 28 2014 10:00 EDT:


$0.0045701365287363 \pm 0.0000000000000001$ s

at midnight Nov 30 2019 it will be $0.0045701365300853 \pm 0.0000000000000001$ s

GWs perturb the times of arrival of pulses at levels that will be measurable.




A galactic-scale GW detector: the Pulsar Timing Array



GW perturbations are correlated among different pulsars.

Need to observe an ensemble of MSPs to extract the correlated signal from the noise.



The Green Bank Telescope and Arecibo Observatory

Our measurements are made with the two most sensitive radio telescopes in the world:




Arecibo Observatory (AO), PR
World's largest
radio telescope



Green Bank Telescope (GBT), WV
World's largest steerable
radio telescope

Soon CHIME will contribute to our data sets at low frequencies

NANOGrav Observing Status




We observe 42 pulsars at the GBT and Arecibo, roughly every three weeks, at two radio frequencies.

We search for GWs in “timing residuals”, calculated over long time spans.

NANOGrav Activities

GW detector construction and characterization

- Find additional MSPs (200 with at least 20-40 suitable for PTAs) to increase our sensitivity
- More efficient/sensitive pulsar searches
- Fully characterized low-frequency GW detector




GW data set generation and curation

- Regular (18 month) open data releases
- New pulsar timing packages
- Cyber-I data curation system




GW detection and characterization

- First detection of low-frequency GWs or tightest constraints to date
- Comprehensive open-source GW data analysis suite



Our goals


Making a detection of the stochastic background produced by SMBBHs, inaugurating the era of low-frequency GWs.



credit: NANOGrav Collaboration

Our goals


Making a detection of the stochastic background produced by SMBBHs, inaugurating the era of low-frequency GWs.



credit: NANOGrav Collaboration

Our goals

Making a detection of the stochastic background produced by SMBBHs, inaugurating the era of low-frequency GWs.



**We are already in
astrophysically
interesting territory!**

credit: NANOGrav Collaboration


Our goals

As the low-frequency GW sky comes into focus, it will offer a **novel view** of **unique and groundbreaking physics**.

Individual supermassive black hole inspirals and their collective “chorus”: physics of accretion, late inspiral dynamics



Cosmic strings: early universe physics/high energy physics



New physics: expect to be surprised



Black hole merger “memory”: a surprising prediction of strong field general relativity.



Alternative theories of gravity and GWs

Two potential differences:

- 1) Additional GW polarization states: up to 6 modes

In GR only have 2 polarization states. Metric perturbation $h_{\mu\nu}$ has 10 independent components. Can perform a coordinate transformation $x_\mu \rightarrow \tilde{x}_\mu(x_\mu)$ that removes 4 components. In GR can work in Lorentz gauge, $\partial_\mu \bar{h}^{\mu\nu} = 0$ removing 4 additional components. In general cannot impose Lorentz gauge.

- 2) GWs may propagate with speeds $< c$. E.g. in massive gravity GWs will have a dispersion relation.

Additional polarization states






Figure
from
Nishizawa
et al. 2009

Antenna pattern response functions for pulsar-Earth system



Effect on stochastic backgrounds


Change the expected shape of the Hellings-Downs curve



Blue curve is
Hellings-Downs
curve (1983)

Effect on stochastic backgrounds

Change the expected shape of the Hellings-Downs curve



Blue curve is
Hellings-Downs
curve (1983)

Conclusions

- Pulsar timing experiments can be used to observe the low frequency GW universe
- A detection is possible within the decade
- These experiments can be used to test gravity in a new way