

Title: Quantum information geometric foundations: beyond the spectral paradigm

Date: May 14, 2015 09:00 AM

URL: <http://pirsa.org/15050090>

Abstract: In the last decade there were proposed several new information theoretic frameworks (in particular, symmetric monoidal categories and "operational" convex sets), allowing for an axiomatic derivation of finite dimensional quantum mechanics as a specific case of a larger universe of information processing theories. Parallel to this, there was an influential development of quantum versions of bayesianism and causality, and relationships between quantum information and space-time structure. In the face of structural problems encountered when moving beyond finite dimensional quantum mechanics, as well as the lack of a mathematically and predictively sound nonperturbative framework for quantum field theories, a question appears: which of the existing structural assumptions of quantum information theory should be relaxed, and how?

In this talk I will present a new approach to the information theoretic foundations of a "general" quantum theory (i.e., beyond quantum mechanics), that is a specific answer to the above question, with a hope to reconstruct both emergent space-times and emergent QFTs. Its mathematical setting is based on using quantum information geometry and integration over noncommutative algebras as structural and conceptual replacements of spectral theory and probability theory, respectively. This corresponds to a paradigmatic change: considering expectation values as more fundamental than eigenvalues. We construct a nonlinear generalisation of quantum kinematics using quantum relative entropies and spaces of states over W^* -algebras. Unitary evolution is generalised to nonlinear hamiltonian flows, while Bayes' and Lueders' rules are generalised to constrained relative entropy maximisations. Combined together, they provide a framework for nonlinear causal inference (information dynamics), that is a generalisation and replacement of completely positive maps. As a result, we construct a large class of information processing theories, containing Hilbert space based QM and probability theory as two special cases. On the conceptual level, we propose a new approach to quantum bayesianism, that is ontically agnostic, intersubjective, and concerned with the relationships between experimental design, model construction, and their mutual predictive verifiability. Finally, we propose a procedure for the emergence of space-times from the geometry of quantum correlations and quantum causality structure, and discuss (briefly) the possibility of reconstructing emergent QFTs.

Quantum information geometric foundations: Beyond the spectral paradigm

Ryszard Paweł Kostecki
Perimeter Institute for Theoretical Physics

arXiv:1505.soon!

Information Theoretic Foundations for Physics
Perimeter Institute, Waterloo

14 May 2015

«Unlike the Riemannian manifolds the quantum mechanical unit spheres do not differ one from another: they are all isomorphic. The worlds of the present-day quantum mechanics thus present a picture of structural monotony: they are all 'painted' on the same standard ideally symmetric surface. The formalism of the quantum theory of infinite systems and quantum field theory is not very different from that. (...) the basic structural framework of the theory is conserved at the cost of quantitative multiplication: when meeting a new level of physical reality the quantum theory responds by simply producing infinite tensor products of its basic structure. (...) It may be that present day quantum theory still represents a relatively primitive stage of development and lacks some essential evolutionary steps leading towards structural flexibility. If this were so, further development would involve a programme opposite to the 'quantization of gravity': instead of modifying general relativity to fit quantum mechanics one should rather modify quantum mechanics to fit general relativity.»

Bogdan Mielnik, 1976, *Quantum logic: is it necessarily orthocomplemented?*

«Unlike the Riemannian manifolds the quantum mechanical unit spheres do not differ one from another: they are all isomorphic. The worlds of the present-day quantum mechanics thus present a picture of structural monotony: they are all 'painted' on the same standard ideally symmetric surface. The formalism of the quantum theory of infinite systems and quantum field theory is not very different from that. (...) the basic structural framework of the theory is conserved at the cost of quantitative multiplication: when meeting a new level of physical reality the quantum theory responds by simply producing infinite tensor products of its basic structure. (...) It may be that present day quantum theory still represents a relatively primitive stage of development and lacks some essential evolutionary steps leading towards structural flexibility. If this were so, further development would involve a programme opposite to the 'quantization of gravity': instead of modifying general relativity to fit quantum mechanics one should rather modify quantum mechanics to fit general relativity.»

Bogdan Mielnik, 1976, *Quantum logic: is it necessarily orthocomplemented?*

$$\frac{\text{SR}}{\text{GR}} = \frac{\text{QM}}{\text{General quantum theory}} ?$$

Plan

1 Nonlinear generalisation of quantum dynamics

- ▶ Geometric structures on quantum states: relative entropies & Poisson brackets
- ▶ Lüders' rules → constrained relative entropy maximisations
- ▶ Unitary evolution → nonlinear hamiltonian flows

2 Geometric framework for quantum information theories beyond quantum mechanics

- ▶ Quantum states = integrals on W^* -algebras
- ▶ Quantum theoretic kinematics = a generalisation of probability theory
- ▶ Quantum theoretic dynamics = a generalisation of causal statistical inference
- ▶ Reconstruction of QM and probability theory
- ▶ Quantum theoretic semantics beyond spectral theory, probabilities, and Born rule
- ▶ Intersubjective bayesian coherence

3 Emergence of space-time theories

- ▶ Space-time geometry = geometry of local correlations and causality
- ▶ Emergent QFTs?



Quantum information models and quantum information distances

trace class operators: $\mathcal{T}(\mathcal{H}) := \{\rho \in \mathfrak{B}(\mathcal{H}) \mid \rho \geq 0, \text{tr}_{\mathcal{H}}|\rho| < \infty\}$

we will consider arbitrary sets of denormalised quantum states: $\mathcal{M}(\mathcal{H}) \subseteq \mathcal{T}(\mathcal{H})^+$

Quantum information distances $D : \mathcal{M}(\mathcal{H}) \times \mathcal{M}(\mathcal{H}) \rightarrow [0, \infty]$ s.t. $D(\rho, \sigma) = 0 \iff \rho = \sigma$.

- E.g.

- ▶ $D_1(\rho, \sigma) := \text{tr}_{\mathcal{H}}(\rho \log \rho - \rho \log \sigma)$ [Umegaki'62]
- ▶ $D_{1/2}(\rho, \sigma) := 2\|\sqrt{\rho} - \sqrt{\sigma}\|_{\mathfrak{B}_2(\mathcal{H})}^2 = 4\text{tr}_{\mathcal{H}}(\frac{1}{2}\rho + \frac{1}{2}\sigma - \sqrt{\rho}\sqrt{\sigma})$ (Hilbert–Schmidt norm²)
- ▶ $D_{L_1(\mathcal{N})}(\rho, \sigma) := \frac{1}{2}\|\rho - \sigma\|_{\mathcal{T}(\mathcal{H})} = \frac{1}{2}\text{tr}_{\mathcal{H}}|\rho - \sigma|$ (L_1 /predual norm)
- ▶ $D_{\gamma}(\rho, \sigma) := \frac{1}{\gamma(1-\gamma)}\text{tr}_{\mathcal{H}}(\gamma\rho + (1-\gamma)\sigma - \rho^{\gamma}\sigma^{1-\gamma})$; $\gamma \in \mathbb{R} \setminus \{0, 1\}$ [Hasegawa'93]
- ▶ $D_{\alpha,z}(\rho, \sigma) := \frac{1}{1-\alpha}\log \text{tr}_{\mathcal{H}}(\rho^{\alpha/z}\sigma^{(1-\alpha)/z})^z$; $\alpha, z \in \mathbb{R}$ [Audenaert–Datta'14]

for $\text{ran}(\rho) \subseteq \text{ran}(\sigma)$, and with all $D(\rho, \sigma) := +\infty$ otherwise.


- Various “quantum geometries” will arise from different additional conditions imposed on pairs $(\mathcal{M}(\mathcal{H}), D)$:
 - ▶ Different choices of $\mathcal{M}(\mathcal{H})$ reflect different assumptions on the available possible knowledge (description of experimental situation).
 - ▶ Different choices of D reflect different assumptions regarding the convention of “best/optimal” estimation/inference.
 - ▶ Both choices are case-to-case-dependent and should be operationally justified.

Quantum entropic projections

Let $\mathcal{Q} \subseteq \mathcal{T}(\mathcal{H})^+$ be such that
for each $\psi \in \mathcal{M}(\mathcal{H})$
there exists a unique solution

$$\mathfrak{P}_{\mathcal{Q}}^D(\psi) := \arg \inf_{\rho \in \mathcal{Q}} \{D(\rho, \psi)\}.$$

It will be called an **entropic projection**.



E.g.

- for $D_{1/2}(\rho, \sigma) = 4\text{tr}_{\mathcal{H}}(\frac{1}{2}\rho + \frac{1}{2}\sigma - \sqrt{\rho}\sqrt{\sigma})$,
consider the entropic projections $\mathfrak{P}_{\mathcal{Q}}^{D_{1/2}}$
where \mathcal{Q} are images of closed convex subspaces $\tilde{\mathcal{Q}} \subseteq \mathcal{K}^+ := \mathfrak{G}_2(\mathcal{H})^+$
under the mapping $\tilde{\mathcal{Q}} \ni \sqrt{\rho} \mapsto \rho \in \mathcal{Q}$.
They coincide with the ordinary projection operators in $\mathfrak{B}(\mathcal{K}) \cong \mathfrak{B}(\mathcal{H} \otimes \mathcal{H}^*)$.
- for $D_1(\rho, \sigma) = \text{tr}_{\mathcal{H}}(\rho \log \rho - \rho \log \sigma)$
and $\mathcal{M}(\mathcal{H}) = \mathcal{T}(\mathcal{H})_1^+$, $\psi \in \mathcal{T}(\mathcal{H})_1^+$, $h \in \mathfrak{B}(\mathcal{H})^{\text{sa}}$, then [Araki'77, Donald'90]

$$\exists! \psi^h := \arg \inf_{\rho \in \mathcal{T}(\mathcal{H})_1^+} \{D_1(\rho, \psi) + \text{tr}_{\mathcal{H}}(\rho h)\}.$$

Quantum measurement, bayesianity, and maximum relative entropy

- Lüders' rules:

$$\rho \mapsto \rho_{\text{new}} := \sum_i P_i \rho P_i \quad (\text{'weak'})$$

$$\rho \mapsto \rho_{\text{new}} := \frac{P \rho P}{\text{tr}_{\mathcal{H}}(P \rho)} \quad (\text{'strong'})$$

- Bub'77'79, Caves–Fuchs–Schack'01, Fuchs'02, Jacobs'02: Lüders' rules should be considered as rules of inference (conditioning) that are quantum analogues of

the Bayes–Laplace rule: $p(x) \mapsto p_{\text{new}}(x) := \frac{p(x)p(b|x)}{p(b)}$.

- Williams'80, Warmuth'05, Caticha&Giffin'06: the Bayes–Laplace rule is a special case of

$$p(x) \mapsto p_{\text{new}}(x) := \arg \inf_{q \in \mathcal{Q}} \{D_1(q, p)\}; \quad D_1(q, p) := \int_{\mathcal{X}} \mu(x) q(x) \log \left(\frac{q(x)}{p(x)} \right).$$

- Douven&Romeijn'12: the Bayes–Laplace rule is also a special case of

$$p \mapsto \arg \inf_{q \in \mathcal{Q}} \{D_1(p, q)\} = \mathfrak{P}_{\mathcal{Q}}^{D_0}(p),$$

where $D_0(p, q) = D_0(q, p)$.

Quantum bayesian inference from quantum entropic projections

- RPK'13'14, F.Hellmann–W.Kamiński–RPK'14:

- 1 weak Lüders' rule is a special case of

$$\rho \mapsto \arg \inf_{\sigma \in \mathcal{Q}} \{D_1(\rho, \sigma)\}$$

with

$$\mathcal{Q} = \{\sigma \in \mathcal{T}(\mathcal{H})^+ \mid [P_i, \sigma] = 0 \forall i\}$$

- 2 strong Lüders' rule derived from

$$\rho \mapsto \arg \inf_{\sigma \in \mathcal{Q}} \{D_1(\rho, \sigma)\}$$

with

$$\mathcal{Q} = \{\sigma \in \mathcal{T}(\mathcal{H})^+ \mid [P_i, \sigma] = 0, \text{tr}_{\mathcal{H}}(\sigma P_i) = p_i \forall i\}$$

under the limit $p_2, \dots, p_n \rightarrow 0$.

- 3 hence, weak and strong Lüders' rules are special cases of quantum entropic projection $\mathfrak{P}_{\mathcal{Q}}^{D_0}$ based on relative entropy $D_0(\sigma, \rho) = D_1(\rho, \sigma)$.

Bayes–Laplace and Lüders' conditionings are special cases of entropic projections
 \Rightarrow "quantum bayesianism \subseteq quantum relative entropism".

Quantum Jeffrey's rule

- Caticha&Giffin'06: under more general constraints, one can derive also Jeffrey's rule (generalising the Bayes–Laplace rule):

$$p(x|\eta) \mapsto p_{\text{new}}(x|\eta) := \sum_{i=1}^n p(x|b_i)\lambda_i = \sum_{i=1}^n \frac{p(x \wedge b_i|\eta)}{p(b_i|\eta)}\lambda_i,$$

where $n \in \mathbb{N}$,

- ▶ $\{b_1, \dots, b_n\}$ is a set of exhaustive and mutually exclusive elements of boolean algebra,
 - ▶ $\lambda_i = p_{\text{new}}(b_i|\eta) \forall i \in \{1, \dots, n\}$,
 - ▶ $p(b_i|\eta) \neq 0$.
- RPK'14: derivation of a quantum analogue of Jeffrey's rule:

$$\mathcal{T}(\mathcal{H})_1^+ \ni \rho \mapsto \rho_{\text{new}} := \arg \inf_{\sigma \in \mathcal{Q}} \{D_1(\rho, \sigma)\} = \sum_{i=1}^n \frac{P_i \rho P_i}{\text{tr}_{\mathcal{H}}(\rho P_i)} \lambda_i \in \mathcal{T}(\mathcal{H})_1^+,$$

where $n \in \mathbb{N}$,

- ▶ $\{P_1, \dots, P_n\} \subseteq \text{Proj}(\mathfrak{B}(\mathcal{H}))$, $\sum_{i=1}^n P_i = \mathbb{I}$, $P_i P_j = \delta_{ij} P_i$,
- ▶ $\lambda_i = \text{tr}_{\mathcal{H}}(\rho_{\text{new}} P_i) \forall i \in \{1, \dots, n\}$,
- ▶ $\text{tr}_{\mathcal{H}}(\rho P_i) \neq 0$.

It generalises Lüders' rule.

Quantum Poisson structure

- Consider the space of self-adjoint trace-class operators: $\mathcal{T}(\mathcal{H})^{\text{sa}} := \mathcal{T}(\mathcal{H}) \cap \mathfrak{B}(\mathcal{H})^{\text{sa}}$.
- It can be equipped with a following real Banach smooth manifold structure:
 - ▶ tangent spaces: $\mathbf{T}_{\phi}(\mathcal{T}(\mathcal{H})^{\text{sa}}) \cong \mathcal{T}(\mathcal{H})^{\text{sa}}$
 - ▶ cotangent spaces: $\mathbf{T}_{\phi}^{\circ}(\mathcal{T}(\mathcal{H})^{\text{sa}}) \cong (\mathcal{T}(\mathcal{H})^{\text{sa}})^{\star} \cong \mathfrak{B}(\mathcal{H})^{\text{sa}}$

- Bóna'91,'00: a Poisson manifold structure on $\mathcal{T}(\mathcal{H})^{\text{sa}}$ is defined by a commutator of an algebra:

$$\{h, f\}(\rho) := \text{tr}_{\mathcal{H}}(\rho \mathbf{i}[\mathbf{d}h(\rho), \mathbf{d}f(\rho)]) \quad \forall f, h \in C^{\infty}(\mathcal{T}(\mathcal{H})^{\text{sa}}; \mathbb{R}) \quad \forall \rho \in \mathcal{T}(\mathcal{H})^{\text{sa}}.$$

- So, if $\mathcal{M}(\mathcal{H}) \subseteq \mathcal{T}(\mathcal{H})^{\text{sa}}$ is a smooth submanifold of $\mathcal{T}(\mathcal{H})^{\text{sa}}$, then every $f \in C^{\infty}(\mathcal{M}(\mathcal{H}); \mathbb{R})$ determines a hamiltonian vector field:

$$\mathfrak{X}_f(\rho) = -\{\cdot, f\}(\rho) = \text{tr}_{\mathcal{H}}(\rho \mathbf{i}[\mathbf{d}(\cdot), \mathbf{d}f(\rho)]).$$

- More generally, we can choose arbitrary real Banach Lie subalgebra \mathcal{A} of $\mathfrak{B}(\mathcal{H})$ such that: (i) it has a unique Banach predual \mathcal{A}_{\star} in $\mathcal{T}(\mathcal{H})$; (ii) there exists at least one $\mathcal{M}(\mathcal{H}) \subseteq \mathcal{T}(\mathcal{H})^{\text{sa}}$ which is a smooth submanifold of \mathcal{A}_{\star} .

Nonlinear quantum hamiltonian dynamics

For each hamiltonian vector field, the corresponding Hamilton equation reads

$$\frac{d}{dt}f(\rho(t)) = \{h, f\}(\rho(t)) = i \operatorname{tr}_{\mathcal{H}}([\rho(t), \mathbf{d}h(\rho(t))] \mathbf{d}f(\rho(t))).$$

The above equation is equivalent to the **Bóna equation** ['91'00]

$$i \frac{d}{dt} \rho(t) = [\mathbf{d}h(\rho(t)), \rho(t)].$$

Hence,

The Poisson structure $\{\cdot, \cdot\}$ induced by a commutator of $\mathfrak{B}(\mathcal{H})$ allows to introduce various nonlinear hamiltonian evolutions on spaces $\mathcal{M}(\mathcal{H})$ of quantum states, generated by arbitrary real-valued smooth functions on $\mathcal{M}(\mathcal{H})$.

The solutions of Bóna equation are state-dependent unitary operators $U(\rho, t)$. They do not form a group, but satisfy a cocycle relationship:

$$U(\rho, t + s) = U((\operatorname{Ad}(U(\rho, t)))(\rho), s)U(\rho, t) \quad \forall t, s \in \mathbb{R}.$$

In a special case, when $h(\rho) = \operatorname{tr}_{\mathcal{H}}(\rho H)$ for $H \in \mathfrak{B}(\mathcal{H})^{\text{sa}}$, the Bóna equation turns to the **von Neumann equation**:

$$i \frac{d}{dt} \rho(t) = [H, \rho(t)].$$

Quantum causal inferences by entropic-hamiltonian dynamics

- **Two elementary geometric structures:**
 - ▶ $D(\cdot, \cdot)$ represents the convention of “best estimation/inference”
 - ▶ $\{h, \cdot\}$ represents a convention of causality (“internal dynamics”)
- **Two elementary forms of quantum dynamics:**
 - ▶ entropic projections \mathfrak{P}_Q^D generated by quantum distances $D(\cdot, \cdot)$
 - ▶ hamiltonian flows w_t^h generated by nonlinear hamiltonian vector fields $\{h, \cdot\}$

A general form of quantum dynamics is defined as a causal inference $\mathfrak{P}_Q^D \circ w_t^h$.

- It generalises unitary evolution followed by a “projective measurement”.
- Postulate: consider the setting of causal inferences $\mathfrak{P}_Q^D \circ w_t^h$ as an alternative to the paradigm of semigroups of CPTP maps.
- Basic idea: every CPTP map can be decomposed into:
 - 1) tensor product of initial state with uncorrelated environment,
 - 2) unitary evolution,
 - 3) projective measurement,
 - 4) partial trace.

It remains to prove that 4 and 3+4 are entropic projections (ongoing work with M.Munk-Nielsen).

Towards new foundations

Idea:

- consider spaces $\mathcal{M}(\mathcal{H})$ as fundamental
- allow any nonlinear functions $\mathcal{M}(\mathcal{H}) \rightarrow \mathbb{R}$ as observables
- define geometry of $\mathcal{M}(\mathcal{H})$ by means of $D(\cdot, \cdot)$ and $\{\cdot, \cdot\}$
- define dynamics of $\mathcal{M}(\mathcal{H})$ by means of $\mathfrak{P}_{\mathbb{Q}}^D(\cdot, \cdot)$ and $w_t^{\{h, \cdot\}}$

Questions:

- what's up with Hilbert spaces? (are they necessary? if not, then what?)
- what's up with spectral theory, probability, Born rule, etc?

Answers:

- replace Hilbert spaces by W^* -algebras
- replace density matrices by positive integrals on W^* -algebras
- this setting is an exact generalisation of Kolmogorov's measure theoretic setting for probability theory
- build up all remaining semantics for quantum theory
in the analogy to semantics of probability theory and statistical inference
(hence: no Born rule, no probabilities, no spectral theory)

Towards new foundations

Idea:

- consider spaces $\mathcal{M}(\mathcal{H})$ as fundamental
- allow any nonlinear functions $\mathcal{M}(\mathcal{H}) \rightarrow \mathbb{R}$ as observables
- define geometry of $\mathcal{M}(\mathcal{H})$ by means of $D(\cdot, \cdot)$ and $\{\cdot, \cdot\}$
- define dynamics of $\mathcal{M}(\mathcal{H})$ by means of $\mathfrak{P}_{\mathbb{Q}}^D(\cdot, \cdot)$ and $w_t^{\{h, \cdot\}}$

Questions:

- what's up with Hilbert spaces? (are they necessary? if not, then what?)
- what's up with spectral theory, probability, Born rule, etc?

Answers:

- replace Hilbert spaces by W^* -algebras
- replace density matrices by positive integrals on W^* -algebras
- this setting is an exact generalisation of Kolmogorov's measure theoretic setting for probability theory
- build up all remaining semantics for quantum theory
in the analogy to semantics of probability theory and statistical inference
(hence: no Born rule, no probabilities, no spectral theory)

Probability theory:

- Underlying structure: measure space (\mathcal{X}, μ)
- Main spaces: **Probabilistic models:**

$$\mathcal{M}(\mathcal{X}, \mu) \subseteq L_1(\mathcal{X}, \mu)^+ := \{p : \mathcal{X} \rightarrow \mathbb{R} \mid \int_{\mathcal{X}} \mu |p| < \infty, p \geq 0\}$$

- e.g. Gaussian models: $\{p(x, (m, s)) = \frac{1}{\sqrt{2\pi s}} e^{-\frac{(x-m)^2}{2s^2}} \mid (m, s) \in \Theta \subseteq \mathbb{R} \times \mathbb{R}^+\}$.
- Observables (estimators): functions $f : \mathcal{X} \rightarrow \mathbb{R}$
- The mapping $L_1(\mathcal{X}, \mu) \times L_\infty(\mathcal{X}, \mu) \ni (p, f) \mapsto \int_{\mathcal{X}} \mu p f \in \mathbb{R}$ determines Banach space duality $L_1(\mathcal{X}, \mu)^* \cong L_\infty(\mathcal{X}, \mu)$.

Quantum mechanics:

- Underlying structure: Hilbert space \mathcal{H}
- Main spaces: **Spaces of density matrices:**

$$\mathcal{M}(\mathcal{H}) \subseteq \mathcal{T}(\mathcal{H})^+ := \{\rho \in \mathfrak{B}(\mathcal{H}) \mid \text{tr}_{\mathcal{H}}(|\rho|) < \infty, \rho \geq 0\}$$

- e.g. Gibbs states: $\{e^{-\beta H} \mid \beta \in]0, \infty[\}$, for a fixed self-adjoint H .
- Observables: self-adjoint operators $x : \mathcal{H} \rightarrow \mathcal{H}$
- The mapping $\mathcal{T}(\mathcal{H}) \times \mathfrak{B}(\mathcal{H}) \ni (\rho, x) \mapsto \text{tr}_{\mathcal{H}}(\rho x) \in \mathbb{C}$ determines Banach space duality $\mathcal{T}(\mathcal{H})^* \cong \mathfrak{B}(\mathcal{H})$.

W^* -algebras and integration

- A W^* -algebra \mathcal{N} :
 - ▶ an algebra over \mathbb{R} or \mathbb{C} with unit \mathbb{I} ,
 - ▶ with $*$ operation s.t. $(xy)^* = y^*x^*$, $(x + y)^* = x^* + y^*$, $(x^*)^* = x$, $(\lambda x)^* = \lambda^*x^*$,
 - ▶ that is also a Banach space,
 - ▶ with \cdot , $+$, $*$ continuous in the norm topology (implied by the condition $\|x^*x\| = \|x\|^2$),
 - ▶ such that there exists a Banach space \mathcal{N}_* satisfying the Banach space duality:
 $(\mathcal{N}_*)^* \cong \mathcal{N}$,
- Special cases:
 - ▶ if \mathcal{N} is commutative
then \exists a measure space (\mathcal{X}, μ) s.t. $\mathcal{N} \cong L_\infty(\mathcal{X}, \mu)$ and $\mathcal{N}_* \cong L_1(\mathcal{X}, \mu)$
 - ▶ if \mathcal{N} is "type I factor"
then \exists a Hilbert space \mathcal{H} s.t. $\mathcal{N} \cong \mathfrak{B}(\mathcal{H})$ and $\mathcal{N}_* \cong \mathcal{T}(\mathcal{H})$.
- Hence, the element $\phi \in (\mathcal{N}_*)^+$ provides a joint generalisation of probability density and of density operator. By means of embedding of \mathcal{N}_* into \mathcal{N}^* , it is also an integral on \mathcal{N} .
- **Key fact:** The above setting allows to develop full-fledged integration theory on noncommutative W^* -algebras, which generalises integration theory on measure spaces (with partial integration, conditional expectations, $L_p(\mathcal{N})$ spaces,...).

New kinematics: quantum models and observables

General quantum information models:

For any W^* -algebra \mathcal{N} , $\mathcal{M}(\mathcal{N})$ will be defined as an arbitrary subset of a positive part of a Banach predual space of \mathcal{N} , $\mathcal{M}(\mathcal{N}) \subseteq \mathcal{N}_*^+$.

Special cases:

- \mathcal{N} is commutative $\Rightarrow \mathcal{M}(\mathcal{N}) = \mathcal{M}(\mathcal{X}, \mu)$
- \mathcal{N} is type I factor $\Rightarrow \mathcal{M}(\mathcal{N}) = \mathcal{M}(\mathcal{H})$.

We **do not** assume that:

- $\mathcal{M}(\mathcal{N})$ is convex (\iff probabilistic mixing)
- $\mathcal{M}(\mathcal{N})$ is smooth (\iff asymptotic estimation)
- $\mathcal{M}(\mathcal{N})$ is normalised (\iff frequentist interpretation)

Observables:

Observables are defined as arbitrary functions $f : \mathcal{M}(\mathcal{N}) \rightarrow \mathbb{R}$.

Hence: smooth observables define hamiltonian vector fields.

Each “observable in the old sense” $x \in \mathcal{N}^{\text{sa}}$ determines a corresponding “observable in the new sense” by $f_x(\phi) := \phi(x)$.

New kinematics: quantum information geometry

- **Main change:** Consider expectation values as more fundamental than eigenvalues
⇒ foundational role of spectral theory replaced by quantum information geometry
- **Kinematic setting:**
 - (1) **spaces:** Hilbert spaces \mathcal{H} of **eigenvectors**
→ spaces $\mathcal{M}(\mathcal{N})$ of denormalised **expectation functionals** on W^* -algebras \mathcal{N} .
 - (2) **observables:** linear functions $\mathcal{H} \rightarrow \mathbb{R}$ that have real eigenvalues
→ nonlinear real valued functions $\mathcal{M}(\mathcal{N}) \rightarrow \mathbb{R}$.
 - (3) **geometry:** geometry of Hilbert spaces \mathcal{H} **defined by scalar product** $\langle \cdot, \cdot \rangle$
→ geometry of spaces $\mathcal{M}(\mathcal{N})$ **defined by quantum relative entropies** $D(\cdot, \cdot)$ and **quantum Poisson structures** $\{\cdot, \cdot\}$.
- **Two fundamental geometric structures on $\mathcal{M}(\mathcal{N})$:**
 - a) **Quantum distances** $D(\cdot, \cdot)$
 - ★ large variety of choices
 - ★ allows to derive riemannian geometry (via $\partial_i \partial_j D$) and Hilbert space projective geometry (via $\mathfrak{P}_{\mathbb{Q}}^D$ for $D = D_{1/2}$) as special cases
 - b) **Quantum Poisson structures** $\{\cdot, \cdot\}$
 - ★ depend on the choice of a real Banach Lie subalgebra of \mathcal{N}
 - ★ generalises symplectic geometry
- No Hilbert spaces, no probability theory in foundations (derived as special cases)


Backwards compatibility

1 Reconstruction of quantum mechanics:

- ▶ \mathcal{N} : type I W^* -algebras
- ▶ $\mathcal{M}(\mathcal{N})$: normalised states
- ▶ D : $D_{1/2}$ or D_0
- ▶ $\{, \cdot\}$: generated by Banach Lie algebra \mathcal{N}^{sa}
- ▶ observables: affine functions on $\mathcal{M}(\mathcal{N})$

2 Reconstruction of probability theory:

- ▶ \mathcal{N} : commutative algebras
- ▶ $\mathcal{M}(\mathcal{N})$: normalised states
- ▶ D : arbitrary
- ▶ $\{, \cdot\}$: trivialises for commutative algebras
- ▶ observables: arbitrary or affine functions on $\mathcal{M}(\mathcal{N})$



Smooth quantum information geometries

Under some conditions, D induces a generalisation of smooth riemannian geometry on $\mathcal{M}(\mathcal{N})$.

- Jenčová'05: a general construction of smooth manifold structure on the space of all strictly positive states over arbitrary W^* -algebra.
- E.g. $\mathcal{M}(\mathcal{H}) := \{\rho(\theta) \in \mathcal{T}(\mathcal{H}) \mid \rho(\theta) > 0, \theta \in \Theta \subseteq \mathbb{R}^n \text{ open}, \theta \mapsto \rho(\theta) \text{ smooth}\}$
- Eguchi'83/Ingarden et al'82/Lesniewski–Ruskai'99/Jenčová'04:
Every smooth distance D with positive definite hessian determines a riemannian metric \mathbf{g}^D and a pair $(\nabla^D, \nabla^{D^\dagger})$ of torsion-free affine connections:

$$\begin{aligned}\mathbf{g}_\phi(u, v) &:= -\partial_{u|\phi} \partial_{v|\omega} D(\phi, \omega)|_{\omega=\phi}, \\ \mathbf{g}_\phi((\nabla_u)_\phi v, w) &:= -\partial_{u|\phi} \partial_{v|\phi} \partial_{w|\omega} D(\phi, \omega)|_{\omega=\phi}, \\ \mathbf{g}_\phi(v, (\nabla_u^\dagger)_\phi w) &:= -\partial_{u|\omega} \partial_{w|\omega} \partial_{v|\phi} D(\phi, \omega)|_{\omega=\phi},\end{aligned}$$

which satisfy the characteristic equation of the Norden['37]–Sen['44] geometry,

$$\mathbf{g}^D(u, v) = \mathbf{g}^D(\mathbf{t}_c^{\nabla^D}(u), \mathbf{t}_c^{\nabla^{D^\dagger}}(v)) \quad \forall u, v \in \mathbf{T}\mathcal{M}(\mathcal{N}).$$

- A riemannian geometry $(\mathcal{M}(\mathcal{N}), \mathbf{g}^D)$ has Levi-Civita connection $\bar{\nabla} = (\nabla^D + \nabla^{D^\dagger})/2$.
- E.g., $\mathcal{M}(\mathcal{N}) = \mathcal{T}(\mathcal{H}) \cap \{\rho > 0, \text{tr}_{\mathcal{H}}(\rho) = 1\}$ and $D_1(\rho, \sigma) = \text{tr}(\rho \log \rho - \rho \log \sigma)$ give Mori['55]–Kubo['56]–Bogolyubov['62] \mathbf{g}^{D_1} and Nagaoka['94]–Hasegawa['95] $(\nabla^{D_1}, \nabla^{D_1^\dagger})$:

$$\mathbf{g}_\rho^{D_1}(x, y) = \text{tr} \left(\int_0^\infty d\lambda x \frac{1}{\lambda \mathbb{1} + \rho} y \frac{1}{\lambda \mathbb{1} + \rho} \right), \quad \mathbf{t}_{\rho, \omega}^{\nabla^{D_1}}(x) = x - \text{tr}(\omega x), \quad \mathbf{t}_{\rho, \omega}^{\nabla^{D_1^\dagger}}(x) = x.$$

Quantum mechanics as a local theory

- Apart from tangent bundle $\bigcup_{\phi} \mathbf{T}_{\phi}\mathcal{M}(\mathcal{N})$, there is also a bundle of complex (GNS) Hilbert spaces $\mathcal{H}\mathcal{M}(\mathcal{N}) \rightarrow \mathcal{M}(\mathcal{N})$.
- Vectors in $\mathbf{T}_{\phi}\mathcal{M}(\mathcal{N})$ are defined by self-adjoint operators, which can be represented uniquely as elements of $(\mathcal{H}_{\phi}\mathcal{M}(\mathcal{N}))_{\mathbb{R}}$.
- Under some (mild) conditions: $\mathbf{T}_{\phi}\mathcal{M}(\mathcal{N}) \sqsubseteq (\mathcal{H}_{\phi}\mathcal{M}(\mathcal{N}))_{\mathbb{R}}$.
- Thus, as opposed to C^* -algebraic approach:
 - ▶ Spaces of quantum states are equipped with rich geometric structure, allowing for model construction, state estimation, and nonlinear dynamics.
 - ▶ Quantum mechanics is reconstructed not only as a global special case of a framework, but also is present locally at each point of a manifold, as an extension of a tangent space.
 - ▶ Our framework allows also for a geometric description of renormalisation procedures (see Cedric Bény's talk).


Jacobson'95: Einstein equations *from* space-time thermodynamics

Consider:

- a space-time (\mathcal{M}, g_{ab})
- a point $p \in \mathcal{M}$
- a small 2-dimensional surface element \mathcal{P}
- a Killing vector field χ^a generating local boost orthogonal to \mathcal{P}

Define:

- a **local causal horizon** \mathcal{H} as a boundary of the past of \mathcal{P} , generated by χ^a
- a **heat flow** δQ as an energy flux across a local causal horizon:
$$\delta Q := \int_{\mathcal{H}} d\Sigma^a T_{ab} \chi^b$$
- a **temperature** T as an Unruh temperature associated with a uniformly accelerated observer.



Assume:

- that **entropy** S is proportional to the area of \mathcal{H} : $S = \lambda A$
- that Clausius' law holds: $\delta Q = T dS$.

Then:

$$R_{ab} - \frac{1}{2} R g_{ab} + \Lambda g_{ab} = \frac{2\pi}{\lambda} T_{ab}.$$

Emergent space-times

- **Basic idea:** Consider a principle of equivalence of euclidean QSM with lorentzian QFT via Wick rotation as a fundamental principle, analogous to $m_{\text{grav}} = m_{\text{inert}}$.
- **Basic mathematical data:**
 - ▶ $\mathbf{g}_\rho^D(\cdot, \cdot)$ is a correlation functional, representing a convention of a local (asymptotic) estimation/inference at ρ .
 - ▶ $\{h(\rho), \cdot\}$ is a dynamical evolution, representing a convention of a local temporal causality at ρ .
- **Required assumptions:**
 - ▶ choice of a manifold Σ that is determined by operational parameters of measurement of "space" and "time"
 - ▶ split $\mathcal{M}(\mathcal{N}) \cong \Sigma \times \widetilde{\mathcal{M}}(\mathcal{N})$
 - ▶ $\{h(\rho), \cdot\}$ is well defined on Σ
- **Implementation:**
 - ▶ consider a riemannian metric \mathbf{g}_Σ^D induced by \mathbf{g}^D on Σ
 - ▶ "Poincaré–Wick rotation" of \mathbf{g}_Σ^D to a lorentzian $\hat{\mathbf{g}}_\Sigma^{D,h}$ along a vector field $\{h, \cdot\}$:

$$\mathbf{g}_\Sigma^D = \mathbf{g}_\perp^D + \mathbf{e}_h \otimes \mathbf{e}_h \mapsto \mathbf{g}_\perp^D - \mathbf{e}_h \otimes \mathbf{e}_h =: \hat{\mathbf{g}}_\Sigma^{D,h},$$

where \mathbf{g}_\perp^D is a riemannian metric induced by \mathbf{g}_Σ^D on the submanifolds orthogonal to \mathbf{e}_h ,

while $\mathbf{e}_h := \frac{\mathbf{g}_\Sigma^D(\{h, \cdot\}, \cdot)}{\sqrt{\mathbf{g}_\Sigma^D(\{h, \cdot\}, \{h, \cdot\})}}$ is a normalised 1-form of $\{h, \cdot\}$.

An emergent space-time is a triple $(\Sigma, \hat{\mathbf{g}}_\Sigma^{D,h}, \mathbf{e}_h)$.

Emergent space-times

- **Basic idea:** Consider a principle of equivalence of euclidean QSM with lorentzian QFT via Wick rotation as a fundamental principle, analogous to $m_{\text{grav}} = m_{\text{inert}}$.
- **Basic mathematical data:**
 - ▶ $\mathbf{g}_\rho^D(\cdot, \cdot)$ is a correlation functional, representing a convention of a local (asymptotic) estimation/inference at ρ .
 - ▶ $\{h(\rho), \cdot\}$ is a dynamical evolution, representing a convention of a local temporal causality at ρ .
- **Required assumptions:**
 - ▶ choice of a manifold Σ that is determined by operational parameters of measurement of "space" and "time"
 - ▶ split $\mathcal{M}(\mathcal{N}) \cong \Sigma \times \widetilde{\mathcal{M}}(\mathcal{N})$
 - ▶ $\{h(\rho), \cdot\}$ is well defined on Σ
- **Implementation:**
 - ▶ consider a riemannian metric \mathbf{g}_Σ^D induced by \mathbf{g}^D on Σ
 - ▶ "Poincaré–Wick rotation" of \mathbf{g}_Σ^D to a lorentzian $\hat{\mathbf{g}}_\Sigma^{D,h}$ along a vector field $\{h, \cdot\}$:

$$\mathbf{g}_\Sigma^D = \mathbf{g}_\perp^D + \mathbf{e}_h \otimes \mathbf{e}_h \mapsto \mathbf{g}_\perp^D - \mathbf{e}_h \otimes \mathbf{e}_h =: \hat{\mathbf{g}}_\Sigma^{D,h},$$

where \mathbf{g}_\perp^D is a riemannian metric induced by \mathbf{g}_Σ^D on the submanifolds orthogonal to \mathbf{e}_h ,

while $\mathbf{e}_h := \frac{\mathbf{g}_\Sigma^D(\{h, \cdot\}, \cdot)}{\sqrt{\mathbf{g}_\Sigma^D(\{h, \cdot\}, \{h, \cdot\})}}$ is a normalised 1-form of $\{h, \cdot\}$.

An emergent space-time is a triple $(\Sigma, \hat{\mathbf{g}}_\Sigma^{D,h}, \mathbf{e}_h)$.

Emergent space-times: comments

- Operational assumptions that may lead to derivation of 4-dimensionality of Σ ?
→ see the talk of Markus Müller for very interesting ideas.
- Instead of a split $\mathcal{M}(\mathcal{N}) \cong \Sigma \times \widetilde{\mathcal{M}}(\mathcal{N})$, one can consider also a nontrivial fibre bundle with locally (but not globally) defined operational space-times $\pi : \mathcal{M}(\mathcal{N}) \rightarrow \Sigma$.
- Every section of a bundle $\widetilde{\mathcal{M}}(\mathcal{N})$ over Σ defines a global quantum state $\phi(\xi)$ over space-time, and this determines a bundle $\mathcal{H}\Sigma \rightarrow \Sigma$ of GNS Hilbert spaces $\mathcal{H}_{\phi(\xi)}\Sigma$, $\xi \in \Sigma$.
- This allows to use Prugovečki's approach to defining quantum propagators over a curved space-time. \Rightarrow construction of emergent QFTs over curved space-time.

Overview

1 Nonlinear generalisation of quantum dynamics

- ▶ Geometric structures on quantum states: relative entropies & Poisson brackets
- ▶ Lüders' rules → constrained relative entropy maximisations
- ▶ Unitary evolution → nonlinear hamiltonian flows

2 Geometric framework for quantum information theories beyond quantum mechanics

- ▶ Quantum states = integrals on W^* -algebras
- ▶ Quantum theoretic kinematics = a generalisation of probability theory
- ▶ Quantum theoretic dynamics = a generalisation of causal statistical inference
- ▶ Reconstruction of QM and probability theory
- ▶ Quantum theoretic semantics beyond spectral theory, probabilities, and Born rule
- ▶ Intersubjective bayesian coherence

3 Emergence of space-time theories

- ▶ Space-time geometry = geometry of local correlations and causality
- ▶ Emergent QFTs?

References

Main paper:

- RPK, 2015, Towards quantum information geometric foundations, soon on arXiv!

Earlier results and insights:

- RPK, 2014, Lüders' and quantum Jeffrey's rules as entropic projections, arXiv:1408.3502.
- F. Hellmann, W. Kamiński, RPK, 2014, Quantum collapse rules from maximum relative entropy principle, arXiv:1407.7766.
- RPK, 2013, W^* -algebras and noncommutative integration, arXiv:1307.4818.
- KostECKI R.P., 2013, Information geometric foundations of quantum theory, Ph.D. thesis, Institute for Theoretical Physics, University of Warsaw.
- Duch P., RPK, 2011, Quantum Schwarzschild space-time, arXiv:1110.6566.
- RPK, 2011, Information dynamics and new geometric foundations for quantum theory, Växjö 2011 Proceedings, AIP Conf. Proc. **1424**, 200. arXiv:1110.4492.
- RPK, 2011, On principles of inductive inference, MAXENT 2011 Proceedings, AIP Conf. Proc. **1443**, 22. arXiv:1109.3142.
- RPK, 2011, The general form of γ -family of quantum relative entropies, Open Sys. Inf. Dyn. **18**, 191. arXiv:1106.2225.
- RPK, 2010, Quantum theory as inductive inference, MAXENT 2010 Proceedings, AIP Conf. Proc. **1305**, 28. arXiv:1009.2423.