

Title: Brane webs, 5d gauge theories and 6d  $N=(1,0)$  SCFT's

Date: Nov 24, 2015 02:00 PM

URL: <http://pirsa.org/15110098>

Abstract: <p>Some 5d gauge theories have a 6d  $N=(1,0)$  SCFT as their UV completion. Given such 5d gauge theory we desire to determine its 6d UV completion. In this talk, I will present a method to do this for 5d gauge theories that can be engineered in string theory by brane webs. This can then be applied to study compactification of 6d  $N=(1,0)$  SCFT's on a torus.</p>

# Motivation

- Investigate the relation between 5d  $N=1$  gauge theories and 6d  $N=(1,0)$  SCFT's.
- A 5d  $N=1$  gauge theory sometimes has 6d  $N=(1,0)$  SCFT as its UV completion.
- Would like to determine the 6d  $N=(1,0)$  SCFT UV completion.
- The 5d theory can then be used to study the 6d SCFT.
- This has interesting implications on the compactification of the 6d SCFT to 4d on a torus.

# Outline

## 1. Introduction

- Generalities of 5d gauge theories and their description using brane webs
- 6d  $N=(1,0)$  SCFT's and their description using brane systems
- 5d gauge theories with a 6d SCFT UV completion

## 2. Determining the 6d SCFT UV completion using brane webs

## 3. Implication for the compactification of 6d SCFT's on a torus

## 4. Conclusions

# 1. Introduction: 5d gauge theories

- 5d gauge theories are non-renormalizable.
- Yet, in the  $N=1$  SUSY case a UV fixed point may exist.
- 5d gauge theory generated from a 5d SCFT by a mass deformation associated with the coupling constant.
- Exhibit interesting non-perturbative properties such as enhancement of symmetry.


[Seiberg, 1996]

# Enhancement of symmetry

- In 5d every simple non-abelian gauge group has a topologically conserved  $U(1)$  current:  $J \sim * \text{Tr} F \wedge F$ .
- This current is carried by instanton particles.
- These can provide additional conserved currents leading to an enhancement of symmetry.
- A classic example is an  $SU(2) + N_F F$ , with  $N_F < 8$ , where instantons enhance the classical  $U(1) \times SO(2N_F)$  global symmetry to  $E_{N_F+1}$  [Seiberg, 1996].

# 5-Brane webs

- Convenient way to represent 5d gauge theories.
- Parameters represented by position of external branes.
- Moduli represented by moving internal branes.
- Fixed point realized when all the 5-branes intersect at a point.



[Aharony, Hanany, 1997]

[Aharony, Hanany, Kol, 1997]

## 6d $N=(1,0)$ SCFT's

- Made from: hypermultiplets, vector multiplets and tensor multiplets.
- Tensor multiplet has a self-dual tensor and real scalar as its bosonic degrees of freedom.
- Scalar can acquire a vev leading to a moduli space called the tensor branch.


## Examples of 6d $N = (1,0)$ SCFT's

- Rank  $N$  E-string theories: made of  $N$  tensor multiplets with an  $E_8$  global symmetry. Described in string theory by  $N$  M5-branes near an M9-plane [Ganor, Hanany, 1996].
- 6d gauge theories are non-renormalizable, but some have a SCFT UV completion. This requires adding tensor multiplet with the coupling constant identified with the scalar vev [Seiberg, 1997].
- 6d gauge theories suffer from gauge anomalies whose cancellation usually requires adding a tensor multiplet. These also constrain the matter content.



# Brane configurations for 6d SCFT's

- Consider the system made from an  $O8^-$ -plane and  $2N$  D6 branes partitioned by an NS5-brane. Gives an  $USp(2N) + (2N + 8)F$  gauge theory on the D6 branes [Hanany, Zaffarony, 1998].
- D6-brane charge conservation leads to anomaly cancelation condition.
- Motion of the NS5-brane is identified with the tensor branch.
- Adding more NS5-branes leads to a  $USp \times SU \dots \times SU$  quiver.
- Adding an NS5-brane stuck on the  $O8^-$ -plane changes the first group to  $SU(2N) + 1AS$ .
- One can also add D8-branes, leading to additional flavors.



# The 6d anomaly polynomial

- Besides gauge anomalies there can also be anomalies involving global currents.
- These include the Lorentz group, the  $SU(2)_R$  symmetry and flavor symmetries.
- Conveniently packaged in an 8-form called the anomaly polynomial.
- There are known techniques to calculate it [Ohmori, Shimizu, Tachikawa, Yonekura, 2014].
- For very Higgsable 6d SCFT's compactified on a torus: related to the central charges of the resulting 4d theory [Ohmori, Shimizu, Tachikawa, Yonekura, 2015].

# 5d gauge theories with a 6d SCFT UV completion


- Some 5d  $N=1$  gauge theories have 6d  $N=(1,0)$  SCFT's as their UV completion.
- 5d gauge theory generated from a 6d SCFT by a mass deformation associated with compactification of the 6d SCFT on a circle of radius  $R$ .
- 5d gauge theory has a space of relevant deformations:  $\frac{1}{R}$  and holonomies of background gauge fields.
- By taking all these to zero one approaches the 6d SCFT.

# 5d gauge theories with a 6d SCFT UV completion: examples


- Maximally supersymmetric Yang-Mills and the 6d (2,0) theory [Douglas, 2011][Lambert, Papageorgakis, Schmidt-Sommerfeld, 2011].
- $SU(2) + 8F$  and the rank 1 E-string theory [Ganor, Morrison, Seiberg, 1997].
- Apparent from the string theory construction: a D4-brane immersed in an  $O8^- + 8D8$ -branes.
- Another hint from the gauge theory perspective: instantons lead to an enhanced affine  $E_8^{(1)}$  global symmetry.

# Example: $SU(N + 2)_0 + (2N + 8)F$ and the $(D_{N+4}, D_{N+4})$ conformal matter

[Hayashi, Kim, Lee, Taki, Yagi, 2015]



T - duality  
+  $O7^-$  resolution




$(D_{N+4}, D_{N+4})$  conformal matter: one dimensional tensor branch leading to 6d gauge theory  $USp(2N) + (2N + 8)F$ .

$SU(N + 2) + (2N + 8)F$  gauge.  
Has an affine  $D_{2N+8}^{(1)}$  global symmetry.


## 2. Determining the 6d SCFT UV completion using brane webs

- Many other 5d gauge theories suspected to lift to 6d.
- With brane web presentations: quivers of  $SU$  groups with fundamentals, or with  $USp$  or  $SU + 1AS$  ends.
- We wish to determine the 6d SCFT also for these.
- The strategy is to first generalize the previous example to a certain class of theories.
- Additional theories can then be determined by connecting them to theories in this class via Higgs branch flows.
- By following the breakings we can determine the 6d SCFT.

# Generalization: adding NS5-branes




# Higgs branch flow



- Higgs branch flow in the 5d theory described by breaking a 5-brane on a 7-brane. This forces several 5-branes to end on the same 7-brane.
- Correspondingly in the 6d theory: breaking a D6-brane on a D8-brane.




# Higgs branch flow and quiver tails



- This can be used to map the starting quiver to more general cases.
- Previous construction instruct us to do the same mapping on the 6d SCFT.

# A simple example




# For what theories can this be applied?


- Appears to be applicable only for linear  $SU$  quivers with fundamentals.
- However, the method can still be applied by manipulating the brane web to one related to the previous cases by Higgs branch flows.
- This works for a large class of 5d gauge theories.
- This leads us to conjecture that this can be applied to any 5d gauge theory with a brane web presentation.

### 3. Implication to the compactification of the 6d SCFT to 4d on a torus


- This method can also be used to determine the result of compactifying the 6d SCFT on a torus.
- This can be checked by comparing the central charges with the ones expected from the 6d anomaly polynomial.




# Implication to the compactification of the 6d SCFT to 4d on a torus



# Implication to the compactification of the 6d SCFT to 4d on a torus




# Implication to the compactification of the 6d SCFT to 4d on a torus



- Starting from the brane web for the 5d  $(N + 2)F + SU(N)X \dots X SU(N) + (N + 1)F$  quiver gauge theory, we perform a long sequence of 7-brane manipulations consisting of moving 7-branes accompanied by Hanany-Witten transitions.

# Implication to the compactification of the 6d SCFT to 4d on a torus



- After intense 7-brane manipulation we arrive to the web on the left. For ease of presentation only the external legs are drawn.
- Right: the associated 4d class S theory.



## 4. Conclusions

- Some 5d gauge theories have a 6d SCFT UV completion. This 6d SCFT can be determined using Higgs branch flow for 5d gauge theories with a brane web representation.
- This can be used to determine the 4d SCFT resulting from the compactification of the 6d SCFT on a torus.

## Open questions

- Can every 5d gauge theory, with an brane web representation, be treated in this way?
- Additional systems: webs in the presence of an  $O7$ -plane,  $O5$ -plane.

## 4. Conclusions

- Some 5d gauge theories have a 6d SCFT UV completion. This 6d SCFT can be determined using Higgs branch flow for 5d gauge theories with a brane web representation.
- This can be used to determine the 4d SCFT resulting from the compactification of the 6d SCFT on a torus.

## Open questions

- Can every 5d gauge theory, with an brane web representation, be treated in this way?
- Additional systems: webs in the presence of an  $O7$ -plane,  $O5$ -plane.