

Title: Limits on locality from gravitational dressing

Date: Nov 26, 2015 10:30 AM

URL: <http://pirsa.org/15110100>

Abstract: <p>In quantum gravity, observables must be diffeomorphism-invariant. Such observables are nonlocal, in contrast with the standard assumption of locality in flat spacetime quantum field theory. I will show how to construct 'gravitationally dressed' observables in linearized gravity that become local in the weak gravity limit, and whose corrections to exact locality are characterized by the Newtonian potential. One can attempt to make these observables more local by concentrating gravitational field lines into a smaller solid angle. In AdS₃ gravity I will show that nonperturbatively there are sharp limits to how much the gravitational dressing can be concentrated.</p>

<p>Based on arXiv:1507.07921 with Steve Giddings, and arXiv:1510.00672 with Don Marolf and Eric Mintun.</p>

Limits on locality from gravitational dressing

arXiv 1507.07921 WD & Steve Giddings

1510.00672 WD, Don Marolf, Eric Mintun

Quantum gravity

QFT local operators $\phi(x)$.

$$[\phi(x), \phi(x')] = 0, \quad (x-x')^2 \rightarrow 0$$

GR diffeomorphism-invariant.

Locality

7.07921

10.00672

W-

WD, D...

Quantum gravity

QFT local operators $\phi(x)$.

$$[\phi(x), \phi(x')] = 0 \quad (x-x')^2 > 0$$

GR

diffeomorphism-invariant.

$$\mathbb{I} = \int d^4x \sqrt{g} \mathcal{O}(x)$$

Plan

① 3+1 linearized gravity + scalar.

$$\bar{\Phi}(x) \xrightarrow{G \rightarrow 0} \phi(x)$$

$\bar{\Phi}$ invariant.

$$[\bar{\Phi}(x), \bar{\Phi}(x')].$$


② 2+1 AdS.

QED Dirac (1953)

$$\phi(x) \rightarrow e^{iq\Lambda(x)} \phi(x)$$



$$A_n(x) \rightarrow A_n(x) - \partial_n \Lambda(x)$$

$$\bar{\Phi}_w(x) = \phi(x) e^{iq \int_0^\infty ds A_z(x+s\hat{z})}$$



$$[E^z(x), \bar{\Phi}_w(x')] = -q \int d^2(x_\perp - x'_\perp) \Theta(z-z') \bar{\Phi}_w(x')$$

$$\bar{\Phi}_c(x) = \phi(x) \exp(i \int d^3x' E_c^i(x'-x) A_i(x'))$$



Gravity

$$g_{\mu\nu} = \eta_{\mu\nu} + \kappa h_{\mu\nu} \quad \kappa = \sqrt{32\pi G}$$

$$\phi \rightarrow \phi - \kappa \xi^\mu \partial_\mu \phi$$


$$h_{\mu\nu} \rightarrow h_{\mu\nu} - \partial_\mu \xi_\nu - \partial_\nu \xi_\mu$$

Wilson line

$$X^\mu(z) = x^\mu + s \hat{z}^\mu + V^\mu(s)$$

$$V_W^\mu = - \int_0^\infty ds s \Gamma_{zz}^\mu(x + s \hat{z}) \quad \tau \text{ s.t.}$$

$$\bar{\Phi}_W(x) = \phi(x + V_W) = \phi(x) + V_W^\mu(x) \partial_\mu \phi(x)$$



Coulomb

$$V_C^m(o) = -\frac{1}{4\pi} \int d^3x \frac{\Gamma^m}{rr} \cdot \frac{1}{r}$$

[P⁴_{ADM}, Φ]

$$\partial^\mu \phi = \rho^\mu \phi$$

$$\Phi_C'(o) = \phi(o) + V_C^m(o) \partial_m \phi(o)$$


$$[h_{\mu\nu}(x), \Phi_C(o)] = [h_{\mu\nu}(x), V^{\lambda}(o)] \partial_\lambda \phi(o)$$

Particle at rest $\partial_o \phi = im\phi$

$$= \left[\underbrace{\frac{\chi m}{16\pi r} (\hat{t}_m \hat{t}_\nu + \hat{r}_m \hat{r}_\nu)}_{(1m) \text{ Schwarzschild}} + \underbrace{\partial_m \zeta_\nu + \partial_\nu \zeta_m}_{\text{diff}} \right] \phi(o)$$


Commutators (equal time)

$$\begin{aligned}
 [\dot{\Phi}_W(x), \Phi_W(x')] &= [\dot{V}_W^m(x), V_W^y(x')] \partial_m \phi(x) \partial_y \phi(x') + \dots \\
 &= -\frac{i4G}{\pi^2} \partial_z \phi(x) \partial_z \phi(x') \delta^2(x_\perp - x'_\perp) \chi_{z = \max(z, z')}
 \end{aligned}$$



Coulomb


$$\begin{aligned}
 [\dot{\Phi}_C(x), \Phi_C(x')] &= [\dot{V}_C^0(x), V_C^0(x')] \dot{\phi}(x) \dot{\phi}(x') \\
 &= i \frac{G m^2}{|x - x'|} \dot{\phi}(x) \dot{\phi}(x')
 \end{aligned}$$



AdS₃

How much can we focus?


Conical defect



$$\tilde{T}_{tt} = 0$$

$$\tilde{T}_{tt} = \frac{1}{(\varphi')^2} \left(T_{tt} + \frac{c}{12\pi\ell^2} \{ \varphi; \varphi \} \right)$$

$$\{ \varphi; \varphi \} = \frac{\varphi'''}{\varphi'} - \frac{3}{2} \left(\frac{\varphi''}{\varphi'} \right)^2$$



Zero displacement

$$\Delta\phi < 2\alpha$$

- $\alpha \rightarrow \alpha_{min}, E \rightarrow \infty$
- Minimize \bar{E} at $\alpha > 0$
 $\sim -\log \alpha$
- $M = -1 + \epsilon, \bar{E} = \bar{E}_0 + \mathcal{O}(\epsilon^2)$