

Title: PSI 2015/2016 Standard Model - Lecture 15

Date: Jan 22, 2016 10:15 AM

URL: <http://pirsa.org/16010024>

Abstract:

Problems with the SM

Problems with the SM

- (1) Neutrino mass: ν 's do have a small but nonzero mass ($\lesssim \text{eV}$)
Why are ν 's so much lighter than other fermions in SM? How do they get a mass?

$$\mathcal{L}_{\text{Yukawa}} = - Y_{ij}^{(\nu)} \bar{\nu}_R^i L_L^j \epsilon H + \text{h.c.}$$

Problems with the SM

- (1) Neutrino mass: ν 's do have a small but nonzero mass ($\lesssim \text{eV}$)
Why are ν 's so much lighter than other fermions in SM? How do they get a mass?

$$\begin{aligned}\mathcal{L}_{\text{Yukawa}} &= - Y_{ij}^{(\nu)} \bar{\nu}_R^i L_L^j \epsilon H + \text{h.c.} \\ &= - \frac{Y_{ij}^{(\nu)}}{\sqrt{2}} \bar{\nu}_R^i \nu_L^j \left(1 + \frac{h}{v}\right)\end{aligned}$$

Introduce RH ν fields
 $\nu_R^i \sim (1, 1, 0)$

Problems with the SM

(1) Neutrino mass: ν 's do have a small but nonzero mass ($\lesssim \text{eV}$)

Why are ν 's so much lighter than other fermions in SM? How do they get a mass?

$$\begin{aligned} \mathcal{L}_{\text{Yukawa}} &= - Y_{ij}^{(\nu)} \bar{\nu}_R^i L_L^{jT} \epsilon H + \text{h.c.} \\ &= - \frac{Y_{ij}^{(\nu)}}{\sqrt{2}} \bar{\nu}_R^i \nu_L^j \left(1 + \frac{h}{v}\right) \end{aligned}$$

Introduce RH ν fields
 $\nu_R^i \sim (1, 1, 0)$

Diagonalize mass matrix

$$\nu_{L,R}^i = (U_{\nu_{L,R}})_{ij} \nu_{L,R}^{'j}$$

Diagonalize mass matrix

$$\nu_{L,R}^i = (U_{\nu_{L,R}})_{ij} \nu_{L,R}^{'j}$$

$$\mathcal{L}_{\text{Yukawa}} = - m_i^{(\nu)} (\bar{\nu}_R^{'i} \nu_L^{'i} + \text{h.c.}) \left(1 + \frac{h}{v}\right)$$

New CC interaction is no longer diagonal:

$$\mathcal{L}_{\text{CC}} = - \bar{\nu}_L^i \gamma^\mu \nu_L^i$$

Diagonalize mass matrix

$$\nu_{L,R}^i = (U_{\nu_{L,R}})_{ij} \nu_{L,R}^{'j}$$

$$\mathcal{L}_{\text{Yukawa}} = - m_i^{(\nu)} (\bar{\nu}_R^{'i} \nu_L^{'i} + \text{h.c.}) \left(1 + \frac{h}{v}\right)$$

New CC interaction is no longer diagonal:

$$\mathcal{L}_{\text{CC}} = -\frac{g}{\sqrt{2}} \bar{\nu}_L^i \gamma^\mu e_L^i$$

Diagonalize mass matrix

$$\nu_{L,R}^i = (U_{\nu_{L,R}})_{ij} \nu_{L,R}^{'j}$$

$$\mathcal{L}_{\text{Yukawa}} = - m_i^{(\nu)} (\bar{\nu}_R^{'i} \nu_L^{'i} + \text{h.c.}) \left(1 + \frac{h}{v}\right)$$

New CC interaction is no longer diagonal:

$$\mathcal{L}_{\text{CC}} = -\frac{g}{\sqrt{2}} \bar{\nu}_L^i W^+ e_L^i + \text{h.c.} =$$

matrix

$$L_R = \left(U_{\nu L R} \right)_{ij} \nu_{L R}^{\prime j} \\ (\bar{\nu}_R^{\prime i} \nu_L^{\prime i} + h.c.) \left(1 + \frac{h}{v} \right)$$

ation is no longer diagonal:

$$\bar{\nu}_L^i W^+ e_L^i + h.c. = - \frac{g}{\sqrt{2}} \bar{\nu}_L^{\prime i} (U_{\nu L}^*)_{ij} (U_{e L})_{jk} W^+ e_L^{\prime k} + h.c.$$

matrix

$$L_R = (U_{VL,R})_{ij} \nu'_{L,R}{}^j$$

$$(\bar{\nu}'_R{}^i \nu'_L{}^i + \text{h.c.}) \left(1 + \frac{h}{v}\right)$$

ation is no longer diagonal:

$$\bar{\nu}_L^i W^+ e_L^i + \text{h.c.} = -\frac{g}{\sqrt{2}} \bar{\nu}_L^i \underbrace{(U_{\nu L}^*)_{ij} (U_{eL})_{jk}}_{U^+} W^+ e_L^k + \text{h.c.}$$

$$\nu_{L,R}^i = (U_{\nu_{L,R}})_{ij} \nu_{L,R}^{'j}$$

$$\mathcal{L}_{\text{Yukawa}} = - m_i^{(\nu)} (\bar{\nu}_R^{'i} \nu_L^{'i} + \text{h.c.}) \left(1 + \frac{h}{v}\right)$$

New CC interaction is no longer diagonal:

$$\mathcal{L}_{\text{CC}} = -\frac{g}{\sqrt{2}} \bar{\nu}_L^i W^+ e_L^i + \text{h.c.} = -\frac{g}{\sqrt{2}} \bar{\nu}_L^i \underbrace{(U_{\nu_L}^*)_{ij} (U_{e_L})_{jk}}_{U^+}$$

Mixing matrix for lepton $U = U_{e_L}^\dagger U_{\nu_L} = \text{PMNS matrix}$

Mixing matrix for lepton

$$U = U_{eL}^\dagger U_{\nu L} = \text{PMNS matrix}$$

We can write down a Majorana mass term for ν_R

Mixing matrix for lepton $U = U_{eL}^\dagger U_{\nu L} = \text{PMNS matrix}$

We can write down a Majorana mass term for ν_R^i

$$\mathcal{L}_{\text{Majorana}} = -\frac{1}{2} M_{ij} \left(\bar{\nu}_R^i \nu_R^{c j} + \bar{\nu}_R^{i c} \nu_R^j \right)$$

$C =$ charge conjugation

$$\nu_R^c = (\nu_R)^c = (P_R \nu)^c$$

$$\mathcal{L}_{\nu \text{ mass}} = -m (\bar{\nu}_R \nu_L + \bar{\nu}_L \nu_R) - \frac{1}{2} M (\bar{\nu}_R \nu_R^c + \bar{\nu}_R^c \nu_R)$$

$$m = \text{Dirac mass} = \frac{y_\nu}{\sqrt{2}}$$

$$\mathcal{L}_{\nu \text{ mass}} = -m (\bar{\nu}_R \nu_L + \bar{\nu}_L \nu_R) - \frac{1}{2} M (\bar{\nu}_R \nu_R^c + \bar{\nu}_R^c \nu_R)$$

$$m = \text{Dirac mass} = \frac{y_\nu}{\sqrt{2}}$$

$$\text{Note: } \bar{\nu}_R \nu_L = \bar{\nu}_L^c \nu_R^c, \quad \bar{\nu}_L \nu_R = \bar{\nu}_R^c \nu_L^c$$

$$\mathcal{L}_{\nu \text{ mass}} = -m(\bar{\nu}_R \nu_L + \bar{\nu}_L \nu_R) - \frac{1}{2} M (\bar{\nu}_R \nu_R^c + \bar{\nu}_R^c \nu_R)$$

$$m = \text{Dirac mass} = \frac{y \nu}{\sqrt{2}}$$

$$\text{Note: } \bar{\nu}_R \nu_L = \bar{\nu}_L^c \nu_R^c, \quad \bar{\nu}_L \nu_R = \bar{\nu}_R^c \nu_L^c$$

$$\mathcal{L}_{\nu \text{ mass}} = -(\bar{\nu}_L^c, \bar{\nu}_R) \begin{pmatrix} 0 & m \\ m & M \end{pmatrix} \begin{pmatrix} \nu_L \\ \nu_R^c \end{pmatrix} + \text{h.c.}$$

Note: $\bar{\nu}_R \nu_L = \bar{\nu}_L^c \nu_R^c$, $\bar{\nu}_L \nu_R = \bar{\nu}_R^c \nu_L^c$

$$\mathcal{L}_{\text{mass}} = - (\bar{\nu}_L^c, \bar{\nu}_R) \begin{pmatrix} 0 & m \\ m & M \end{pmatrix} \begin{pmatrix} \nu_L \\ \nu_R^c \end{pmatrix} + \text{h.c.}$$

$$\Psi = \begin{pmatrix} \nu_R \\ \eta^+ \end{pmatrix} \quad \Psi^c = \begin{pmatrix} \eta \\ \frac{1}{2}^+ \end{pmatrix}$$

$$\Psi_R = \begin{pmatrix} 0 \\ \eta^+ \end{pmatrix} \quad \Psi_R^c = (\Psi_R)^c = \begin{pmatrix} \eta \\ 0 \end{pmatrix}$$

Compute eigenvalue & eigenvectors (mass eigen states)

Assume $M \gg m$.

$$\nu \approx \nu_L$$

$$N \approx \nu_R$$

$$\mathcal{L}_{\text{mass}} = -\frac{1}{2} m_\nu \bar{\nu}^c \nu - \frac{1}{2} m_N \bar{N}^c N$$

$$\mathcal{L}_{\text{mass}} = -m(\bar{\nu}_R \nu_L + \bar{\nu}_L \nu_R) - \frac{1}{2} M (\bar{\nu}_R \nu_R^c + \bar{\nu}_R^c \nu_R)$$

$$m = \text{Dirac mass} = \frac{y_\nu}{\sqrt{2}}$$

$$\text{Note: } \bar{\nu}_R \nu_L = \bar{\nu}_L^c \nu_R^c, \quad \bar{\nu}_L \nu_R = \bar{\nu}_R^c \nu_L^c$$

$$\mathcal{L}_{\text{mass}} = -(\bar{\nu}_L^c, \bar{\nu}_R) \begin{pmatrix} 0 & m \\ m & M \end{pmatrix} \begin{pmatrix} \nu_L \\ \nu_R^c \end{pmatrix} + \text{h.c.}$$

$$\Psi = \begin{pmatrix} \nu_R \\ \eta^+ \end{pmatrix}$$

$$\Psi^c = \begin{pmatrix} \eta \\ \bar{\eta}^+ \end{pmatrix}$$

$$\Psi_R = \begin{pmatrix} 0 \\ \eta^+ \end{pmatrix} \quad \Psi_R^c = (\Psi_R)^c = \begin{pmatrix} \eta \\ 0 \end{pmatrix}$$

$$\nu_R = \begin{pmatrix} 0 \\ \eta^+ \end{pmatrix} \quad \nu_L = \begin{pmatrix} \xi \\ 0 \end{pmatrix}$$

Compute eigenvalue & eigenvectors (mass eigenstates)

Compute eigenvalue & eigenvectors (mass eigen states)

Assume $M \gg m$.

$$\nu \approx \nu_L \quad m_\nu \approx \frac{m^2}{M}$$

$$N \approx \nu_R \quad m_N \approx M$$

$$\mathcal{L}_{\text{mass}} = -\frac{1}{2} m_\nu \bar{\nu}^c \nu - \frac{1}{2} m_N \bar{N}^c N$$

See-saw mechanism:

Neutrinos are light because ν_μ have a
very large Majorana mass.

$$y_V \sim 1 \text{ GeV}$$

$$M \sim 10^9 \text{ GeV}$$

$$m_\nu \sim 1 \text{ eV}$$

See-saw mechanism:

Neutrinos are light because ν_R have a
very large Majorana mass.

$$y_V \sim 1 \text{ GeV}$$

$$y \sim 10^{-2}$$

$$M \sim 10^9 \text{ GeV}$$

$$m_\nu \sim 1 \text{ eV}$$

See-saw mechanism:

Neutrinos are light because ν_μ have a
very large Majorana mass.

$$y_V \sim 1 \text{ GeV}$$


$$y \sim 10^{-2}$$

$$M \sim 10^9 \text{ GeV}$$

$$m_\nu \sim 1 \text{ eV}$$

For Dirac neutrinos.

$$y \lesssim 10^{-12}$$



(2) What is the origin of flavor?
why are there 3 generations?
why are the CKM & PMNS matrices so different?

$$|V_{CKM}| = \begin{pmatrix} 0.97 & 0.23 & 0.004 \\ 0.23 & 0.97 & 0.04 \\ 0.009 & 0.04 & 0.999 \end{pmatrix}$$

ent?

closely aligned with 1.

Why are the CKM & PMNS matrices so different?

$$|V_{\text{CKM}}| = \begin{pmatrix} 0.97 & 0.23 & 0.004 \\ 0.23 & 0.97 & 0.04 \\ 0.009 & 0.04 & 0.999 \end{pmatrix}$$

$$|U_{\text{PMNS}}| = \begin{pmatrix} 0.82 & 0.54 & 0.15 \\ 0.23 & 0.70 & 0.62 \\ 0.44 & 0.45 & 0.77 \end{pmatrix}$$

the CKM & PMNS matrices so different:

$$|V_{CKM}| = \begin{pmatrix} 0.97 & 0.23 & 0.004 \\ 0.23 & 0.97 & 0.04 \\ 0.009 & 0.04 & 0.999 \end{pmatrix}$$

closely aligned with $\mathbb{1}$.

$$|U_{PMNS}| = \begin{pmatrix} 0.82 & 0.54 & 0.15 \\ 0.23 & 0.70 & 0.62 \\ 0.44 & 0.45 & 0.77 \end{pmatrix}$$

"anarchy"

(3) Why are there 3 gauge group?

Why does $SU(2)_L$ violate P ?

Can quarks & leptons be unified in a single representation?

Why do fermions have seemingly ad hoc hypercharge assignments?

Can quarks & leptons be unified in a single representation?

Why do fermions have seemingly ad hoc hypercharge assignments?

- Pati-Salam model

$$SU(4) \times SU(2)_L \times SU(2)_R$$

Can quarks & leptons be unified in a single representation?

Why do fermions have seemingly ad hoc hypercharge assignments?

- Pati-Salam model

Restores parity.

Quarks & lepton are unified.

$$SU(4) \times SU(2)_L \times SU(2)_R$$

$$\begin{aligned}\psi_L &\sim (4, 2, 1) \\ &= \begin{pmatrix} u_L^r & d_L^r \\ u_L^g & d_L^g \\ u_L^b & d_L^b \\ \nu_L & e_L \end{pmatrix}\end{aligned}$$

Can quarks & leptons be unified in a single representation?
Why do fermions have seemingly ad hoc hypercharge assignments?

Pati-Salam model

$$SU(4) \times SU(2)_L \times SU(2)_R$$

Restores parity.

Quarks & lepton are unified.

$$\begin{aligned}\psi_L &\sim (4, 2, 1) \\ &= \begin{pmatrix} u_L^r & d_L^r \\ u_L^g & d_L^g \\ u_L^b & d_L^b \\ \nu_L & e_L \end{pmatrix}\end{aligned}$$

$$\psi_R \sim (4, 1, 2)$$

Can quarks & leptons be unified in a single representation?

Why do fermions have seemingly ad hoc hypercharge assignments?

Pati-Salam model

$$SU(4) \times SU(2)_L \times SU(2)_R \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$$

Restores parity.

Quarks & lepton are unified.

$$\begin{aligned}\psi_L &\sim (4, 2, 1) \\ &= \begin{pmatrix} u_L^r & d_L^r \\ u_L^g & d_L^g \\ u_L^b & d_L^b \\ \nu_L & e_L \end{pmatrix}\end{aligned}$$

$$\psi_R \sim (4, 1, 2)$$

Can quarks & leptons be unified in a single representation?

Why do fermions have seemingly ad hoc hypercharge assignments?

Pati-Salam model

$$SU(4) \times SU(2)_L \times SU(2)_R \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$$

Restores parity.

Quarks & lepton are unified.

Explains hypercharge.

(combination of $SU(4) \times SU(2)_R$ generators)

$$\begin{aligned} \Psi_L &\sim (4, 2, 1) \\ &= \begin{pmatrix} u_L^r & d_L^r \\ u_L^g & d_L^g \\ u_L^b & d_L^b \\ \nu_L & e_L \end{pmatrix} \end{aligned}$$

$$\Psi_R \sim (4, 1, 2)$$

(combination of $SU(4) \times SU(2)$ generators) $\begin{pmatrix} u_L & d_L \\ \nu_L & e_L \end{pmatrix}$

- Georgi-Glashow $SU(5)$ model. (Grand Unified theory (GUT))

$$SU(5) \rightarrow SU(3) \times SU(2) \times U(1)$$

SM fermions fit into two representations.

$$\bar{5} = \begin{array}{|c|} \hline \\ \hline \\ \hline \\ \hline \\ \hline \end{array}$$

(combination of $SU(4) \times SU(2)$ generators) $\begin{pmatrix} u_L & d_L \\ \nu_L & e_L \end{pmatrix}$

• Georgi-Glashow $SU(5)$ model. (Grand Unified theory (GUT))

$$SU(5) \rightarrow SU(3) \times SU(2) \times U(1)$$

SM fermions fit into two representations.

$$\bar{5} = \begin{pmatrix} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{pmatrix} = \begin{pmatrix} (d_R^c)^{\alpha=r,g,b} \\ \text{---} \\ \text{---} \\ L_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \end{pmatrix} \quad \left. \begin{array}{l} \} 3 \text{ colors} \\ \} \text{Lepton doublet} \end{array} \right\}$$

(combination of $SU(4)$ and $SU(2)$ generators) $\begin{pmatrix} u_L & d_L \\ \nu_L & e_L \end{pmatrix}$

• Georgi-Glashow $SU(5)$ model. (Grand Unified theory (GUT))

$$SU(5) \rightarrow SU(3) \times SU(2) \times U(1)$$

SM fermions fit into two representations.

$$\bar{5} = \begin{bmatrix} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{bmatrix} = \begin{pmatrix} (d_R^c)_{\alpha=r,g,b} \\ \text{---} \\ \text{---} \\ L_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \end{pmatrix} \begin{matrix} \} 3 \text{ colors} \\ \\ \} \text{Lepton doublet} \end{matrix}$$

$$10 = \begin{bmatrix} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{bmatrix}$$

ν_L, e_L

Unified theory (GUT)

$\left\{ \begin{array}{l} 3 \text{ colors} \end{array} \right.$

$\left\{ \begin{array}{l} \text{Lepton doublet} \end{array} \right.$

Georgi $SO(10)$

Fit all SM fermions + ν_R in a single rep. of $SO(10)$.

ν_L, e_L

Unified theory (GUT)

} 3 colors

} Lepton doublet

Georgi $SO(10)$

Fit all SM fermions + ν_R in a single rep. of $SO(10)$. 16 dim rep.

2 quarks (u, d) $\times$ 2 chiralities $\times$ 3 colors

+ 2 leptons (e, ν) $\times$ 2 chiralities = 16.


ν_L, e_L

Unified theory (GUT)

} 3 colors

} Lepton doublet

Georgi $SO(10) \begin{matrix} \rightarrow SU(5) \rightarrow SU(3) \times SU(2) \times U(1) \\ \rightarrow P.S. \rightarrow \end{matrix}$
Fit all SM fermions + ν_R in a single rep. of $SO(10)$. 16 dim rep.
2 quarks (u, d) $\times$ 2 chiralities $\times$ 3 colors
+ 2 leptons (e, ν) $\times$ 2 chiralities = 16.



complicated

Higgs sector

Simple

fermion/gauge sector

→ complicated

completely


Higgs sector

SM

→ complicated

Simple

fermion/gauge sector



(combination of $SU(4) \times SU(2)_R$ generators) $\begin{pmatrix} \bar{\nu}_L & e_L \end{pmatrix}$

• Georgi-Glashow $SU(5)$ model. (Grand Unified theory)

$$SU(5) \rightarrow SU(3) \times SU(2) \times U(1)_Y$$

SM fermions fit into two representations.

$$\bar{5} = \begin{pmatrix} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{pmatrix} = \begin{pmatrix} (d_R^c)^{\alpha=r,g,b} \\ \text{---} \\ \text{---} \\ L_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \end{pmatrix} \quad \left. \begin{array}{l} \} 3 \text{ colors} \\ \} \text{Lepton doublet} \end{array} \right\}$$

$$10 = \begin{pmatrix} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{pmatrix}$$

$$1 = \nu_R$$

(combination of $su(4)$ & $su(2)_R$ generators)

String theory

(2) What is the origin of flavor?

Why are there 3 generations?

Why are the CKM & PMNS matrices so different?

$$|V_{CKM}| = \begin{pmatrix} 0.97 & 0.23 & 0.004 \\ 0.23 & 0.97 & 0.04 \\ 0.009 & 0.04 & 0.999 \end{pmatrix}$$

$$|U_{PMNS}| = \begin{pmatrix} 0.82 & 0.54 & 0.15 \\ 0.23 & 0.70 & 0.62 \\ 0.44 & 0.45 & 0.77 \end{pmatrix}$$

$$I = V_R$$

Standard Model Problems: The Dark Matter Puzzle

Sean Tulin

PSI lecture Fri Jan 22



Luminous mass $\neq$ Gravitational mass

**Total mass of what we see
(observed stars and gas)**

**Total mass inferred by how stars
and galaxies move under gravity**

Dark matter proposed by Fritz Zwicky in 1933, but not taken seriously.



Zwicky

*"dunkle
Materie"*

The dark matter issue became a convincing problem after work by Vera Rubin in 1970s.



Rubin

Zwicky's argument for dark matter




Coma cluster (*Sloan Digital Sky Survey*)

Zwicky's argument for dark matter

Galaxies are moving too fast and not enough mass to bind them together

The galaxies should just fly apart!



Coma cluster (*Sloan Digital Sky Survey*)

Zwicky's argument for dark matter


There is extra matter: **gas**
X-ray emission from gas

But *still* too little matter

Mass budget for Coma cluster
Stars and gas = 15% of total mass
85% of total mass = unseen mass
(dark matter)



Rubin's argument for dark matter




NGC 1232 (ESO)


$$v(r)^2 = GM(r)/r$$

v = circular velocity
G = Newton's constant
M = total mass enclosed within radius **r**


Stars are rotating too fast in galaxies



No dark matter




Dark matter
Matches observations



Ingo Berg



Andromeda galaxy



Constant velocity implies $dM/dr = \text{constant}$.


Much more mass in outer halo than the inner halo.

Seeing dark matter through its gravity




Seeing dark matter through its gravity


Cluster 0024+1654 (Hubble)



Mass map inferred by gravitational lensing
Tyson et al (1998)




**All visible galaxies are tiny objects
inside of much bigger, more
massive halos of dark matter**



Dark matter governs the structure of visible matter

**Halo stabilizes thin disk galaxies we observe
Otherwise they are unstable to perturbations**

Is dark matter made from baryons?




Very faint stars, gas, or planets that we cannot detect?

Light isotope abundances depend on the baryon density at $t \sim 1$ min (way before any stars have formed).

**Having baryons be all the dark matter is excluded by measured abundance of deuterium
No Standard Model particle can be dark matter**

Timeline of the Universe 13.8 billion years old




What is the Universe made of?


Energy/mass budget (using $E=mc^2$):

- SM** {
- 0.005% light
 - $\sim$ 0.005% neutrinos
 - 5% atoms (stars, planets, interstellar gas and dust)
- ?** {
- 25% dark matter
 - 70% dark energy

Dark Universe is 95% of the Universe.
Its nature is unknown.





How does dark matter form the structure of the Universe?




Cosmic microwave background shows the initial fluctuations of dark matter before structure formed ($\delta\rho/\rho \sim 10^{-5}$)

plasma of electrons and protons




Evolve the Universe in time

Dark matter N-body simulations



Dark matter N-body simulations



The pattern of visible matter (galaxies) traces the pattern of dark matter

Dark matter has 5x more mass. Visible matter is attracted to where the dark matter is.

The pattern of visible matter (galaxies) traces the pattern of dark matter

Dark matter has 5x more mass. Visible matter is attracted to where the dark matter is.

Dark matter predictions:

“Tiny” dark matter clumps

Dark matter filaments

Giant dark matter nodes

Voids with little dark matter

Observed Universe:


Individual galaxies with DM halos

Galaxies in filaments

Clusters of many galaxies

Voids with few galaxies

Fly through a billion light years in our local galactic neighborhood



Dark matter is a particle physics problem

Everything we know about dark matter comes from its gravitational influence in the Universe

To discover its particle physics properties, we need to know its non-gravitational interactions.

What is the Lagrangian?


What are the symmetries/forces?

LHC event: $pp \rightarrow ZZ + \text{hadrons}$

CMS Experiment at the LHC, CERN

Data recorded: 2011-May-25 08:00:19.229673 GMT(10:00:19 CEST)

Run / Event: 165633 / 394010457



© CERN 2011. All rights reserved.



If one Z went to dark matter, the second Z would appear as a transverse momentum imbalance.

Dark matter at the LHC

For example: Dark matter might be a new very heavy neutrino ($m \sim 1000$ GeV).

It can be produced in high energy collisions via the weak force.

Direct detection




Dark matter and dark forces


SM interactions with dark matter could be very suppressed.

But dark matter could have its own forces.

Feynman diagram for dark matter particles interacting via a dark force




Cuspy halo with no interactions



**Halo with interactions
("core")**

Cored halos leave their imprint on rotation curves



Most dark matter dominated galaxies have cores.
One possible explanation: Interactions among dark matter particles.

Conclusions (?)