Title: PSI 2015/2016 Quantum Information - Lecture 11

Date: Mar 07, 2016 11:30 AM

URL: http://pirsa.org/16030058

Abstract:


Gsuper
$U_+ = I - 2|\omega\rangle\langle\omega|$
$V = 2|\psi\rangle\langle\psi| - I$
$= H U_{0_1} H$


1) Physics Simulation Nielsen Chuang
Open Q Systems
2) Quantum Error Correction
ICLM
3) Quantum Error Correction
4) QKD Nielsen Chuang
5) Private Q. Channel


Grover
N
ω
θ→
|ω⟩
$U_f = I - 2|\omega\rangle\langle\omega|$
$V = 2|\psi\rangle\langle\psi| - I$
$(= H U_{0\perp} H$
$|0\cdots0\rangle \to -|\omega\rangle$
$|x_1 \cdots x_n\rangle \to (-1)^{x_1 \cdots x_n}|x_1 \cdots x_n\rangle$
$\omega = 7$
$|\psi\rangle = \frac{1}{2\sqrt{2}}(|1\rangle + |2\rangle + |3\rangle + |4\rangle + |5\rangle - |6\rangle + |7\rangle - |8\rangle)$
$U_f|\psi\rangle =$
$V U_f|\psi\rangle = (2|\psi\rangle\langle\psi| - \mathbb{1}) U_f|\psi\rangle$
$= 2\left(\frac{1}{2\sqrt{2}}\right)^2 \frac{6-1}{2\sqrt{2}}|\psi\rangle - \frac{1}{2}($ $)$
$= \frac{3}{4\sqrt{2}}|\psi\rangle - \frac{1}{2\sqrt{2}}($ $)$
$\frac{1}{4\sqrt{2}}$ $\frac{5}{4\sqrt{2}}$
Physics Simulat


Physics Simulation
$-i\frac{d|\psi\rangle}{dt} = H|\psi\rangle$
$|\psi_t\rangle = U|\psi\rangle = e^{-i\int H\,dt}|\psi_0\rangle$
$= e^{-i\Delta t H/\hbar}\, e^{-i\Delta t H/\hbar}\, e^{-i\Delta t H/\hbar} \cdots |\psi\rangle$


$\frac{5}{\sqrt{2}}$
$H_5 = \frac{Z_1 Z_2 Z_3}{Z_1 Z_3}$
$Z_1 Z_2 \mathbb{1}_3 \xrightarrow{R_2(Y, 90°)} Z \times \mathbb{1} \xrightarrow{e^{-\frac{\pi}{4} Z_1 Z_3}}$
$e^{-\frac{\pi}{4} Z_1 Z_3} Z \times \mathbb{1} \; e^{+\frac{\pi}{4} Z_2 Z_3}$
$\frac{1}{\sqrt{2}}(\mathbb{1} - Z_2 Z_3) Z_1 X_2 \mathbb{1}_3 (\quad + \quad)$
$Z_1 Y_2 Z_3$


$H_5 = \frac{Z_1 Z_2 Z_3}{Z_1 Z_3}$
$Z_1 Z_2 \mathbb{1}_3 \xrightarrow{R_2(Y,90°)} Z X \mathbb{1}$
$Z_1 Y_2 Z_3$
$\frac{1}{\sqrt{2}}(\mathbb{1} - Z_2 Z_3) Z_1 X_2 \mathbb{1}_3$
$Z_1 Y_2 Z_3$


$H_s = \frac{Z_1Z_2Z_3}{Z_1Z_3}$
$Z_1Z_2\mathbb{1}_3 \xrightarrow{R_2(Y,90°)} ZX\mathbb{1} \xrightarrow{e^{-\frac{\pi}{4}Z_2Z_3}} Z_1Y_2Z_3 \xrightarrow{e^{-\frac{\pi}{4}X_2}} Z_1Z_2Z_3$
$e^{-\frac{\pi}{4}Z_2Z_3}\, ZX\mathbb{1}\, e^{+\frac{\pi}{4}Z_2Z_3}$
$\frac{1}{\sqrt{2}}(\mathbb{1} - Z_2Z_3)Z_1X_2\mathbb{1}_3$
$Z_1Y_2Z_3$
$|\psi_0\rangle$
$|\psi\rangle$
$e^{-i\Delta t H/\hbar}$
Δt
$e^{-i\Delta t Z_1Z_2Z_3} = e^{-\frac{\pi}{4}X_2} e^{-\frac{\pi}{4}Z_2Z_3} e^{-\frac{\pi}{4}Y_2} e^{-i\Delta t Z_1Z_2} e^{+\frac{\pi}{4}Y_2} e^{+\frac{\pi}{4}Z_2Z_3} e^{+\frac{\pi}{4}X_2}$


Fermion simulations
$[X,Y] = 2iZ$
$\{a_i, a_j\} = a_i a_j + a_j a_i = 0$
$\{a_i, a_j^\dagger\} = \delta_{ij}$
$a_i|1\rangle = |0\rangle$
$a_i|0\rangle = 0$
$a_i^\dagger|0\rangle = |1\rangle$
$a_i^\dagger|1\rangle = 0$
$a_1 = \sigma_-^1$ $a_1^\dagger = \sigma_+^1$ $\sigma = X + iY$
$a_2 = \sigma_z^1 \sigma_-^2$ $a_2^\dagger = \sigma_z^1 \sigma_+^2$
$a_3 = \sigma_z^1 \sigma_z^2 \sigma_-^3$ $a_3^\dagger = \sigma_z^1 \sigma_z^2 \sigma_+^3$


Fermion simulations
[X,Y] = 2iZ
{a_i, a_j} = a_i a_j + a_j a_i = 0
{a_i, a_j^†} = δ_ij
a_1|1⟩ = |0⟩
a_1|0⟩ = 0
a_1^†|0⟩ = |1⟩
a_1^†|1⟩ = 0
a_1 = σ_1^1 σ_1^+ = σ_1^1 σ = ½(X+iY)
a_2 = σ_z^1 σ_-^2 a_2^† = σ_z^1 σ_+^2
a_3 = σ_z^1 σ_z^2 σ_-^3 a_3^† = σ_z^1 σ_z^2 σ_+^3
Q' = X+iY =
(0 1; 1 0)(0 1; -1 0)
(2 0; 0 2)


$= \frac{3}{4\sqrt{2}}|4\rangle - \frac{1}{2\sqrt{2}}(\quad)$
$\frac{1}{\sqrt{2}} \quad \frac{5}{4\sqrt{2}}$
$Y] = 2iZ$
$a_1 = \sigma_1^- \quad a_1^+ = \sigma_1^+ \quad \sigma_- = \frac{1}{2}(X + iY)$
$a_2 = \sigma_z^1 \sigma_-^2 \quad a_2^+ = \sigma_z^1 \sigma_+^2$
$a_3 = \sigma_z^1 \sigma_z^2 \sigma_-^3 \quad a_3^+ = \sigma_z^1 \sigma_z^2 \sigma_+^3$
$\sigma_-^1 = X + iY =$
$\{a_1, a_2\} = a_1 a_2 + a_2 a_1$
$= \sigma_-^1 \sigma_z^1 \sigma_-^2 + \sigma_z^1 \sigma_-^1 \sigma_-^2$
$= (\sigma_-^1 \sigma_z^1 + \sigma_z^1 \sigma_-^1)\sigma_-^2$
$= 0$

$U_+ = I - 2|w><w|$

$V = 2|+><+|I - I$

$(= H U_{01} H$

↳ $|0\cdots0> \to -|\infty>$

$|x_1 x_2 x_3> \to |x_1 x_2>$

$= 2\left(\frac{1}{\sqrt{2}}\right)^2 \frac{6-1}{2\sqrt{2}}|+> - \frac{1}{2}($ $)$

$= \frac{3}{4\sqrt{2}}|+> - \frac{1}{2\sqrt{2}}($ $)$

$\frac{1}{4\sqrt{2}}$ $\frac{5}{4\sqrt{2}}$

Fermion simulations

$[X,Y] = 2iZ$

$\{a_0, a_j\} = a_0 a_j + a_j a_0 = 0$

$\{a_i, a_j^\dagger\} = \delta_{ij}$

$a_1|1> = |0>$

$a_1|0> = 0$

$a_1^\dagger|0> = |1>$

$a_1^\dagger|1> = 0$

$a_1 = \sigma_-^1 \quad a_1^\dagger = \sigma_+^1 \quad \mathbb{C} = \frac{1}{2}(X+iY)$

$a_2 = \sigma_z^1 \sigma_-^2 \quad a_2^2 = \sigma_z^1 \sigma_+^2$

$a_3 = \sigma_z^1 \sigma_z^2 \sigma_-^3 \quad a_3^2 = \sigma_z^1 \sigma_z^2 \sigma_+^3$

$\mathbb{C}' = X+iY =$

$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$

$\begin{pmatrix} 2i \\ 2 \end{pmatrix}$

$\{a_1, a_2\} = a_1 a_2 + a_2 a_1$

$= \sigma_-^1 \sigma_z^1 \sigma_-^2 + \sigma_z^1 \sigma_-^1 \sigma_-^2$

$= (\sigma_-^1 \sigma_z^1 + \sigma_z^1 \sigma_-^1)\sigma_-^2$

$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$

$\begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$

$= 0$

ZZ_3

$X_2 \to \frac{1}{-1}\frac{Z}{Z}\frac{Y}{3}$

$\downarrow$

$Z_1 Z_2 X_3$

Quantum operations

A q. operation is a map sends density matrices to density matrices

$\mathcal{E}$:
- be linear $\mathcal{E}(p\rho + q\sigma) = p\mathcal{E}(\rho) + q\mathcal{E}(\sigma)$
- preserve the trace $tr[\mathcal{E}(\rho)] = tr[\rho]$
- preserve positivity $\rho \geq 0$ $\mathcal{E}(\rho) \geq 0$
- completely positive $(\mathcal{E} \otimes I)\rho \geq 0$

Quantum operations

A q. operation is a map sends density matrices to density matrices

$\mathcal{E}$:
- be linear $\mathcal{E}(p\rho + q\sigma) = p\mathcal{E}(\rho) + q\mathcal{E}(\sigma)$
- preserve the trace $tr[\mathcal{E}(\rho)] = tr[\rho]$
- preserve positivity $\rho \geq 0 \quad \mathcal{E}(\rho) \geq 0$
- completely positive $(\mathcal{E} \otimes I)\rho \geq 0$

$$\begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \longrightarrow \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

- Preserve positivity $\rho \geq 0 \quad \mathcal{E}(\rho) \geq 0$

i) Depolarizing channel

$$\rho = (1-p)\rho + \frac{p}{3}\left[X\rho X + Y\rho Y + Z\rho Z\right]$$





ii) Dephasing

$$\rho = (1-p)\rho + Z\rho Z$$

$$\begin{pmatrix} \alpha\alpha^* & \alpha\beta^* e^{-t/T_2} \\ \alpha^*\beta e^{-t/T_2} & \beta\beta^* \end{pmatrix}$$


① $\mathcal{E}$ is CPTP
② $\mathcal{E}(\rho) = \sum_i A_i \rho A_i^\dagger$ $\sum A_i^\dagger A_i = I$
③ $\exists U$ $\mathcal{E}(\rho) = \mathrm{tr}_B [U(\rho \otimes |0\rangle\langle 0|)U^\dagger]$
Amplitude damping (relaxation to ground state)
$\rho = A_0 \rho A_0^\dagger + A_1 \rho A_1^\dagger$
$A_0 = \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{1-\gamma} \end{pmatrix}$
$A_1 = \begin{pmatrix} 0 & \sqrt{\gamma} \\ 0 & 0 \end{pmatrix}$
$|0\rangle_e|0\rangle_f \to |0\rangle|0\rangle$
$|1\rangle_e|0\rangle_f$
$\to \sqrt{1-\gamma}\,|1\rangle_e|0\rangle_f + \sqrt{\gamma}\,|0\rangle_e|1\rangle_f$