

Title: Gravitational Waves: The Birth of a New Area of Astronomy

Date: Oct 20, 2016 01:00 PM

URL: <http://pirsa.org/16100061>

Abstract: <p>On September 14th and December 26th, 2015, the Advanced LIGO detectors observed two gravitational wave signals, each from the merger of stellar-mass black holes. These two observations have given us the first glimpse in to the population of stellar mass black holes. In this talk I will discuss these first detections of gravitational waves including the non-detection of gravitational waves from the merger of binary neutron star and neutron star black holes systems. I will also describe the LIGO interferometers, their current state and the future of this exciting new field of gravitational-wave astronomy.</p>

Gravitational wave propagation

$$h(t) = Ae^{i(2\pi ft - \mathbf{k} \cdot \mathbf{r})}$$

Gravitational wave strength is given in strain:

$$h \sim \delta L / L$$


Typical strains from astrophysical sources are very small by the time they arrive at Earth:


$$h \sim 10^{-21}$$

GW luminosity of recent BBH detections are huge. Peak luminosity:

$$L_{\text{GW}} \sim 10^{56} \text{ erg/s}$$

(GRB luminosity $\sim 10^{49-52}$ erg/s)





Proxima Centauri

4.2 light years

Imagine measuring this distance to a precision of **ten microns**

LIGO

Laser Interferometer Gravitational-wave Observatory

LIGO-Hanford



LIGO-Livingston




LIGO

Wave Observatory

LIGO-Livingston




The design




PRL 116, 131103 (2016)

5

Sensitivity: sources of noise




Sensitivity: past, present and future



PRL 116, 131103 (2016)

9

GW150914



PRL 116, 061102 (2016)

- Observed on September 14th, 2015 at 09:40:45 UTC
- First observed in LIGO-Livingston then 7ms later at LIGO-Hanford
- Over 0.2 seconds the signal increases in frequency and amplitude over ~8 cycles from 35Hz to peak amplitude at 150 Hz

11

How do we know 2 black holes collided?

Chirp mass:

$$\mathcal{M} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}} = \frac{c^3}{G} \left[\frac{5}{96} \pi^{-8/3} f^{-11/3} \dot{f} \right]^{3/5}$$

$\mathcal{M} \sim 30 M_\odot$ implies total mass $M \gtrsim 70 M_\odot$ (detector frame)

Components must reach an orbital frequency of 75 Hz without touching. Equal Newtonian point masses orbiting at this frequency would be ~ 350 km apart

Black holes are the only known objects compact enough to be this close without touching

The big announcement...

PRL 116, 061102 (2016)

Selected for a Viewpoint in *Physics*
PHYSICAL REVIEW LETTERS

week ending
12 FEBRUARY 2016



Observation of Gravitational Waves from a Binary Black Hole Merger

B. P. Abbott *et al.**

(LIGO Scientific Collaboration and Virgo Collaboration)

(Received 21 January 2016; published 11 February 2016)

On September 14, 2015 at 09:50:45 UTC the two detectors of the Laser Interferometer Gravitational-Wave Observatory simultaneously observed a transient gravitational-wave signal. The signal sweeps upwards in frequency from 35 to 250 Hz with a peak gravitational-wave strain of 1.0×10^{-21} . It matches the waveform predicted by general relativity for the inspiral and merger of a pair of black holes and the ringdown of the resulting single black hole. The signal was observed with a matched-filter signal-to-noise ratio of 24 and a false alarm rate estimated to be less than 1 event per 203 000 years, equivalent to a significance greater than 5.1σ . The source lies at a luminosity distance of 410_{-180}^{+160} Mpc corresponding to a redshift $z = 0.09_{-0.04}^{+0.03}$. In the source frame, the initial black hole masses are $36_{-4}^{+5} M_{\odot}$ and $29_{-4}^{+4} M_{\odot}$, and the final black hole mass is $62_{-4}^{+4} M_{\odot}$, with $3.0_{-0.5}^{+0.5} M_{\odot} c^2$ radiated in gravitational waves. All uncertainties define 90% credible intervals. These observations demonstrate the existence of binary stellar-mass black hole systems. This is the first direct detection of gravitational waves and the first observation of a binary black hole merger.

DOI: 10.1103/PhysRevLett.116.061102

I. INTRODUCTION

In 1916, the year after the final formulation of the field


The discovery of the binary pulsar system PSR B1913+16 by Hulse and Taylor [20] and subsequent observations of its energy loss by Taylor and Weisberg [21] demonstrated

The day after the announcement...



Image Credit: Calum Torrie et al.

GW151226



PRL 116, 241103 (2016)


17

To detect signals from compact-object binaries, we construct a bank template waveforms and matched-filter the data

$$\rho = \frac{\langle s|h \rangle}{\sqrt{\langle h|h \rangle}}$$

$$\langle a|b \rangle = 4\text{Re} \int_{f_{\text{low}}}^{f_{\text{high}}} \frac{\tilde{a}(f)\tilde{b}(f)}{S_n(f)} df$$

Apply additional waveform-consistency tests to separate signal from noise



Allen, ..., DAB, et al. Phys Rev D 85 122006 (2012)


Babak, ..., DAB, et al. Phys Rev D 87 024033 (2013)

Usman, ..., DAB, et al. arXiv:1508.02357
Capano, et al. arXiv:1602.03509

Abbott, ..., DAB, et al. arXiv:1602.03839
DAB, et al., Phys. Rev. D 86 084017 (2012)

Calculating Significance


- Determined by rate at which detector noise produces an event with a detection statistic value equal to or higher than the candidate event
- Background set of data is created from coincident data from multiple detectors
- Slide the timestamps of one detector's data by many multiples of 0.1s and computing a new set of coincident events



Usman et al., arXiv: 1508.02357 (2015)


22

Results from the first observing run



Parameters of the BBH systems


Posterior probability densities of the masses, spins and distance to the three events



Abbott et al. arXiv: 1606.04856 (2016)


25

Electromagnetic Follow-Up




Timeline of observations of GW150914, separated by band and relative to the time of the gravitational wave event

Localisation



Sky localization depends on:

- the location and orientation of the detectors
- time delay between signal arrival at spatially separated sites




Abbott et al. arXiv: 1606.04856 (2016)

out BBH
 depend
 which we
 y well
 different
 ons
 merger
 ange
 1


Searching for BNS and NS-BH systems

During O1 we looking for gravitational waves from binary neutron star (BNS) and neutron star - black hole (NS-BH) systems



- O1 90% upper limit NS-BH rate compared to other published rates
- Dark blue assumes 1.4-5 M_⊙ and light blue 1.4-10 M_⊙.
- Constrain the merger rate of NS-BH systems with BH at least 5 M_⊙ to be less than 3,600 Gpc⁻³yr⁻¹ (assuming isotropic distribution of component spins)
- O2 and O3 BNS ranges are assumed to be 1-1.9 and 1.9-2.7 times larger than O1

Advanced LIGO's sensitivity was at the upper end of that predicted for the first observing run



Abbott et al. Living Reviews in Relativity **19**, 1 (2016)