Title: Tensor networks and the renormalization group

Date: Nov 09, 2016 02:00 PM

URL: http://pirsa.org/16110045

Abstract: Tensor networks offer an efficient representation of many-body wave-functions in an exponentially large Hilbert space by exploiting the area law of ground state quantum entanglement. I will start with a gentle introduction to the tensor network formalism. Then I will describe its application to realizing Wilson's renormalization group directly on quantum lattice models (e.g. quantum spin chains), with emphasis on the RG fixed points corresponding to conformal field theories. We will see how to define both global and local scale transformations on the lattice in such a way that, intriguingly, conformal invariance can be tested (and conformal data extracted) right at the UV cut-off scale.

Perimeter Nov 9th 2016

COLLOQUIUM

Tensor networks and the renormalization group





Guifre Vidal

PERIMETER INSTITUTE FOR THEORETICAL PHYSICS

TENSOR NETWORKS INITIATIVE

SIMONS FOUNDATION

NSERC

compute canada | calcul canada

Outline

1 – Entanglement and tensor networks

$S(A) \sim |\partial A|$

area law

2 – Application: The renormalization group





Quantum Mechanics

1920-1930

Niels Bohr



Albert Einstein

Wolfgang Pauli

Paul Dirac

$$i\hbar \frac{\partial}{\partial t}\Psi = \hat{H}\Psi$$

Schrodinger equation

Erwin Schrodinger

Enrico Fermi

Werner Heisenberg

Richard Feynman

There is a problem…

Paul Dirac

"The fundamental laws necessary for the mathematical treatment of a large part of physics and the whole of chemistry are thus completely known,

and the difficulty lies only in the fact that application of these laws leads to equations that are too complex to be solved."

number of spins		Hilbert space dimension	
$N = 1$	●	$2^1 = 2$	$\|\Psi\rangle = c_1\|\uparrow\rangle + c_2\|\downarrow\rangle$
$N = 2$	● ●	$2^2 = 4$	$\|\Psi\rangle = c_1\|\uparrow\uparrow\rangle + c_2\|\uparrow\downarrow\rangle + c_3\|\downarrow\uparrow\rangle + c_4\|\downarrow\downarrow\rangle$
$N = 3$	● ● ●	$2^3 = 8$	$\|\Psi\rangle = c_1\|\uparrow\uparrow\uparrow\rangle + c_2\|\uparrow\uparrow\downarrow\rangle + c_3\|\uparrow\downarrow\uparrow\rangle + c_4\|\uparrow\downarrow\downarrow\rangle + c_5\|\downarrow\uparrow\uparrow\rangle + c_6\|\downarrow\uparrow\downarrow\rangle + c_7\|\downarrow\downarrow\uparrow\rangle + c_8\|\downarrow\downarrow\downarrow\rangle$
$N = 4$	● ● ● ●	$2^4 = 16$	$\|\Psi\rangle = c_1\|\uparrow\uparrow\uparrow\uparrow\rangle + c_2\|\uparrow\uparrow\uparrow\downarrow\rangle + c_3\|\uparrow\uparrow\downarrow\uparrow\rangle + c_4\|\uparrow\uparrow\downarrow\downarrow\rangle + c_5\|\uparrow\downarrow\uparrow\uparrow\rangle + c_6\|\uparrow\downarrow\uparrow\downarrow\rangle + c_7\|\uparrow\downarrow\downarrow\uparrow\rangle + c_8\|\uparrow\downarrow\downarrow\downarrow\rangle + c_9\|\downarrow\uparrow\uparrow\uparrow\rangle + c_{10}\|\downarrow\uparrow\uparrow\downarrow\rangle + c_{11}\|\downarrow\uparrow\downarrow\uparrow\rangle + c_{12}\|\downarrow\uparrow\downarrow\downarrow\rangle + c_{13}\|\downarrow\downarrow\uparrow\uparrow\rangle + c_{14}\|\downarrow\downarrow\uparrow\downarrow\rangle + c_{15}\|\downarrow\downarrow\downarrow\uparrow\rangle + c_{16}\|\downarrow\downarrow\downarrow\downarrow\rangle$
N	● ● ● ··· ●	$dim = 2^N$	



example: $N = 16 \times 16 = 256$ spins

$\dim = 2^{256} \approx 10^{77}$ complex numbers

10GB laptop

10^9 complex numbers

$N = 32$ spins

petabyte (10^{15} bytes) supercomputer

10^{14} complex numbers

$N = 48$ spins



# of atoms in

human being

10^{28}

earth

10^{50}

solar system

10^{57}



milky way

10^{69}

observable universe

10^{80}



Entanglement

$\mathcal{H}^A \otimes \mathcal{H}^B$

quantum spin chain

$$|\Psi^{AB}\rangle \stackrel{?}{=} |\psi^A\rangle \otimes |\varphi^B\rangle$$

Def.: product state $|\Psi^{AB}\rangle = |\psi^A\rangle \otimes |\varphi^B\rangle$

example: $|\uparrow\uparrow \cdots \uparrow\uparrow^A\rangle \otimes |\uparrow\uparrow \cdots \uparrow\uparrow^B\rangle$

Def.: entangled state $|\Psi^{AB}\rangle \neq |\psi^A\rangle \otimes |\varphi^B\rangle$

example:

$\frac{1}{\sqrt{2}}|\uparrow\uparrow \cdots \uparrow\uparrow^A\rangle \otimes |\uparrow\uparrow \cdots \uparrow\uparrow^B\rangle + \frac{1}{\sqrt{2}}|\downarrow\downarrow \cdots \downarrow\downarrow^A\rangle \otimes |\downarrow\downarrow \cdots \downarrow\downarrow^B\rangle$

How much entanglement does $|\Psi^{AB}\rangle$ a state have?

Def.: entanglement entropy $\quad S(\Psi^{AB}) \equiv -Tr\,(\rho^A \log_2 \rho^A)$

$$\rho^A \equiv tr_B\,(|\Psi^{AB}\rangle\langle\Psi^{AB}|)$$

example: $\frac{1}{\sqrt{2}}|\uparrow\uparrow\cdots\uparrow\uparrow^A\rangle \otimes |\uparrow\uparrow\cdots\uparrow\uparrow^B\rangle + \frac{1}{\sqrt{2}}|\downarrow\downarrow\cdots\downarrow\downarrow^A\rangle \otimes |\downarrow\downarrow\cdots\downarrow\downarrow^B\rangle$

$$\rho^A = \frac{1}{2}|\uparrow\uparrow\cdots\uparrow\uparrow^A\rangle\langle\uparrow\uparrow\cdots\uparrow\uparrow^A| + \frac{1}{2}|\downarrow\downarrow\cdots\downarrow\downarrow^A\rangle\langle\downarrow\downarrow\cdots\downarrow\downarrow^A|$$

$S(\Psi^{AB}) = 1$ — 1 unit of entanglement

example: $|\uparrow\uparrow\cdots\uparrow\uparrow^A\rangle \otimes |\uparrow\uparrow\cdots\uparrow\uparrow^B\rangle$ $\quad S(\Psi^{AB}) = 0$ — 0 units of entanglement

$$\rho^A = |\uparrow\uparrow\cdots\uparrow\uparrow^A\rangle\langle\uparrow\uparrow\cdots\uparrow\uparrow^A|$$

Scaling of entanglement

quantum spin chain



generic state $S(L) \approx L$

product state $S(L) = 0$

$|\uparrow\uparrow \cdots \uparrow\uparrow\rangle$

ground state of
local Hamiltonians

massive/gapped $S(L) = const.$

critical/gapless $S(L) = \log(L)$



Ground states of local Hamiltonians obey an entanglement area law





area law
for entanglement entropy

$$S(A) \sim |\partial A| \sim L^{D-1}$$

instead of
volume law

$$S(A) \sim |A| \sim L^{D}$$

sometimes,
logarithmic corrections

$$S(A) \sim L^{D-1} \log(L)$$

Ground states of local Hamiltonians are special/non-generic states



Tensor network state: variational many-body wavefunction

many-body wave-function

$$|\Psi\rangle = \sum_{i_1, i_2, \cdots, i_N} \Psi_{i_1 i_2 \cdots i_N} \, |i_1 i_2 \cdots i_N\rangle$$



graphical notation





$$T_{ij} = \sum_k R_{ik} S_{kj} \qquad a = \vec{y}^{\dagger} \cdot M \cdot \vec{x} \qquad tr(ABCD)$$

$$|\Psi\rangle = \sum_{i_1, i_2, \cdots, i_N} \Psi_{i_1 i_2 \cdots i_N} |i_1 i_2 \cdots i_N\rangle$$

1 tensor
of size 2^N



=



$O(N)$ tensors
of size χ^p independent of N

2^N
parameters
inefficient

→

$O(N)$
parameters
efficient





Scaling of entanglement

matrix product state

(MPS)



multi-scale entanglement renormalization ansatz

(MERA)



network connectivity:

$S(A) \approx const$

area law!

just as ground state of massive/gapped 1d system

$S(A) \approx \log L$

logarithmic correction!

just as ground state of critical/gapless 1d system

CURRENT APPLICATIONS

tensor network =
sparse data structure



PERIMETER PI INSTITUTE FOR THEORETICAL PHYSICS

ENGLISH | FR

ABOUT RESEARCH TRAINING OUTREACH PEOPLE EVENTS VIDEO LIBRARY NEWS SUPPORT PI

Home » Research » Research Initiatives » Tensor Networks Initiative

RESEARCH

RESEARCH AREAS

RESEARCH INITIATIVES

TENSOR NETWORKS

EHT

FROM DISCRETUUM TO CONTINUUM

TENSOR NETWORKS INITIATIVE

FROM ENTANGLED QUANTUM MATTER TO EMERGENT SPACE TIME

Tensor networks have in recent years emerged as a powerful tool to describe strongly entangled quantum many-body systems. Applications range from the study of collective phenomena in condensed matter to providing a discrete realization of the holographic principle in quantum gravity.

TENSOR NETWORKS INITIATIVE

Ash Milsted
PI postdoc

1 – Entanglement and tensor networks

$S(A) \sim |\partial A|$

area law

2 – Application: The renormalization group





Part II

Application: the Renormalization Group

A – The renormalization group and its fixed points



B – Global scale transformation/invariance



Glen Evenbly
University of
Sherbrooke

C – Local scale transformation/invariance



The Renormalization Group

Hamiltonian

$$H[\vec{k}] \longrightarrow H[\vec{k}(s)]$$

scale

coupling constants

$$\vec{k} = (k_1, k_2, k_3, \cdots)$$

Ising model

$$H[J, \lambda] = J \sum_i \sigma_i^x \sigma_j^x + \lambda \sum_i \sigma_i^z$$

$$H[k_1, k_2, 0, 0, \cdots] = k_1 \sum_i \sigma_i^x \sigma_j^x + k_2 \sum_i \sigma_i^z$$

Leo Kadanoff

Kenneth Wilson

RG flow in the space of Hamiltonians



RG flow in the space of Hamiltonians



Leo Kadanoff





block spin

+ some rule: majority vote, etc

Change of scale?

coarse-graining transformation

Kenneth Wilson

$$Z = \int_{|p|\leq\Lambda} \mathcal{D}\phi e^{-H[\phi,\vec{k}]}$$

$$e^{-H[\phi,\vec{k}']} = \int_{\Lambda' \leq |p| \leq \Lambda} \mathcal{D}\phi e^{-H[\phi,\vec{k}]}$$

$$Z = \int_{|p|\leq\Lambda'} \mathcal{D}\phi e^{-H[\phi,\vec{k}']}$$



exact renormalization group equation (ERGE)

MERA as a Renormalization Group transformation



MERA defines an RG flow in the space of wave-functions



$$|\Psi\rangle \to |\Psi'\rangle \to |\Psi''\rangle \to \cdots$$




$\mathcal{L}'$
$\mathcal{L}$
w isometry
u disentangler
$\mathbb{V}^{\otimes N/2}$
$\uparrow W^\dagger$
$\mathbb{V}^{\otimes N}$


$\mathcal{O}$
$\mathcal{O}' = W^\dagger \mathcal{O} W$
scale transf.
local operator on $\mathcal{L}$
local operator on $\mathcal{L}'$


u
$u^\dagger$
=
w
$w^\dagger$
=


$\mathcal{L}'$
$\mathcal{L}$
w isometry
u disentangler
$\mathbb{V}^{\otimes N/2}$
$\uparrow W^\dagger$
$\mathbb{V}^{\otimes N}$


$\mathcal{O}$
$\mathcal{O}' = W^\dagger \mathcal{O} W$
scale transf.
local operator on $\mathcal{L}$
local operator on $\mathcal{L}'$


u
$u^\dagger$
=
w
$w^\dagger$
=


=

MERA defines an RG flow in the space of wave-functions:

$$|\Psi\rangle \to |\Psi'\rangle \to |\Psi''\rangle \to \cdots$$





What is a proper RG transformation on the lattice?

why use this?



and not use this? (closer to Kadanoff spin-blocking)



MERA



$S(A) \approx \log L$

tree tensor network (TTN)



$S(A) \approx const.$

What is a proper RG transformation on the lattice?

MERA RG flow seems to be qualitatively correct
= consistent with expectation from QFT

e.g.: when adding a mass m in a free particle theory, mass grows with coarse-grainining transformation as expected:





Is the MERA RG flow quantitatively correct?

Let us analyze critical fixed-points

In the continuum, conformal field theory (CFT)

critical RG fixed-point = Conformal field theory (CFT)

1) scaling operators $\phi_\alpha(x)$

complex plane



$\phi_\alpha(0)$

scale transformation

re-scaling

$z \to e^{\lambda} z$



$e^{\Delta_\alpha \lambda} \phi_\alpha(0)$

scaling dimension

Lorentz boost

(= rotation in Euclidean time)

rotation

$z \to e^{i\theta} z$



$e^{i s_\alpha \theta} \phi_\alpha(0)$

conformal spin

example: CFT for critical Ising model

$$H = \sum_i \sigma_i^x \sigma_j^x + \sum_i \sigma_i^z$$

primary fields

spin

$\{\mathbb{I}, \sigma, \varepsilon\}$

identity

energy density

scaling dimensions and conformal spins

$(\Delta_{\mathbb{I}}, s_{\mathbb{I}}) = (0,0)$

$(\Delta_\sigma, s_\sigma) = (1/8,0)$

$(\Delta_\varepsilon, s_\varepsilon) = (1,0)$

Ising CFT

spin

primary fields

$\{\mathbb{I}, \sigma, \varepsilon\}$

identity

energy density

global scale/rotation invariance + locality = local scale/rotation invariance (conformal group)

2) conformal towers

under conformal transformations, scaling operators mix



conformal towers = irreducible representations of the conformal group

primary field = highest weight state

3) Operator product expansion (OPE)



$$\phi_\alpha(x)\phi_\beta(y) \sim \sum_\gamma C_{\alpha\beta\gamma}(x-y)\phi_\gamma(x)$$

CFT completely specified by conformal data:

- central charge c
- list of primary fields $\{\phi_\alpha\}$ with their
  - scaling dimensions $\Delta_\alpha \equiv h_\alpha + \bar{h}_\alpha$
  - conformal spin $s_\alpha \equiv h_\alpha - \bar{h}_\alpha$
  - OPE coefficients $C_{\alpha\beta\gamma}$

example: Ising CFT

central charge $c = \frac{1}{2}$

primary fields $\{\mathbb{I}, \sigma, \varepsilon\}$ (spin)

- $\mathbb{I}$: identity
- ε: energy density

scaling dimensions and conformal spins

$(\Delta_{\mathbb{I}}, s_{\mathbb{I}}) = (0,0)$

$(\Delta_\sigma, s_\sigma) = (1/8,0)$

$(\Delta_\varepsilon, s_\varepsilon) = (1,0)$

OPE

$C_{\epsilon\sigma\sigma} = \frac{1}{2}$

With notion of scale/RG transformation on the lattice,



$\equiv W$

define notion of (discrete) **scale invariance**

Def.: $|\Psi\rangle$ is scale invariant

iff $\quad W|\Psi\rangle = |\Psi\rangle$

example: scale-invariant wavefunction



$|\Psi\rangle$

Intriguing...

scale invariance with a length scale (UV cut-off)

Is this a useful notion of scale transformation / scale invariance ?

We can access the correct *conformal data* of underlying CFT from the lattice



$= 2^{-\Delta_\alpha}$

scaling operators

- scaling dimensions $\quad \Delta_\alpha \equiv h_\alpha + \bar{h}_\alpha$
- OPE coefficients $C_{\alpha\beta\gamma}$

e.g. critical Ising model

critical Hamiltonian

$$H = \sum_i \sigma_i^x \sigma_j^x + \sum_i \sigma_i^z$$



1) variational optimization

$\min_{u,w} \langle \Psi | H | \Psi \rangle$

(approx. an hour on your laptop)

ansatz $|\Psi\rangle \leftrightarrow (u, w)$

isometry

disentangler

2) diagonalize linear map





scaling dimensions

$(\Delta_{\mathbb{I}} = 0)$

$\Delta_\sigma \approx 0.124997$

$\Delta_\varepsilon \approx 0.99993$

$\Delta_\mu \approx 0.125002$

$\Delta_\psi \approx 0.500001$

$\Delta_{\bar{\psi}} \approx 0.500001$

OPE

$C_{\epsilon\sigma\sigma} = \frac{1}{2}$

$C_{\epsilon\mu\mu} = -\frac{1}{2}$

$C_{\epsilon\psi\bar{\psi}} = i$

$C_{\epsilon\bar{\psi}\psi} = -i$

$C_{\psi\mu\sigma} = \frac{e^{-\frac{i\pi}{4}}}{\sqrt{2}}$

$C_{\bar{\psi}\mu\sigma} = \frac{e^{\frac{i\pi}{4}}}{\sqrt{2}}$

$(\pm 6 \times 10^{-4})$

What is a proper RG transformation on the lattice?



MERA RG flow seems to be qualitatively correct
= consistent with expectation from QFT



At a critical RG fixed-point, MERA accurately reproduces the universal data of the underlying CFT

critical fixed point = CFT

RG transformation $\sim$ global scale transformation



- scaling dimensions Δ_α
- OPE coefficients $C_{\alpha\beta\gamma}$

Glen Evenbly

- conformal spins s_α?
- which scaling operators ϕ_α are primary operators?

answer will come from defining/studying local scale transformations on the lattice

another **CFT fact** :



A CFT on a circle of perimeter L has energy and momentum spectra given by

$$E_\alpha = \frac{2\pi}{L}\left(\Delta_\alpha - \frac{c}{12}\right)$$

$$p_\alpha = \frac{2\pi}{L}\, s_\alpha$$

continuum → discrete





$$H|\Psi_\alpha\rangle = E_\alpha|\Psi_\alpha\rangle \qquad H|\Psi_\alpha\rangle = E_\alpha|\Psi_\alpha\rangle$$

continuous time translations

$$P|\Psi_\alpha\rangle = p_\alpha|\Psi_\alpha\rangle \qquad T|\Psi_\alpha\rangle = e^{ip_\alpha}|\Psi_\alpha\rangle$$

discrete space translations

DVRC

John Cardy 1989

$$E_\alpha = \frac{2\pi}{L}\left(\Delta_\alpha - \frac{c}{12}\right)$$

$$p_\alpha = \frac{2\pi}{L}\, s_\alpha$$



example:
(coarse-grained)
Ising model
$L = 8$ sites

- scaling dimensions Δ_α
- conformal spins s_α

great! However,

- primary operators ϕ_α?
- conformal towers ?

global scale transformation



local scale transformation v1.0 (incorrect)



local scale transformation v2.0 (correct)





tensor network equalities

example 1: generic local scale transformation

















conformal tower of identity

$\mathbb{I}$

conformal tower of spin

σ

conformal tower of energy density

ε

Cardy 1989



Ashley 2016





$$p_\alpha = \frac{2\pi}{L} s_\alpha$$

$$E_\alpha = \frac{2\pi}{L}\left(\Delta_\alpha - \frac{c}{12}\right)$$

use tensor networks to build and apply
local scale transformations on the lattice



example 2: local scale transformation corresponding to global conformal generators





Virasoro L_0 $L_{\pm 1}$ $L_{\pm 2}$ $L_{\pm 3}$ $\cdots$

global conformal

$$q_1 \equiv (l_1 - \bar{l}_{-1}) + (l_{-1} - \bar{l}_1)$$
$$\sim cos\left(\frac{2\pi}{L}x\right)\partial_x$$

$$Q_1 \equiv (L_1 - \bar{L}_{-1}) + (L_{-1} - \bar{L}_1)$$

$e^{iQ_1} \sim$ mixing within global conformal towers





identify quasi-primaries and their global conformal towers

Summary

1 - Entanglement and tensor networks

$S(A) \sim |\partial A|$

area law

2 - Application: The renormalization group





What else?

Euclidean path integrals
(Lagrangian formalism)

Continuous
tensor networks
for QFTs

Tensor network holography: AdS/CFT