Title: Testing early Universe physics with upcoming observations

Date: Mar 28, 2017 11:00 AM

URL: http://pirsa.org/17030093

Abstract: Cosmology has seen great progress thanks to precision measurements and is bound to greatly benefit from upcoming Large Scale Structure and Cosmic Microwave Background data. I will point out a number of interesting directions. In particular, I discuss how the microphysics of inflation may be tested in galaxy surveys through â€œfossilâ€• signatures originating from squeezed primordial correlations. I further elaborate on the constraining power of CMB spectral distortions on small-scale cosmological fluctuations and on particle decays in the very early Universe in relation to reheating. I also describe some of the possible constraints on inflation and reheating from future B-mode observations.

Outline:

ED, M. Fasiello, M. Kamionkowski - *2015*
ED, M. Fasiello, D. Jeong, M. Kamionkowski - *2014*

Fossils from inflation :
signatures of long-short mode couplings in galaxy surveys

J. Chluba, **ED**, M.A. Amin, M. Kamionkowski - *2016*
ED, R. Emami - *2016*
ED, L.M. Krauss, J. Chluba - *2015*
R. Emami, **ED**, J. Chluba, M. Kamionkowski - *2015*

Spectral distortions of the CMB:
testing small scales primordial perturbations and BSM physics

ED, M. Fasiello, T. Fujita - *2016*
M. Biagetti, **ED**, M. Fasiello, M. Peloso - *2014*

Inflation and primordial gravitational waves:
production mechanisms and r-H correspondence

A glance at cosmic history …



3

Inflation

- Era of *accelerated (nearly exponential) expansion* in the primordial Universe : $a(t) \sim e^{Ht}$



Hubble radius: $\sim 1/H \sim$ const.

Physical scales are stretched by the expansion!

- Mechanism for the generation of cosmological fluctuations

$$\phi(\vec{x}, t) = \varphi(t) + \delta\phi(\vec{x}, t)$$

Inflaton quantum fluctuations

↓

CMB perturbations
and LSS of the Universe





Inflation

- Simplest realization: single-scalar field in slow-roll (SFSR)

Accelerated expansion : $\ddot{a} > 0$

Einstein eqs. : $\dfrac{\ddot{a}}{a} = -\dfrac{1}{2\,M_P^2}\,(\rho + 3\,p)$ $\Bigg\}$ $p < -\dfrac{1}{3}\rho$

Scalar field :

$$p_\varphi = \frac{\dot{\varphi}^2}{2} - V(\varphi) \approx -V(\varphi)$$

$$\rho_\varphi = \frac{\dot{\varphi}^2}{2} + V(\varphi) \approx V(\varphi)$$

$\Bigg\}$ $p_\varphi \approx -\rho_\varphi$

Slow-roll

Einstein eqs. : $\longrightarrow$ $H^2 = \dfrac{\rho_\varphi}{3\,M_P^2} \approx \dfrac{V(\varphi)}{3\,M_P^2} \approx$ const.

$$\ddot{\varphi} + 3H\dot{\varphi} + V'(\varphi) = 0$$

Slow-roll parameters

$$\begin{cases} \epsilon \equiv -\dfrac{\dot{H}}{H^2} = \dfrac{1}{2}\left(\dfrac{M_P V_\varphi}{V}\right)^2 \ll 1 \\ \eta \equiv \dfrac{M_P^2\, V_{\varphi\varphi}}{V} \ll 1 \end{cases}$$



Predictions: power spectra (SFSR)

Scalar fluctuations : $\delta\phi \longrightarrow (\delta T/\bar{T} - \delta\rho/\bar{\rho})$ $\quad P_S \sim \frac{1}{\epsilon}\left(\frac{H}{M_P}\right)^2 k^{n_s-1}$

Tensor fluctuations / primordial gravity waves : $\quad P_T \sim \left(\frac{H}{M_P}\right)^2 k^{n_T}$

Energy-scale of inflation !

What have we learned from observations?

Scalar fluct. :

- nearly-scale invariant $n_s = 0.968 \pm 0.006$

Tensor fluct. :
tensor-to-scalar ratio
r < 0.07

BICEP2/KECK +Planck



Predictions: power spectra (SFSR)

Scalar fluctuations : $\delta\phi \longrightarrow (\delta T/\bar{T} - \delta\rho/\bar{\rho})$ $\quad P_S \sim \frac{1}{\epsilon}\left(\frac{H}{M_P}\right)^2 k^{n_s-1}$

Tensor fluctuations / primordial gravity waves : $\quad P_T \sim \left(\frac{H}{M_P}\right)^2 k^{n_T}$

Energy-scale of inflation !

What have we learned from observations?

Scalar fluct. :

- nearly-scale invariant $n_s = 0.968 \pm 0.006$

Tensor fluct. :
tensor-to-scalar ratio
r < 0.07

BICEP2/KECK +Planck



One or more active fields

How do fields interact

Any spectator field

?

Energy scale of inflation

What are the masses of the extra fields

How does inflation end

What happens at Reheating

…

Microphysics of inflation: primordial non-Gaussianity



local



$f_{\rm nl}^{\rm loc} = 0.8 \pm 5$

orthogonal



$f_{\rm nl}^{\rm orth} = -26 \pm 21$

equilateral



$f_{\rm nl}^{\rm eq} = -4 \pm 43$

$$\begin{cases} f_{\rm nl}^{\rm loc} \gtrsim \mathcal{O}(1) \longrightarrow \cancel{\text{single-clock}} \\ f_{\rm nl}^{\rm eq,\,orth} \gtrsim \mathcal{O}(1) \longrightarrow \cancel{\text{slow-roll}} \end{cases}$$

thresholds potentially at reach for upcoming galaxy surveys!

There is more:
e.g. ***intermediate shapes***



Case study: Quasi-Single-Field

tangent: inflaton
radial: massive isocurvaton

(also SUSY -motivated)





Bispectrum shape between local and equil.



$m_{iso} \to 0$

$m_{iso} \to 3\,H/2$

$$\mathcal{B}|_{squ.} \sim \frac{1}{k_1^3 k_3^3}\left(\frac{k_3}{k_1}\right)^{3/2-f(m_{iso})}$$

$$f(m_{iso}) \equiv \sqrt{\frac{9}{4} - \frac{m_{iso}^2}{H^2}}$$

bispectrum shape delivers info on mass spectrum!

[Chen, Wang, 2010]

What happens next?

- Galaxy surveys : *photometric/spectroscopic* (DESI, LSST, Euclid, SPHEREx, …) and *21cm* (CHIME, …)

  Microphysics of inflation (**primordial non-Gaussianity**, running/features of primordial power spectrum, …) → Fossils from inflation

- CMB experiments :

  Spectral distortions (PRISM, PIXIE, …)

  Testing post- and **pre-recombination** physics → A window into primordial perturbations on small scales

  B-modes (BICEP/KECK series, SPTPol, SPIDER, PIXIE, …)

  Origin of B-modes? What can we learn about the **early Universe**? → Inflation and primordial gravitational waves

‘Fossils’ : signatures from **multi-clock** inflationary dynamics



Effect of a long mode on local observables :



- Gaussian initial conditions + statistical homog./isotropy → $\langle \delta_{\vec{k}_1} \delta_{\vec{k}_2} \rangle = \delta^{(3)}(\vec{k}_1 + \vec{k}_2) P(k_1)$ diagonal 2p corr.

- Local/squeezed non-Gaussianity



→



→ $\langle \delta_{\vec{k}_1} \delta_{\vec{k}_2} \rangle_{\vec{K}} = \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{K}) \times f(\vec{k}_1, \vec{k}_2) A(K)$

off-diagonal 2p corr.!

Probing σ through off-diagonal correlation :



Azimuthal dependence:

- Scalar → no φ dependence
- Vector → $\cos\varphi$, $\sin\varphi$
- Tensor → $\cos 2\varphi$, $\sin 2\varphi$

[Jeong, Kamiomkowski, 2012]

What if σ = tensor mode of the metric:



Learning about primordial tensors
through non-Gaussian effects!

[ED, Fasiello, Jeong, Kamionkowski - 2014]
[ED, Fasiello, Kamionkowski - 2015]

Probing σ through off-diagonal correlation :



Azimuthal dependence:

- Scalar → no φ dependence
- Vector → $\cos\varphi$, $\sin\varphi$
- Tensor → $\cos 2\varphi$, $\sin 2\varphi$

[Jeong, Kamiomkowski, 2012]

What if σ = tensor mode of the metric:



Learning about primordial tensors
through non-Gaussian effects!

[ED, Fasiello, Jeong, Kamionkowski - 2014]
[ED, Fasiello, Kamionkowski - 2015]

- Optimal estimator for a single mode : inverse variance weighting

$$\widehat{\gamma_p(\vec{K})} = \sigma^2 \sum_{\vec{k}} \frac{|B_p(\vec{K},\vec{k},\vec{K}-\vec{k})/P_\gamma^p(K)|^2}{2\,V\,P^{tot}(k)P^{tot}(|\vec{K}-\vec{k}|)}\delta_{\vec{k}}\delta_{\vec{K}-\vec{k}}$$

$$\sigma^2 = \left[\sum_{\vec{k}} \frac{|B_p(\vec{K},\vec{k},\vec{K}-\vec{k})/P_\gamma(K)|^2}{2\,V\,P^{tot}(k)P^{tot}(|\vec{K}-\vec{k}|)}\right]^{-1}$$

random variable

variance

$$\{y_i, \sigma_i^2\}$$

$$\hat{y} = \frac{\sum_i (y_i/\sigma_i^2)}{\sum_j (1/\sigma_j^2)} \qquad \sigma^2 = \left(\sum_i \frac{1}{\sigma_i^2}\right)^{-1}$$

- Optimal estimator for power amplitude: stochastic GW background with $P_p(K) = A_\gamma\, P^f_\gamma(K)$



$$\widehat{A_\gamma} = \sigma_\gamma^2 \sum_{p,\vec{K}} \frac{(P^f_\gamma(K))^2}{(P^n_p(K))^2} \left(\frac{|\widehat{\gamma_p(\vec{K})}|^2}{V} - P^n_p(K) \right)$$

optimal sum over different K-modes

$$\begin{cases} \sigma_\gamma^{-2} = \sum_{p,\vec{K}} \frac{(P^f_\gamma(K))^2}{2(P^n_p(K))^2} \\ P^n_p \equiv \left[\sum_{\vec{k}} \frac{|B_p(\vec{K},\vec{k},\vec{K}-\vec{k})/P_\gamma(K)|^2}{2V P^{tot}(k) P^{tot}(|\vec{K}-\vec{k}|)} \right]^{-1} \end{cases}$$

$$\mathcal{B}_{\gamma\zeta\zeta} \simeq \beta P_\gamma P_\zeta$$

$$\sigma_\gamma \propto \frac{1}{\beta^2} \left(\frac{k_{max}}{k_{min}} \right)^{-3}$$

Which classes of models predict these signatures at observable levels?

~~single-clock ccs~~

- (n+1)-point function fixed in terms of n-point functions (soft limit for one of the modes)
- Apply if super-horizon modes freeze + standard initial conditions (rescaling of background for short modes)
- Derived from symmetries of the action (invariance under space diffs) [Maldacena 2003, Creminelli-Zaldarriaga 2004, Goldberger et al 2013, . . .]

- *Isocurvature* modes (multi-field)
- *Non-Bunch Davies* initial states [Holman - Tolley 2007, Brahma-Nelson-Shandera 2013, …]
- *Broken space diffs* (e.g. solid inflation/space-dependent background) [Endlich et al., 2013, Bartolo et al, 2015]

Fossil signatures at reach for upcoming observations (e.g. Euclid or 21cm)!

[ED, Fasiello, Jeong, Kamionkowski - 2014]
[ED, Fasiello, Kamionkowski - 2015]

Summary and prospects

- *Squeezed limit*: powerful discriminant for inflationary models (*single vs multi-clock*)
- Correlations in squeezed limit from inflation: *local observables* affected by *long modes (Anisotropic* effects / *Off-diagonal* correlations*)*
- Necessary to understand effects of long wl. modes:
  - — test existence of a long-short correlation
  - — correctly map observations to theory

Next :

- Building catalogue of fossil signatures for multi-field models
  → estimating masses/spin/interactions of primordial fields
- Lots more to do on tensor non-Gaussianities (especially timely given B-modes searches!)

A window into primordial perturbations on *small scales*

Cosmic Microwave spectrum from COBE/FIRAS



Energy/photon injections into CMB at $z \lesssim 2 \times 10^6$

↓

Distortion of the blackbody spectrum

$$\Delta I_\nu \approx \int G_{\rm th}(\nu, z') \frac{d(Q/\rho_\gamma)}{dz'} dz'$$

energy-injection rate



Deviations allowed from FIRAS:

$|\mu| \lesssim 9 \times 10^{-5}$

$|y| \lesssim 1.5 \times 10^{-5}$

[Fixsen et al., 1996]

[Chluba, 2013]

Analytic approximations for visibility functions

$$\frac{\Delta\rho_\gamma}{\rho_\gamma} \approx \left[\frac{\Delta\rho_\gamma}{\rho_\gamma}\right]_\mu + \left[\frac{\Delta\rho_\gamma}{\rho_\gamma}\right]_y$$

$$\frac{\mu}{1.4} \approx \int_0^\infty \mathcal{J}(z')\mathcal{J}_\mu(z')\frac{d(Q/\rho_\gamma)}{dz'}dz'$$

$$4\,y \approx \int_0^\infty \mathcal{J}(z')\mathcal{J}_y(z')\frac{d(Q/\rho_\gamma)}{dz'}dz'$$



$$\mathcal{J}(z) \approx e^{-[z/(2\times10^6)]^{5/2}}$$

$$\mathcal{J}_\mu(z) \approx 1 - \mathcal{J}_y(z)$$

$$\mathcal{J}_y(z) \approx \left[1 + \left(\frac{1+z}{6\times10^4}\right)^{2.58}\right]^{-1}$$

- for energy injection + no residual SDs!

Why spectral distortions matter?

CMB anisotropies:
(~20 yrs)



Spectral distortions:

COBE/FIRAS



$|\mu| \lesssim 9 \times 10^{-5}$
$|y| \lesssim 1.5 \times 10^{-5}$
[Fixsen et al., 1996]

PIXIE



$|\mu| \lesssim 2 \times 10^{-8}$
$|y| \lesssim 4 \times 10^{-9}$
[Kogut et al., 2011]

Mechanisms producing SDs include:

- Cosmological recombination
- Reionization, structure formation
- Dissipation of magnetic fields
- Decay/annihilation of particles
- Dissipation of primordial fluctuations
- …

Why spectral distortions matter?

CMB anisotropies:
(~20 yrs)



Spectral distortions:

COBE/FIRAS



$|\mu| \lesssim 9 \times 10^{-5}$

$|y| \lesssim 1.5 \times 10^{-5}$

[Fixsen et al., 1996]

PIXIE



$|\mu| \lesssim 2 \times 10^{-8}$

$|y| \lesssim 4 \times 10^{-9}$

[Kogut et al., 2011]

Mechanisms producing SDs include:

- Cosmological recombination
- Reionization, structure formation
- Dissipation of magnetic fields
- Decay/annihilation of particles
- Dissipation of primordial fluctuations
- …

Gravitinos in the early Universe

- predicted in SUGRA
- spin 3/2 super-partners of gravitons
- SUSY-breaking scale: $F \sim \sqrt{m_{3/2} M_P}$



$V(\phi)$ dominates → inflaton decays → $T_{\rm rh}$ → ordinary radiation/ matter eras

relic gravitinos quickly *diluted*

gravitinos production by:
- *rapid inflaton oscillations*
- *scatterings in the hot plasma*

Input: DVI - 1920x1080p@60Hz
Output: SDI - 1920x1080i@60Hz

Example : $G \to \gamma + \tilde{\gamma}$



$B_{[G \to \gamma + \tilde{\gamma}]} = 1$

[**ED**, L. M. Krauss, J. Chluba -2015]

Summary and prospects

- Competitive bounds from SDs on particle decays/annihilations
- Gravitinos from Reheating: constraints on SUSY/ scale of inflation

Next :

- *Residual (r-type) distortions:* more observables!

  $(10^4 \lesssim z \lesssim \text{few} \times 10^5)$

  $p = \{\mu, y, \mu_k\}$ $\begin{cases} \mu_1^{(\mathrm{DL})} \simeq 3 \times 10^{-7} \\ \mu_2^{(\mathrm{DL})} \simeq 2 \times 10^{-6} \\ \mu_3^{(\mathrm{DL})} \simeq 7 \times 10^{-6} \end{cases}$

- *Photon (vs energy) injection :*
  different treatment of thermalization problem

- *Post-recombination decays :*
  different treatment of energy exchange as plasma recombines

- *Complete particles cascade :*
  likely to strengthen current bounds

SDs from diffusion damping

Primordial fluctuations re-enter Hubble radius after inflation

Isotropization by photon diffusion



[Hu&White, 1996]

[Planck]



Effective **energy release** into CMB

[Hu et al., 1994]
[Chluba et al., 2012]

$$\frac{d(Q/\rho_\gamma)}{dz} \sim \int dk\, k^4 \mathcal{P}_\zeta(k)\, e^{-[2\,k^2/k_D^2(z)]} \approx \left(4\times 10^{-6}(1+z)^{3/2}\mathrm{Mpc}^{-1}\right)^2$$

(damping scale)

Probing scales k: $\begin{cases} [50, 10^4]\,\mathrm{Mpc}^{-1} & \text{(with } \mu \text{ distortion)} \\ [1, 50]\,\mathrm{Mpc}^{-1} & \text{(with } y \text{ distortion)} \end{cases}$

probing ~ **10** extra inflationary e-folds!

Power spectrum constraints: SFSR

$$\mathcal{P}_\zeta(k) = A_\zeta \left(\frac{k}{k_*}\right)^{n_s - 1 + \frac{1}{2}\{\mathrm{d}n_s/\mathrm{d}\ln k\}\ln(k/k_*)}$$

$$\begin{cases} n_s = 0.968 \pm 0.006 \\ \mathrm{d}n_s/\mathrm{d}\ln k = -0.006 \pm 0.007 \end{cases}$$

Planck temperature + polarization data on large angular scales (68% CL)

[Khatri, Sunyaev, Chluba, 2012]



E.g. for ns = 0.96: $\dfrac{Q^{z=5\times10^4}_{z=2\times10^6}}{\rho_\gamma} \simeq \mathcal{O}(10^{-8})$

Sensitivity for μ distortion

$$\sigma_\mu = (1/n) \times 10^{-8}$$

E.g. : n=1 ← PIXIE, n=10 ← PRISM

3 x PIXIE = *guaranteed discovery*

detection (95% CL) | exclusion of SFSR (95% CL)

[Cabass, Melchiorri, Pajer, 2016]

Power spectrum constraints: ~~SESR~~

$$\mu \approx 2.2 \int \mathcal{P}_\zeta \left[e^{-\left(\frac{k\cdot \mathrm{Mpc}}{5400}\right)^2} - e^{-\left(\frac{k\cdot \mathrm{Mpc}}{31.6}\right)^2} \right] d\ln k$$

$$y \approx 0.4 \int \mathcal{P}_\zeta(k) e^{-\left(\frac{k\cdot \mathrm{Mpc}}{31.6}\right)^2} d\ln k$$





- e.g. from particle production
- bump of amplitude $\mathcal{A}_{\zeta,i}$, localized around k_i
- intermediate distortions to remove degeneracies

[Chluba et al., 2015]

General note:

bounds from SDs competitive w.r.t.those from PBHs and UCMHs!

30

Bispectrum constraints

- Bispectrum in the squeezed limit from distortion-temperature correlation

[Pajer - Zaldarriaga, 2012]

Example: local ansatz $\mathcal{R}(\vec{x}) = r(\vec{x}) + \frac{3}{5} f_{nl}\, r^2(\vec{x})$

Long-short mode decomposition: $\mathcal{R} = \mathcal{R}_L + \mathcal{R}_s$

$$\downarrow$$

$$\mathcal{R}_s(\vec{x}) \approx r_s(\vec{x}) \left[1 + \frac{6}{5} f_{nl}\, \mathcal{R}_L(\vec{x}) \right]$$

$$\frac{\Delta T}{T} \sim \frac{\mathcal{R}}{5} \qquad \frac{\Delta d}{d} \sim \frac{\delta\langle \mathcal{R}^2 \rangle}{\langle \mathcal{R}^2 \rangle} \longrightarrow C_l^{(T,d)} \sim f_{nl}(k_d)\, C_l^{TT}$$

k_d: $(1-50)\mathrm{Mpc}^{-1}$ for y; $(50-10^4)\mathrm{Mpc}^{-1}$ for μ

k_T: $\lesssim 0.01\,\mathrm{Mpc}^{-1}$ (large angle CMB temp. fluct.)

J. Chluba, ED, M.A. Amin, M. Kamionkowski - 2016, ED, R. Emami - 2016
R. Emami, ED, J. Chluba, M. Kamionkowski - 2015

31

Scale-dependence of non-Gaussianity

$\lesssim 0.01\mathrm{Mpc}^{-1}$ k_T k_d $(50-10^4)\,\mathrm{Mpc}^{-1}$ for μ, $(1-50)\,\mathrm{Mpc}^{-1}$ for y

Smallest detectable μ-T and y-T correlations (1-σ):

- template: $B_{\mathcal{R}} \simeq \frac{12}{5} f_{nl}(k_s) P_{\mathcal{R}}(k_s) P_{\mathcal{R}}(k_L)$
- uniform f_{nl} signal on μ and y scales

$$f_{nl}^{(\mu)} \simeq 10^2 \left(\frac{\mu_{\min}}{10^{-9}}\right)\left(\frac{\langle\mu\rangle}{2\times 10^{-8}}\right)^{-1}$$
$$f_{nl}^{(y)} \simeq 10^2 \left(\frac{y_{\min}}{2\times 10^{-10}}\right)\left(\frac{\langle y\rangle}{4\times 10^{-9}}\right)^{-1}$$

smallest detectable monopoles for a given experiment

$$\langle\mu\rangle \approx \int d\log k\, \Delta^2_{\mathcal{R}}(k) W_\mu(k)$$
$$\langle y\rangle \approx \int d\log k\, \Delta^2_{\mathcal{R}}(k) W_y(k)$$

[Emami, **ED**, Chluba, Kamionkowski - 2015]

Summary and prospects

SDs from diffusion damping gives access to ~10 extra inflationary e-folds!

→ **Vast range of modes** to uncover statistics of **primordial fluctuations**

	k/Mpc	*fnl*
CMB anisot./galaxy surveys	$(10^{-4} - 1)$	bispectrum / off-diagonal correlations, …
y distortion	$(1 - 50)$	y -T correlation
μ distortion	$(50 - 10^4)$	μ -T correlation

Next :

— **More non-Gaussian *observables*, e.g. :**

- Cross-correlating distortions with *polarization*
- Including *residual* *(r-type)* distortions $(10^4 \lesssim z \lesssim \text{few} \times 10^5)$

— **Evaluating *non-primordial* distortions:**
e.g. from black-holes, (BSM/DM)particles decays/annihilations, …

Inflation and primordial *gravitational waves*

B-modes polarization



BICEP2/KECK+Planck

$$r < 0.07$$

Stage-IV CMB experiments : $\sigma(r) \sim 0.001$

35

GWs from inflation: SFSR

- Production from tensor modes of the metric

$$ds^2 = a^2(\tau)\left[-d\tau^2 + (\delta_{ij} + \gamma_{ij}(\tau, \vec{x}))dx^i dx^j\right]$$

transverse & traceless

- Primordial tensor power spectrum

$$\mathcal{P}_\gamma \sim \left(\frac{H}{M_P}\right)^2 k^{n_T}$$

energy scale of inflation

$$V_{\rm infl}^{1/4} \approx 10^{16}{\rm GeV}\,(r/0.01)^{1/4}$$

- Consistency relations

$$n_T \simeq -2\epsilon = -r/8$$

(red-spectrum)

- Lyth bound $\Delta\phi/M_P \gtrsim (r/0.01)^{1/2}$ (small vs large field)

Alternative inflationary production mechanisms

- Spectator fields with small sound speed
  [Biagetti, Fasiello, Riotto 2012, Biagetti, ED, Fasiello, Peloso 2014]

$$\left[\partial_\tau^2 + k^2 - \frac{a''}{a}\right](a\,\gamma_{ij}) = S_{ij} \qquad S_{ij} \sim \mathcal{O}(\delta\sigma^2/c_s^n)$$

source of GW
(= 0 if free-field)

Main challenge: sourcing tensors to observable level without affecting scalar sector (e.g. large nG)!

- Particle production / gauge fields:
  — Auxiliary scalars with time-varying masses : $\sum_i \chi_i^2\,(\phi - \phi_i)^2$
  [Chung et al., 2000, Senatore et al, 2011, Pearce et al, 2016]

  — Gauge fields coupled to axions : $\chi\, F\,\tilde{F}$
  [Sorbo 2011, Mukohyama et al. 2012-2014, ED-Fasiello-Fujita 2016, …]

Important : inflaton <u>not directly</u> coupled to source!

* Breaking standard r ⟷ H correspondence
* Distinct signature can arise: blue spectrum (⟶ interesting also for direct detection!); parity-violation (⟶ TB, EB)

GWs from inflation: SFSR

- Production from tensor modes of the metric

$$ds^2 = a^2(\tau)\left[-d\tau^2 + (\delta_{ij} + \gamma_{ij}(\tau, \vec{x}))dx^i dx^j\right]$$

transverse & traceless

- Primordial tensor power spectrum

$$\mathcal{P}_\gamma \sim \left(\frac{H}{M_P}\right)^2 k^{n_T}$$

energy scale of inflation

$$V_{\rm infl}^{1/4} \approx 10^{16}{\rm GeV}\,(r/0.01)^{1/4}$$

- Consistency relations

$$n_T \simeq -2\epsilon = -r/8$$

(red-spectrum)

- Lyth bound $\Delta\phi/M_P \gtrsim (r/0.01)^{1/2}$ (small vs large field)

Alternative inflationary production mechanisms

- Spectator fields with small sound speed
  [Biagetti, Fasiello, Riotto 2012, Biagetti, ED, Fasiello, Peloso 2014]

$$\left[\partial_\tau^2 + k^2 - \frac{a''}{a}\right](a\,\gamma_{ij}) = S_{ij} \qquad S_{ij} \sim \mathcal{O}(\delta\sigma^2/c_s^n)$$

source of GW
(= 0 if free-field)

Main challenge: sourcing tensors to observable level without affecting scalar sector (e.g. large nG)!

- Particle production / gauge fields:
  — Auxiliary scalars with time-varying masses : $\sum_i \chi_i^2\,(\phi - \phi_i)^2$
  [Chung et al., 2000, Senatore et al, 2011, Pearce et al, 2016]

  — Gauge fields coupled to axions : $\chi\, F\,\tilde{F}$
  [Sorbo 2011, Mukohyama et al. 2012-2014, ED-Fasiello-Fujita 2016, …]

Important : inflaton <u>not directly</u> coupled to source!

* Breaking standard r ⟷ H correspondence

* Distinct signature can arise: blue spectrum (⟶ interesting also for direct detection!); parity-violation (⟶ TB, EB)

Many possibilities for discovery …

An exciting road ahead
for cosmology!

39

Final summary

Fossils from inflation : signatures of long-short mode couplings in galaxy surveys

Spectral distortions: testing primordial perturbations on small scales and BSM physics

Inflation and primordial GWs: production mechanisms and the r-H correspondence