

Title: Positive Representations of Split Real Quantum Groups

Date: May 01, 2017 02:00 PM

URL: <http://pirsa.org/17050032>

Abstract: <p>The notion of Positive Representations is a new research program devoted to the representation theory of split real quantum groups, initiated in a joint work with Igor Frenkel. It is a generalization of the special class of representations considered by J. Teschner for  $U_q(\mathfrak{sl}(2, \mathbb{R}))$  in Liouville theory, where it exhibits a strong parallel to the finite-dimensional representation theory of compact quantum groups, but at the same time also serves some new properties that are not available in the compact case. In this talk, I will survey the recent developments and describe some of its relations to other areas of mathematics.&nbsp;</p>

## Motivation

[Drinfeld, Jimbo (1985)]

Simple Lie algebra  $\mathfrak{g} \rightsquigarrow$  quantum group  $\mathcal{U}_q(\mathfrak{g})$

- Search for solutions to Yang-Baxter equation

Finite dimensional representation theory  $\longrightarrow$  many applications!

- Knot and 3-manifolds invariant:  
Reshetikhin-Turaev's TQFT, braided tensor category
- Categorification: Khovanov homology, Nakajima's quiver variety...
- Kazhdan-Lusztig theory:  $Rep(\widehat{\mathfrak{g}}) \longleftrightarrow Rep(\mathcal{U}_q(\mathfrak{g}))$
- Lusztig's Canonical basis / Kashiwara's Crystal basis
- Many more...

## Motivation

[Drinfeld, Jimbo (1985)]

Simple Lie algebra  $\mathfrak{g} \rightsquigarrow$  quantum group  $\mathcal{U}_q(\mathfrak{g})$

- Search for solutions to Yang-Baxter equation

Finite dimensional representation theory  $\longrightarrow$  many applications!

- Knot and 3-manifolds invariant:  
Reshetikhin-Turaev's TQFT, braided tensor category
- Categorification: Khovanov homology, Nakajima's quiver variety...
- Kazhdan-Lusztig theory:  $Rep(\widehat{\mathfrak{g}}) \longleftrightarrow Rep(\mathcal{U}_q(\mathfrak{g}))$
- Lusztig's Canonical basis / Kashiwara's Crystal basis
- Many more...

## Motivation

Classical Lie theory: two important **real forms**:  $\mathfrak{g}_\mathbb{C}$  and  $\mathfrak{g}_\mathbb{R}$

$\mathfrak{g}_\mathbb{C} \longleftrightarrow$  compact groups (e.g.  $SU(n), SO(2n)$ )

- Finite dimensional representation theory is **well-behaved**
  - Highest weight representations  $V_\lambda : \lambda \in P^+$
  - Closure under tensor product:

$$V_\lambda \otimes V_\mu \simeq \bigoplus_{\nu} c_{\lambda\mu}^{\nu} V_{\nu}$$

- Peter-Weyl's Theorem:

$$\mathbb{C}[G] \simeq \bigoplus_{\lambda} V_{\lambda} \otimes V_{\lambda}^*$$

- Generalized nicely to corresponding **quantum groups**  $\mathcal{U}_q(\mathfrak{g}_\mathbb{C})$ 
  - Universal  $R$  matrix: Braiding:  $V_\lambda \otimes V_\mu \simeq V_\mu \otimes V_\lambda$   
 $\longrightarrow$  Braided Tensor Category

## Motivation

$\mathfrak{g}_{\mathbb{R}} \longleftrightarrow$  split real groups (e.g.  $SL(n, \mathbb{R}), SO(n, n)$ )

- Much more complicated [Harish-Chandra]

Ex:  $G=SL(2, \mathbb{R})$ :

- Principal series  $P_{\lambda}^{\epsilon}$ , discrete series, complementary series
- Tensor product:

$$P_{\lambda} \otimes P_{\mu} \simeq \int_{\mathbb{R}^+}^{\oplus} P_{\nu} d\mu(\nu) \bigoplus \text{(discrete part)}$$

- Peter-Weyl's Theorem:

$$L^2(SL(2, \mathbb{R})) \simeq \int_{\mathbb{R}^+}^{\oplus} P_{\lambda} \otimes P_{\lambda}^* d\mu(\lambda) \bigoplus \text{(discrete part)}$$

- Quantum group level - involving self-adjoint operators
- Largely open due to analytic difficulties and spectral properties (non-compactness, unbounded operators etc.)
- Foundation work from Physics by Faddeev, Kashaev, Volkov, .....

## Representation theory of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

Simplest case:  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

Some **special class** of representations  $\mathcal{P}_\lambda = \text{GOOD!}$

Studied by **Teschner *et al.*** from **quantum Liouville theory**:

- Parameterized by  $\lambda \in \mathbb{R}_{\geq 0}$
- Generators = **positive self-adjoint** operators on  $L^2(\mathbb{R})$
- **Integrable representations** in the sense of [**Schmüdgen (1999)**].
- **No classical limit  $q \rightarrow 1$**

## Representation theory of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

Parallel to representation theory of compact case!!

- (Continuous) Braided tensor category structure  
[Ponsot-Teschner, Bytsko-Teschner]:

$$\mathcal{P}_\alpha \otimes \mathcal{P}_\beta \simeq \int_{\mathbb{R}_+}^{\oplus} \mathcal{P}_\gamma d\mu(\gamma)$$

$$\mathcal{P}_\alpha \otimes \mathcal{P}_\beta \simeq \mathcal{P}_\beta \otimes \mathcal{P}_\alpha$$

- Harmonic analysis gives Peter-Weyl type theorem  
[Ip (2013)]:

$$L^2(SL_q^+(2, \mathbb{R})) \simeq \int_{\mathbb{R}_+}^{\oplus} \mathcal{P}_\gamma \otimes \mathcal{P}_\gamma^* d\mu(\gamma)$$

## Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

New research program started in [Frenkel-Ip]

- Generalization of Teschner's representations to higher rank
- Generators = positive self-adjoint operators on  $L^2(\mathbb{R}^N)$

Many new phenomena **not present** in compact case:

- Faddeev's modular double and quantum dilogarithm
- Langlands duality as simple analytic relations
- Theory of multiplier Hopf algebra and locally compact quantum group from  $C^*$ -algebra
- Discriminant variety for Positive Casimirs
- Connection to quantum Teichmüller theory and cluster algebra

We expect many applications of finite dimensional representation theory of  $\mathcal{U}_q(\mathfrak{g}_c)$  can be generalized to  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ .

Definition of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ 

$q = e^{\pi i b^2}$  not root of unity,  $0 < b^2 < 1$ .

## Definition [Drinfeld-Jimbo]

$\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R})) = \text{Hopf-}^* \text{ algebra } \langle E, F, K^{\pm 1} \rangle \text{ such that}$

$$KE = q^2 EK, \quad KF = q^{-2} FK, \quad [E, F] = \frac{K - K^{-1}}{q - q^{-1}}$$

*Coproduct:*

$$\Delta(K) = K \otimes K, \quad \Delta(E) = E \otimes K + 1 \otimes E, \quad \Delta(F) = F \otimes 1 + K^{-1} \otimes F$$

*Real form:*

$$K^* = K, \quad E^* = E, \quad F^* = F$$

(Also counit  $\epsilon$ , antipode  $S$ )

For higher rank, also Serre relations,  $K_i E_j = q_i^{a_{ij}} E_j K_i$  etc.

## Integrable Representations of Quantum Plane

For  $q = e^{\pi i b^2}$ , there is a **canonical representation** for the relation

$$UV = q^2 VU$$

where  $U, V$  are **positive self adjoint** operators:

$$U = e^{2\pi b x}, \quad V = e^{2\pi b p} \quad (\text{i.e. } (Vf)(x) = f(x - ib))$$

**unbounded operators** on  $L^2(\mathbb{R})$ , where  $p = \frac{1}{2\pi i} \frac{d}{dx}$

- Acting on common dense **core**:  $\mathcal{W} := \{e^{-\alpha x^2 + \beta x} P(x)\}_{\text{Re}(\alpha) > 0, \beta \in \mathbb{C}}$
- **Integrable representation [Schmüdgen]**:

$$U^{is} V^{it} = q^{-2st} V^{it} U^{is}, \quad \forall s, t \in \mathbb{R}$$

as unitary operators.

## Faddeev's Modular Double

Recall  $UV = q^2VU$ ,

$$q = e^{\pi i b^2}, \quad U = e^{2\pi b x}, \quad V = e^{2\pi b p}$$

Define

$$\tilde{q} := e^{\pi i b^{-2}}, \quad \tilde{U} := U^{\frac{1}{b^2}}, \quad \tilde{V} := V^{\frac{1}{b^2}}$$

i.e. replacing  $b$  by  $b^{-1}$ . Then

- $\tilde{U}\tilde{V} = \tilde{q}^2\tilde{V}\tilde{U}$
- $\{U, V\}$  commute (weakly) with  $\{\tilde{U}, \tilde{V}\}$

Together  $\langle U, V, \tilde{U}, \tilde{V} \rangle$  generates the **Modular Double**.

- **Faddeev's Idea:** should extend to quantum group level.
- Liouville theory:  $q = e^{\pi i b^2} \longleftrightarrow c = 1 + 6(b + b^{-1})^2$

## Ponsot-Teschner's representation

- $b \longleftrightarrow b^{-1}$  gives  $\{\tilde{E}, \tilde{F}, \tilde{K}\}$  a representation of  $\mathcal{U}_{\tilde{q}}(\mathfrak{sl}(2, \mathbb{R}))$ .
- $\{E, F, K\}$  commute (weakly) with  $\{\tilde{E}, \tilde{F}, \tilde{K}\}$
- Define

$$\mathbf{e} := \left( \frac{i}{q - q^{-1}} \right)^{-1} E, \quad \mathbf{f} := \left( \frac{i}{q - q^{-1}} \right)^{-1} F,$$

we have

$$\mathbf{e}^{\frac{1}{b^2}} = \tilde{\mathbf{e}}, \quad \mathbf{f}^{\frac{1}{b^2}} = \tilde{\mathbf{f}}, \quad K^{\frac{1}{b^2}} = \tilde{K},$$

called the "transcendental relations".

Hence  $\mathcal{P}_\lambda$  is a representation of the **Modular Double**

$$\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(2, \mathbb{R})) := \mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R})) \otimes \mathcal{U}_{\tilde{q}}(\mathfrak{sl}(2, \mathbb{R})).$$

## Properties

Existence of universal  $R$  operator ( $K =: q^H$ ):

Theorem [Bytsko-Teschner (2003)]

$$R = q^{\frac{H \otimes H}{4}} g_b(\mathbf{e} \otimes \mathbf{f}) q^{\frac{H \otimes H}{4}}$$

$$R\Delta = \Delta^{op}R$$

- $R$  is a unitary operator on  $L^2(\mathbb{R}) \otimes L^2(\mathbb{R})$ .
- $R$  is invariant under  $b \longleftrightarrow b^{-1}$ .

Here  $g_b(x)$  is called the **quantum dilogarithm**.

- non-compact version of  $\text{Exp}_q(x)$  and  $\Gamma_q(x)$ .

## Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

New research program started in [Frenkel-Ip]

- Generalization of Teschner's representations to higher rank

Positive representations

- = "Quantization of minimal principal series representations"

Construction:

- Induced rep. of  $\mathcal{U}(\mathfrak{g})$  on  $L^2(U^+)$  by differential operators
- Lusztig's total positive space  $L^2(U_{>0}^+) \simeq L^2(\mathbb{R}_{>0}^{N=\dim U^+})$
- Mellin transformation:  $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$  differential operator  $\rightsquigarrow$  finite difference operator
- Quantization+ "Wick rotation"  
 $\implies$  positive operators  $\mathbf{e}_i, \mathbf{f}_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ 

## Theorem [Ip (2012)]

There exists a family of irreducible representations  $\mathcal{P}_\lambda$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :

- $\lambda \in \mathbb{R}_{\geq 0} P_+ \subset \mathfrak{h}_{\mathbb{R}}^*$  of classical  $\mathfrak{g}_{\mathbb{R}}$  ( $\iff \lambda \in \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ ).
- *Positivity*:  $\{E_i, F_i, K_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^{N=\dim U^+})$
- *Transcendental relations*: Define

$$\tilde{\mathbf{e}} := \mathbf{e}^{\frac{1}{b^2}}, \quad \tilde{\mathbf{f}} := \mathbf{f}^{\frac{1}{b^2}} \quad \tilde{K} := K^{\frac{1}{b^2}}$$

- *simply-laced*:  $\{\tilde{E}_i, \tilde{F}_i, \tilde{K}_i\}$  interchange  $b \longleftrightarrow b^{-1}$
- *non-simply-laced*:  $\{\tilde{E}_i, \tilde{F}_i, \tilde{K}_i\}$  generates the *Langlands dual*  $\mathcal{U}_{\tilde{q}}({}^L \mathfrak{g}_{\mathbb{R}})$
- $\{E_i, F_i, K_i\}$  commute weakly with  $\{\tilde{E}_i, \tilde{F}_i, \tilde{K}_i\}$  up to a sign.
- Does not depend on choice of reduced expression of  $w_0$

Positive Representations of  $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$ 

## Theorem [Ip (2012) cont'd]

Let

$$U_k = e^{2\pi b x_k}, V_k = e^{2\pi b p_k}$$

generate the quantum planes:

$$U_k V_k = q^2 V_k U_k.$$

Then  $\{\mathbf{e}_i, \mathbf{f}_i, K_i\}_{i=1}^n =$  *Laurent polynomials* in  $\{U_k, V_k\}_{k=1}^N$ .

- *Positive coefficients*  $\in \mathbb{Z}_{\geq 0}[q, q^{-1}]$
- $K_i =$  *multiplication operators.*

(Ignoring  $*$ -structure:)

Extends the **Feigin map** of  $\mathcal{U}_q(\mathfrak{b}) \hookrightarrow \mathbb{C}\langle \mathbb{T}^N \rangle$  to the **whole**  $\mathcal{U}_q(\mathfrak{g})$ .

# Relations to Other Areas

## Multiplier Hopf Algebra

$G$  compact,  $\Delta : C[G] \longrightarrow C[G] \otimes C[G]$ :

$$\Delta f(g_1, g_2) = f(g_1 g_2)$$

$G$  locally compact:  $\Delta : C_0(G) \rightarrow C_0(G) \otimes C_0(G)$ !

### Definition

*Multiplier algebra  $M(\mathcal{A})$  of a  $C^*$ -algebra  $\mathcal{A} \subset \mathcal{B}(\mathcal{H})$  is the  $C^*$ -algebra*

$$M(\mathcal{A}) = \{b \in \mathcal{B}(\mathcal{H}) : b\mathcal{A} \subset \mathcal{A}, \mathcal{A}b \subset \mathcal{A}\}$$

### Example

$\mathcal{A} = C_0(G)$ ,  $M(\mathcal{A}) = C_b(G)$ ,  $\Delta : \mathcal{A} \longrightarrow M(\mathcal{A} \otimes \mathcal{A})$

### Definition

*Multiplier Hopf algebra  $\mathcal{A}$ :  $\Delta, \epsilon, S$  extends to homomorphisms of  $M(\mathcal{A})$*

## Multiplier Hopf Algebra

- Idea: work with bounded operators

$U, V$  (positive unbounded operators)

$\rightsquigarrow U^{is}, V^{it}$  (unitary operators)

$\rightsquigarrow \iint_{\mathbb{R} \times \mathbb{R}} f(s, t) U^{is} V^{it} ds dt$  (bounded operators generated by  $U, V$ )

- Non-simple root:

$$T_i(\mathbf{e}_j) := \mathbf{e}_{ij} := \frac{q^{\frac{1}{2}} \mathbf{e}_j \mathbf{e}_i - q^{-\frac{1}{2}} \mathbf{e}_i \mathbf{e}_j}{q - q^{-1}}$$

Also positive self-adjoint in  $\mathcal{P}_\lambda$ !

- Continuous PBW basis:

$$\prod_{\alpha \in \Delta_+}^{\rightarrow} E_\alpha^{k_\alpha} \rightsquigarrow \prod_{\alpha \in \Delta_+}^{\rightarrow} \mathbf{e}_\alpha^{it_\alpha}$$

## Drinfeld-Jimbo quantum groups in $C^*$ -algebra!

Techniques from  $C^*$ -algebra and Non-Commutative Geometry:

- Multiplier Hopf algebra [van Daele]
- Locally compact quantum groups [Kustermans-Vaes]
- Multiplicative unitary  $W$  [Woronowicz]:

$$\Delta(x) = W(x \otimes 1)W^*, \quad x \in \mathcal{A}$$

- Gelfand-Naimark-Segal (GNS) representations

**Theorem [Ip (2013)]**

*Peter-Weyl theorem:*

$$L^2(SL_{q\tilde{q}}^+(2, \mathbb{R})) \simeq \int_{\mathbb{R}_+}^{\oplus} \mathcal{P}_\gamma \otimes \mathcal{P}_\gamma^* d\mu(\gamma)$$

*as regular  $\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(2, \mathbb{R}))$  representation.*

*(GNS representation + Drinfeld Double + Hopf pairing)*

## Positive Casimirs

- Center of  $\mathcal{U}(\mathfrak{sl}_2)$ :  $C = FE + \frac{1}{2}(H + 1)$
- Center of  $\mathcal{U}_q(\mathfrak{sl}_2)$ :  $C = FE + \left[\frac{1}{2}(H + 1)\right]_q^2$   

$$= FE + \frac{qK + q^{-1}K^{-1}}{(q - q^{-1})^2} + \text{const}$$

- Rescale by  $\left(\frac{i}{q - q^{-1}}\right)^2$ :

$$C = \mathbf{fe} - qK - q^{-1}K^{-1}$$

- $C \curvearrowright \mathcal{P}_\lambda$  as **positive scalar**:

$$C = e^{2\pi b\lambda} + e^{-2\pi b\lambda} > 0$$

Spectral decomposition of  $\Delta(C)$  on  $\mathcal{P}_\lambda \otimes \mathcal{P}_\mu$

$\implies$  **Tensor product decomposition!**

## Positive Casimirs

- Center of  $\mathcal{U}(\mathfrak{sl}_2)$ :  $C = FE + \frac{1}{2}(H + 1)$
- Center of  $\mathcal{U}_q(\mathfrak{sl}_2)$ :  $C = FE + \left[\frac{1}{2}(H + 1)\right]_q^2$   

$$= FE + \frac{qK + q^{-1}K^{-1}}{(q - q^{-1})^2} + \text{const}$$

- Rescale by  $\left(\frac{i}{q - q^{-1}}\right)^2$ :

$$C = \mathbf{fe} - qK - q^{-1}K^{-1}$$

- $C \curvearrowright \mathcal{P}_\lambda$  as **positive scalar**:

$$C = e^{2\pi b\lambda} + e^{-2\pi b\lambda} > 0$$

Spectral decomposition of  $\Delta(C)$  on  $\mathcal{P}_\lambda \otimes \mathcal{P}_\mu$

$\implies$  **Tensor product decomposition!**

## Positive Casimirs

Center of  $\mathcal{U}_q(\mathfrak{g})$  generated by  $n = \text{rank } \mathfrak{g}$  Casimir elements

- [Zhang-Gould-Bracken]

$$C_k := (\text{Tr}_q|_{V_k})(R_{21}R), \quad k = 1, \dots, n$$

- $V_k = k$ th fundamental representation of  $\mathcal{U}_q(\mathfrak{g})$

### Theorem [Ip (2016)]

Each  $C_k$  acts on  $\mathcal{P}_\lambda$  as *positive* scalar  $C_k(\vec{\lambda})$

$$C_k(\vec{\lambda}) = \sum_{V_k^\mu \subset V_k} e^{-4\pi b \mu(\vec{\lambda} \cdot \vec{W})} \geq \dim V_k$$

where  $\mu = \text{weight of } V_k^\mu$ ,  $\vec{W} = \text{fundamental coweight}$ .

Idea: *virtual highest weights*  $\longleftrightarrow$  “analytic continuation” from  $\mathcal{U}_q(\mathfrak{g})!$

## Positive Casimirs

Center of  $\mathcal{U}_q(\mathfrak{g})$  generated by  $n = \text{rank } \mathfrak{g}$  Casimir elements

- [Zhang-Gould-Bracken]

$$C_k := (\text{Tr}_q|_{V_k})(R_{21}R), \quad k = 1, \dots, n$$

- $V_k = k$ th fundamental representation of  $\mathcal{U}_q(\mathfrak{g})$

### Theorem [Ip (2016)]

Each  $C_k$  acts on  $\mathcal{P}_\lambda$  as *positive* scalar  $C_k(\vec{\lambda})$

$$C_k(\vec{\lambda}) = \sum_{V_k^\mu \subset V_k} e^{-4\pi b \mu(\vec{\lambda} \cdot \vec{W})} \geq \dim V_k$$

where  $\mu = \text{weight of } V_k^\mu$ ,  $\vec{W} = \text{fundamental coweight}$ .

Idea: *virtual highest weights*  $\longleftrightarrow$  “analytic continuation” from  $\mathcal{U}_q(\mathfrak{g})!$

## Example: Type $A_2$

Center of  $\mathcal{U}_q(\mathfrak{sl}_3)$  generated by  $C_1$  and  $C_2$ :

$$C_1 = K_1^{\frac{1}{3}} K_2^{-\frac{1}{3}} (q^{-2} K_1 K_2 + K_1^{-1} K_2 + q^2 K_1^{-1} K_2^{-1} - q^{-1} K_2 \mathbf{e}_1 \mathbf{f}_1 - q K_1^{-1} \mathbf{e}_2 \mathbf{f}_2 + \mathbf{e}_{21} \mathbf{f}_{12})$$

$$C_2 = K_1^{-\frac{1}{3}} K_2^{\frac{1}{3}} (q^{-2} K_1 K_2 + K_1 K_2^{-1} + q^2 K_1^{-1} K_2^{-1} - q K_2^{-1} \mathbf{e}_1 \mathbf{f}_1 - q^{-1} K_1 \mathbf{e}_2 \mathbf{f}_2 + \mathbf{e}_{12} \mathbf{f}_{21})$$

On  $\mathcal{P}_\lambda$ , each  $C_k$  reduced from 35 terms to **positive** scalar:

$$C_1 \curvearrowright \mathcal{P}_\lambda = e^{\frac{8\pi b\lambda_1}{3} + \frac{4\pi b\lambda_2}{3}} + e^{-\frac{4\pi b\lambda_1}{3} + \frac{4\pi b\lambda_2}{3}} + e^{-\frac{4\pi b\lambda_1}{3} - \frac{8\pi b\lambda_2}{3}} \geq 3$$

$$C_2 \curvearrowright \mathcal{P}_\lambda = e^{\frac{4\pi b\lambda_1}{3} + \frac{8\pi b\lambda_2}{3}} + e^{\frac{4\pi b\lambda_1}{3} - \frac{4\pi b\lambda_2}{3}} + e^{-\frac{8\pi b\lambda_1}{3} - \frac{4\pi b\lambda_2}{3}} \geq 3$$




Figure: Type  $A_2$  Region

## Example: Type $A_2$

Center of  $\mathcal{U}_q(\mathfrak{sl}_3)$  generated by  $C_1$  and  $C_2$ :

$$C_1 = K_1^{\frac{1}{3}} K_2^{-\frac{1}{3}} (q^{-2} K_1 K_2 + K_1^{-1} K_2 + q^2 K_1^{-1} K_2^{-1} - q^{-1} K_2 \mathbf{e}_1 \mathbf{f}_1 - q K_1^{-1} \mathbf{e}_2 \mathbf{f}_2 + \mathbf{e}_{21} \mathbf{f}_{12})$$

$$C_2 = K_1^{-\frac{1}{3}} K_2^{\frac{1}{3}} (q^{-2} K_1 K_2 + K_1 K_2^{-1} + q^2 K_1^{-1} K_2^{-1} - q K_2^{-1} \mathbf{e}_1 \mathbf{f}_1 - q^{-1} K_1 \mathbf{e}_2 \mathbf{f}_2 + \mathbf{e}_{12} \mathbf{f}_{21})$$

On  $\mathcal{P}_\lambda$ , each  $C_k$  reduced from 35 terms to **positive** scalar:

$$C_1 \curvearrowright \mathcal{P}_\lambda = e^{\frac{8\pi b\lambda_1}{3} + \frac{4\pi b\lambda_2}{3}} + e^{-\frac{4\pi b\lambda_1}{3} + \frac{4\pi b\lambda_2}{3}} + e^{-\frac{4\pi b\lambda_1}{3} - \frac{8\pi b\lambda_2}{3}} \geq 3$$

$$C_2 \curvearrowright \mathcal{P}_\lambda = e^{\frac{4\pi b\lambda_1}{3} + \frac{8\pi b\lambda_2}{3}} + e^{\frac{4\pi b\lambda_1}{3} - \frac{4\pi b\lambda_2}{3}} + e^{-\frac{8\pi b\lambda_1}{3} - \frac{4\pi b\lambda_2}{3}} \geq 3$$




Figure: Type  $A_2$  Region

## Discriminant Variety

Region in  $\mathbb{R}^n$  defined by  $\Phi : (\lambda_1, \dots, \lambda_n) \mapsto (C_1(\vec{\lambda}), \dots, C_n(\vec{\lambda}))$

- Singularity at  $\Phi(0, \dots, 0)$
- Boundary described by **discriminant** of some polynomial

Example of type  $A_2$ :

- Boundary:

$$(XY + 9)^2 = 4(X^3 + Y^3 + 27)$$

$$\iff \text{Disc}_z(z^3 + Xz^2 + Yz + 1) = 0$$

- Type  $A_2$  singularity

Relation to simple (du Val) singularity theory?

# Quantum Teichmüller Theory




Given  $S =$  surface with punctures

- $\mathcal{T}_S =$  Teichmüller space with Weil-Petersson Poisson structure
- Mapping class group  $MCG(S) \curvearrowright \mathcal{T}_S$
- Quantization  $\mathcal{T}_S^q$  represented on space of states  $\mathbf{H} = \otimes_{\tau \in \Delta} \mathcal{H}$
- $g \in MCG(S) \rightsquigarrow$  unitary operator  $\rho(g) \curvearrowright \mathbf{H}$
- Goal: Projective unitary representations of  $MCG(S)$

Remark

$\mathbf{H} \simeq$  Space of conformal blocks in Liouville CFT

# Quantum Teichmüller Theory



The pentagon equation for  $T$



# Quantum Teichmüller Theory

Two main approaches based on triangulations of  $S$ :

- **Kashaev's** coordinate:  $(x, p)$  on each  $\Delta$   
 $\implies$  Kashaev's groupoid  $\{\mathbf{T}_{jk}, \mathbf{A}_j\}_{j,k \in \Delta}$  associated to change of dotted ideal triangulations  $\Delta$
- **Thurston's** shear coordinate (**Penner's** lambda length)  
 $\lambda_a, \lambda_b, \lambda_c$  on each edge  
 $\implies$  Fock-Goncharov cluster varieties  $\mathcal{X}_{PSL(2, \mathbb{R}), S}$

## Higher Rank Construction

Quantum plane = Borel part of  $\mathcal{U}_{q\bar{q}}(\mathfrak{sl}(2, \mathbb{R}))!$

- Replace  $\mathcal{H}$  by  $\mathcal{P}_\lambda^b$ , the restriction of  $\mathcal{P}_\lambda$  from  $\mathcal{U}_{q\bar{q}}(\mathfrak{g}_{\mathbb{R}})$  to  $\mathcal{U}_{q\bar{q}}(\mathfrak{b}_{\mathbb{R}})$
- $\mathcal{P}^b := \mathcal{P}_\lambda^b$  does not depend on  $\lambda$

### Theorem

$\mathcal{P}^b$  is closed under tensor product:

$$\mathcal{P}^b \otimes \mathcal{P}^b \simeq \mathcal{P}^b \otimes M$$

$\implies$  gives quantum mutation operators  $\mathbf{T}$ .

(Ongoing) Construct Kashaev's  $\mathbf{A}$  operators by dual representations.

$\implies$  Candidate for quantum higher Teichmüller theory

$\implies$  New projective unitary representations of  $MCG(S)$ .

(?) Relation to Fock-Goncharov's cluster varieties

## Higher Rank Construction

Quantum plane = **Borel part** of  $\mathcal{U}_{q\bar{q}}(\mathfrak{sl}(2, \mathbb{R}))!$

- Replace  $\mathcal{H}$  by  $\mathcal{P}_\lambda^b$ , the restriction of  $\mathcal{P}_\lambda$  from  $\mathcal{U}_{q\bar{q}}(\mathfrak{g}_{\mathbb{R}})$  to  $\mathcal{U}_{q\bar{q}}(\mathfrak{b}_{\mathbb{R}})$
- $\mathcal{P}^b := \mathcal{P}_\lambda^b$  does not depend on  $\lambda$

### Theorem

$\mathcal{P}^b$  is closed under tensor product:

$$\mathcal{P}^b \otimes \mathcal{P}^b \simeq \mathcal{P}^b \otimes M$$

$\implies$  gives *quantum mutation operators*  $\mathbf{T}$ .

(Ongoing) Construct Kashaev's  $\mathbf{A}$  operators by dual representations.

$\implies$  Candidate for quantum higher Teichmüller theory

$\implies$  New projective unitary representations of  $MCG(S)$ .

(?) Relation to **Fock-Goncharov's** cluster varieties



## Cluster realization of $\mathcal{U}_q(\mathfrak{g})$


- Teichmüller theory  $\longleftrightarrow$  “Moduli space of  $PSL(2, \mathbb{R})$ -local system”
- “Higher Teichmüller theory”  $\longleftrightarrow$  “Moduli space of  $G$ -local system”

Fock-Goncharov’s cluster varieties  $\mathcal{X}_{G,S}$  on surface  $S$

- Poisson structure on  $\mathcal{X}_{G,S}$ : described by quivers on  $\Delta$
- $\rightsquigarrow$  Quantum torus algebra  $\mathcal{X}_{G,S}^q$

### Example

$\mathcal{X}_{sl_4, \Delta}^q$ -quiver:



# Cluster realization of $\mathcal{U}_q(\mathfrak{g})$

Positive Representations  $\implies$

**Theorem [Schrader-Shapiro, Ip (2016)]**


*There is embedding for  $\mathfrak{g}$  of any simple type:*


$$\mathcal{U}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_{G,S}^q / \sim$$

*$S = \text{disk with 1 puncture and 2 marked points}$*


**Theorem [Schrader-Shapiro, Ip (2016)]**


*Universal R matrix  $\longleftrightarrow$  Quiver mutations giving half-Dehn twist*






The quantum cluster algebra  $\mathcal{X}_{\mathfrak{sl}_4, S}^q$ :





The quantum cluster algebra  $\mathcal{X}_{\mathfrak{sl}_4, S}^q$ :



Embedding of  $F_i \in \mathcal{U}_q(\mathfrak{sl}_4) \longrightarrow \mathcal{X}_{\mathfrak{sl}_4, S}^q$   
 [Schrader-Shapiro (2016)]

# Cluster realization of $\mathcal{U}_q(\mathfrak{g})$




Figure: Cluster realization of Quantum Groups of type  $E_8$

## Modular Double of $\mathcal{U}_q(\mathfrak{osp}(1|2))$

$Cl_2 =$  Clifford algebra  $\langle \xi, \eta \rangle$  such that

$$\xi^2 = \eta^2 = 1, \quad \eta\xi + \xi\eta = 1$$

Spinor trick:  $Rep(\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))) \longrightarrow Rep(\mathcal{U}_q(\mathfrak{osp}(1|2)))$

**Theorem [Ip-Zeitlin (2013)]**

Let  $q_* = iq$ . Given a representation of  $\mathcal{U}_{q_*}(\mathfrak{sl}(2, \mathbb{R}))$ , there exists a representation of  $\mathcal{U}_q(\mathfrak{osp}(1|2)) = \langle \mathcal{E}, \mathcal{F}, \mathcal{K}^{\pm 1} \rangle$  by


$$\mathcal{E} = \alpha E\xi, \quad \mathcal{F} = F\eta, \quad \mathcal{K} = K i\eta\xi$$

where  $\alpha = i \frac{q+q^{-1}}{q-q^{-1}} > 0$ .

In particular, induces **Modular Double** structure:

$$\mathcal{U}_{q_*, \tilde{q}_*}(\mathfrak{sl}(2, \mathbb{R})) \implies \mathcal{U}_{q, \tau(q)}(\mathfrak{osp}(1|2)), \quad \tau(q) = -i\tilde{q}_*$$

# Summary



# Future Perspectives

