

Title: Spectrum of conformal gauge theories on a torus

Date: Jun 16, 2017 10:00 AM

URL: <http://pirsa.org/17060058>

Abstract: <p>Many model quantum spin systems have been proposed to realize critical points or phases described by 2+1 dimensional conformal gauge theories. On a torus of size L and modular parameter τ , the energy levels of such gauge theories equal $(1/L) \times$ universal functions of τ . We compute the universal spectrum of QED3, a U(1) gauge theory with N_f two-component massless Dirac fermions, in the large- N_f limit. We also allow for a Chern-Simons term at level k , and show how the topological k -fold ground state degeneracy in the absence of fermions transforms into the universal spectrum in the presence of fermions; these computations are performed at fixed N_f/k in the large- N_f limit.</p>

Spectrum of conformal gauge theories on a torus

Alex Thomson

AT, S. Sachdev, Phys. Rev. B 95, 205128 (2017)
arXiv:1607.05279

1

QED₃: a critical spin liquid

- ✧ QED₃ is a strongly coupled conformal field theory (CFT) in 2+1d describing a U(1) gauge boson coupled to N_f two-component Dirac spinors:

$$\mathcal{L}_{\text{qed}} = -\bar{\psi}_\alpha i\gamma^\mu (\partial_\mu - iA_\mu) \psi_\alpha + \frac{1}{4e^2} (\partial_\mu A_\nu - \partial_\nu A_\mu)^2$$

It has a SU(N_f) symmetry under which the fermion flavours rotate into one another.

QED₃: a critical spin liquid

- ✧ QED₃ is a strongly coupled conformal field theory (CFT) in 2+1d describing a U(1) gauge boson coupled to N_f two-component Dirac spinors:

$$\mathcal{L}_{\text{qed}} = -\bar{\psi}_\alpha i\gamma^\mu (\partial_\mu - iA_\mu) \psi_\alpha + \frac{1}{4e^2} (\partial_\mu A_\nu - \partial_\nu A_\mu)^2$$

It has a SU(N_f) symmetry under which the fermion flavours rotate into one another.

- ✧ It describes a gapless, critical, U(1) spin liquid

➡ Might describe:

- ✧ the spin-1/2 kagome Heisenberg antiferromagnet
 - ✧ the J_1 - J_2 antiferromagnet on the triangle lattice.
- ✧ It is also the proposed critical theory of a number of deconfined critical points.

Universality

- ✧ QED_3 is a conformal field theory and is therefore characterized by a number of *universal* properties
 - ✧ Depends only on IR properties, no UV dependence
- ✧ Torus spectrum may be useful to help identify the CFT in a numerical simulation.
 - ➡ Should depend only on IR cutoffs such as size and shape of torus, not the details of the lattice being simulated.

Previous work has been done for the Ising & $O(N)$ CFTs...


Schuler et al., PRL 117, 210401 (2016)
Whitsitt, Sachdev, Phys. Rev. B 94, 085134 (2016)
Whitsitt et al., arXiv:1701.03111

3

Ising spectrum on torii

Critical Ising model:
$$H_{\text{TFI}} = - \sum_{\langle ij \rangle} \sigma_i^z \sigma_j^z - g_c \sum_i \sigma_i^x$$



ED and QMC on multiple lattices:



UV data

Spectrum depends only on modular parameter, τ , length L , and speed of light c .

IR data



Schuler, Whitsitt, Henry, Sachdev, Läuchli, PRL 117, 210401 (2016)

Plan

- ✧ Kagome antiferromagnet and QED_3
- ✧ Path integral approach to torus spectrum
 - ✧ QED_3 in the large- N_f limit
- ✧ QED_3 Spectrum


Kagome antiferromagnet

$$H_H = J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j$$

✧ Parton construction:

$$\mathbf{S}_i = \frac{1}{2} f_{i\alpha}^\dagger \boldsymbol{\sigma}_{\alpha\beta} f_{i\beta}, \quad 1 = \sum_{\alpha} f_{i\alpha}^\dagger f_{i\alpha}$$

$$\longrightarrow H_H = -\frac{J}{2} \sum_{\langle ij \rangle} f_{i\alpha}^\dagger f_{j\alpha} f_{i\beta}^\dagger f_{j\beta}$$




Kagome antiferromagnet

$$H_H = J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j$$

✧ Parton construction:

$$\mathbf{S}_i = \frac{1}{2} f_{i\alpha}^\dagger \boldsymbol{\sigma}_{\alpha\beta} f_{i\beta}, \quad 1 = \sum_{\alpha} f_{i\alpha}^\dagger f_{i\alpha}$$

$$\longrightarrow H_H = -\frac{J}{2} \sum_{\langle ij \rangle} f_{i\alpha}^\dagger f_{j\alpha} f_{i\beta}^\dagger f_{j\beta}$$



✧ Mean field theory: decouple the 4-fermion interaction with a Hubbard-Stratonovich transformation.

$$H_{\text{MF}} = - \sum_{ij} t_{ij} f_{i\alpha}^\dagger f_{j\alpha} + H.c. \quad t_{ij} = J \langle f_{i\alpha}^\dagger f_{j\alpha} \rangle$$

✧ Ansatz with π -flux per hexagon gives 2 Dirac cones per spin:

$$\mathcal{L}_{\text{MF}} = -\bar{\psi}_{\sigma a} \gamma^\mu \partial_\mu \psi_{\sigma a} \quad \underbrace{\sigma = \uparrow, \downarrow}_{\text{spin}} \quad \underbrace{a = \pm}_{\text{valley: } \pm \mathbf{Q}}$$

Critical spin liquid

- ✧ Spin operators invariant under $f_{i\alpha} \rightarrow e^{i\phi_i} f_{i\alpha}$: the slave fermions carry a gauge charge.

→
$$\mathcal{L}_{\text{qed}} = -\bar{\psi}_\alpha i\gamma^\mu (\partial_\mu - iA_\mu) \psi_\alpha + \frac{1}{4e^2} (\partial_\mu A_\nu - \partial_\nu A_\mu)^2$$

- ✧ QED₃ with $N_f=4$ is **strongly coupled**. We work in the $N_f=\infty$ limit.

- ✧ We can also add a Chern-Simons (CS) term to the action:

$$\mathcal{L}_{\text{CS}} = \frac{ik}{4\pi} \epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho$$

This CFT can describe certain phase transitions between topological phases.

Path integral definition on square torus

Partition
function


$$Z = \frac{1}{\text{Vol}(G)} \int DA D\psi e^{-S[A,\psi]}$$

$\text{Vol}(G)$ = gauge group
volume

✧ Decompose gauge field as $A_\mu = a_\mu + B_\mu + \partial_\mu \phi$ $\partial^2 \phi = 0$

✧ a_μ is the **zero momentum mode**: not pure gauge on the torus

$a_\mu \rightarrow a_\mu + \frac{2\pi}{L}$ \Rightarrow a_μ is a periodic
constant, $a_\mu \in [0, 2\pi/L)$



✧ B_μ is the finite-momentum **gauge-fixed potential**: $\partial^\mu B_\mu = 0$.

✧ $\partial_\mu \phi$ parametrizes an extra **gauge degree of freedom**.

Path integral normalization

$$Z = \frac{1}{\text{Vol}(G)} \int DA D\psi e^{-S[A,\psi]} \quad A_\mu = a_\mu + B_\mu + \partial_\mu \phi$$

✧ Measure of integration:

$$DA = Da DB D(d\phi) = Da DB D'\phi \sqrt{\det'(-\nabla^2)} \quad \leftarrow \text{Faddeev-Popov determinant}$$

✧ Gauge group volume: $\text{Vol}(G) = 2\pi \sqrt{\beta L^2} \int D'\phi$

$$\Rightarrow Z = \frac{\beta L^2}{2\pi} \sqrt{\det'(-\nabla^2)} \int d^3 a DB D\psi e^{-S[\psi, B, a]}$$

Large- N_f QED₃

✧ Couple gauge field to N_f two-component Dirac fermions:

QED₃

$$S_{\text{qed}}[\psi, A] = - \int d^3r \bar{\psi}_\alpha i\gamma^\mu (\partial_\mu - iA_\mu) \psi_\alpha$$

N_f even

Chern-Simons

$$S_{\text{CS}}[A] = \frac{ik}{4\pi} \int d^3r \epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho$$

$k = 0$ or $k \sim \mathcal{O}(N_f)$

✧ S_{qed} is quadratic in the fermions, so integrate them out:

$$Z = \frac{\beta L^2}{2\pi} \sqrt{\det'(-\nabla^2)} \int d^3a DB e^{-S_{\text{CS}}[B] + N_f \log \det(i\cancel{\partial} + \cancel{\partial} + B)},$$

Saddle-point configurations

$$Z = \frac{\beta L^2}{2\pi} \sqrt{\det'(-\nabla^2)} \int d^3 a DB e^{-S_{CS}[B] + N_f \log \det(i\cancel{D} + \cancel{A} + \cancel{B})},$$

✧ Plane: $A_\mu = \text{const.}$ are all gauge equivalent

➡ $A_\mu = 0$ is the saddle-point solution.

✧ Torus: finite zero mode a_μ exists

➡ must minimize the free energy with respect to a_μ

✧ Rescale $B \rightarrow B/\sqrt{N_f}$ and expand in powers of N_f :

$$N_f \log \det(i\cancel{D}) = \underbrace{N_f \text{tr} \log(i\cancel{D} + \cancel{A})}_{\text{free fermion free energy } F_0(a)} + \underbrace{\sqrt{N_f} \text{tr} \left(\frac{1}{i\cancel{D} + \cancel{A}} \cancel{B} \right)}_{\text{vanishes}} - \underbrace{\frac{1}{2} \text{tr} \left(\frac{1}{i\cancel{D} + \cancel{A}} \cancel{B} \frac{1}{i\cancel{D} + \cancel{A}} \cancel{B} \right)}_{\text{Gauge field fluctuations Suppressed by } 1/N_f} + \dots$$

Minimization of free energy

$$N_f \log \det (i\mathcal{D}) = N_f \text{tr} \log (i\mathcal{D} + \phi) - \frac{1}{2} \text{tr} \left(\frac{1}{i\mathcal{D} + \phi} \mathcal{B} \frac{1}{i\mathcal{D} + \phi} \mathcal{B} \right) + \dots$$


✧ The leading contribution is simply the free Dirac free energy:

$$F_0(a) = -\text{tr} \log (i\mathcal{D} + \phi) = -\sum_p \log (p + a)^2$$

✧ Find that anti-periodic boundary conditions minimize F_0 :

$$\bar{a}_{x,y} = 0$$

$$p_{x,y} = \frac{2\pi}{L} \left(n_{x,y} + \frac{1}{2} \right)$$



12

Free Dirac-like states

Free Dirac Hamiltonian $\mathcal{H}_D = \sum_{\mathbf{p}} |\mathbf{p}| \left[\chi_{+\alpha}^\dagger(\mathbf{p}) \chi_{+\alpha}(\mathbf{p}) - \chi_{-\alpha}^\dagger(\mathbf{p}) \chi_{-\alpha}(\mathbf{p}) \right]$

particles
holes

✧ Constraint 1:

- ✧ $\chi_{\pm\alpha}(\mathbf{p})$ are not gauge invariant. The lowest energy states fermion states relevant to QED₃ are

$$\chi_{+\alpha}^\dagger(\mathbf{p} + \mathbf{q}) \chi_{-\beta}(\mathbf{p}) |0\rangle, \quad \chi_{+\alpha}^\dagger(-\mathbf{p}) \chi_{-\beta}(-\mathbf{p} - \mathbf{q}) |0\rangle$$

with energy $E_f(\mathbf{q}, \mathbf{p}) = |\mathbf{q} + \mathbf{p}| + |\mathbf{p}|$.

- ✧ The free theory has $2N_f^2$ such states (except when $\mathbf{p} + \mathbf{q} = -\mathbf{p}$, in which case there are N_f^2).

Gauge constraint

$$\chi_{+\alpha}^\dagger(\mathbf{p} + \mathbf{q})\chi_{-\beta}(\mathbf{p}) |0\rangle, \quad \chi_{+\alpha}^\dagger(-\mathbf{p})\chi_{-\beta}(-\mathbf{p} - \mathbf{q}) |0\rangle$$

✧ Constraint 2:

- ✧ The saddle-point approximation is equivalent to imposing the classical equations of motion:

$$\underbrace{\sum_{\alpha} \bar{\psi}_{\alpha} \gamma^{\mu} \psi_{\alpha}}_{J_{\mu}} = \frac{i}{e^2} \partial_{\nu} F^{\mu\nu} \quad \boxed{\text{Gauss-Ampère Law}}$$

Gauge constraint

$$\chi_{+\alpha}^\dagger(\mathbf{p} + \mathbf{q})\chi_{-\beta}(\mathbf{p}) |0\rangle, \quad \chi_{+\alpha}^\dagger(-\mathbf{p})\chi_{-\beta}(-\mathbf{p} - \mathbf{q}) |0\rangle$$

✧ Constraint 2:

- ✧ The saddle-point approximation is equivalent to imposing the classical equations of motion:

$$\underbrace{\sum_{\alpha} \bar{\psi}_{\alpha} \gamma^{\mu} \psi_{\alpha}}_{J_{\mu}} = \cancel{\frac{i}{\epsilon} \partial_{\nu} T^{\mu\nu}} \quad \boxed{\text{Gauss-Ampère Law}}$$

- ✧ All states corresponding to the poles of $\langle J_{\mu}(x) J_{\nu}(0) \rangle$ are removed. This corresponds to the removal of a single state for each momentum pair \mathbf{p} & \mathbf{q} .


➡ There are $2N_f^2 - 1$ free fermion-like states in QED₃ (except when $\mathbf{p} + \mathbf{q} = -\mathbf{p}$, in which case there are $N_f^2 - 1$).

Fermion energies

◇ Fermion energies: $E_f(\mathbf{q}, \mathbf{p}) = |\mathbf{q} + \mathbf{p}| + |\mathbf{p}|$.

$$\mathbf{q} = \frac{2\pi}{L} \left(m_x + \frac{1}{2}, m_y + \frac{1}{2} \right), \quad \mathbf{p} = \frac{2\pi}{L} \left(n_x + \frac{1}{2}, n_y + \frac{1}{2} \right),$$

$$E_f(\mathbf{q}, \mathbf{p}) = \frac{2\pi}{L} \left[\sqrt{(n_x + m_x + 1)^2 + (n_x + m_y + 1)^2} + \sqrt{(n_x + 1/2)^2 + (n_y + 1/2)^2} \right]$$



$\bar{\mathbf{q}} = (0,0)$	
\bar{E}_f	d_f
1.414214	$4N_f^2 - 2$
3.162278	$8N_f^2 - 4$
4.242640	$4N_f^2 - 2$

$$\bar{\mathbf{q}} = L\mathbf{q}/2\pi,$$

$$\bar{E}_f = LE_f/2\pi$$

Photon contribution

→ the spectrum is non-extensive in N_f , so $\mathcal{O}(1)$ corrections must be included as well.

$$N_f \log \det (i\mathcal{D}) = N_f \text{tr} \log (i\mathcal{D} + \not{a}) - \frac{1}{2} \text{tr} \left(\frac{1}{i\mathcal{D} + \not{a}} \not{B} \frac{1}{i\mathcal{D} + \not{a}} \not{B} \right) + \dots$$

✧ The only contribution at next-to-leading order is from the photon:

$$S_{\text{eff}}[B] = \frac{1}{2} \sum_q B_\mu(-q) \Pi^{\mu\nu}(q) B_\nu(q),$$

$$\Pi^{\mu\nu} = \Pi_f^{\mu\nu} + \Pi_{\text{CS}}^{\mu\nu}$$

$$\Pi_f^{\mu\nu}(q) = \mu \begin{array}{c} \alpha \quad \alpha \\ \curvearrowright \quad \curvearrowleft \\ \alpha \quad \alpha \end{array} \nu$$

= quadratic term in determinant expansion.

$$\Pi_{\text{CS}}^{\mu\nu}(q) = \frac{1}{2\pi\lambda} \epsilon^{\mu\nu\rho} q_\rho$$

$$\lambda = \frac{N_f}{k}$$

Photon states

$$S_{\text{eff}}[B] = \frac{N_f}{2} \sum_q B_\mu(-q) \Pi^{\mu\nu}(q) B_\nu(q), \quad \Pi^{\mu\nu}(q) = \Pi_f^{\mu\nu}(q) + \Pi_{\text{CS}}^{\mu\nu}(q)$$

✧ When $k=0$, this is simply the expansion:

$$\begin{array}{c} \mu \\ \text{~~~~~} \\ \nu \end{array} = \begin{array}{c} \mu \\ \text{.....} \\ \nu \end{array} + \begin{array}{c} \mu \\ \text{.....} \end{array} \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \begin{array}{c} \nu \\ \text{.....} \end{array} + \begin{array}{c} \mu \\ \text{.....} \end{array} \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \begin{array}{c} \nu \\ \text{.....} \end{array} + \dots$$

Spectrum from path integral

$$\text{Harmonic Oscillator} \quad H = \frac{1}{2}p^2 + \frac{1}{2}\Omega^2 q^2 \quad Z = \sum_{n=0}^{\infty} e^{-\beta(n+\frac{1}{2})\Omega}$$


✧ Path integral:

$$Z = \int Dq \exp \left[-\frac{1}{2} \int_0^\beta d\tau q(\tau) (-\partial_\tau^2 + \Omega^2) q(\tau) \right] = \left[\det \left(\frac{-\partial_\tau^2 + \Omega^2}{2\pi} \right) \right]^{-1/2}$$

$$\begin{aligned} \Rightarrow F &= -\frac{1}{\beta} \log Z = \frac{1}{2\beta} \log \det(-\partial_\tau^2 + \Omega^2) \\ &= \frac{1}{2\beta} \sum_{\omega_n} \log [\omega_n^2 + \Omega^2] = -\frac{1}{\beta} \log \left(\frac{e^{-\beta\Omega/2}}{1 - e^{-\beta\Omega}} \right) \end{aligned}$$

Photon states in QED₃


✧ Argument of log for $\mathbf{q}=0$: $\mathcal{F}(\omega, \mathbf{q} = 0) = -\frac{\Pi_f^{xx}\Pi_f^{yy}}{\omega^2} + \frac{1}{4\pi^2\lambda}$



Plot of $\Pi_f^{xx}(\omega, \mathbf{q}=0)$ and $\pm|\omega|/\lambda$ for $\lambda=N_f/k=4$.

Photon states in QED₃

✧ Argument of log for $\mathbf{q}=0$: $\mathcal{F}(\omega, \mathbf{q} = 0) = -\frac{\Pi_f^{xx}\Pi_f^{yy}}{\omega^2} + \frac{1}{4\pi^2\lambda}$




✧ Π_f^{xx} diverges at E_f :

$$\Pi_f^{xx}(\omega, 0) \sim \frac{\Pi_\gamma(\omega^2 - \omega_\gamma^2)}{\Pi_{\mathbf{p}}(\omega^2 - E_f(0, \mathbf{p})^2)}$$

➡ These diverges effectively remove the states created by J^μ


Plot of $\Pi_f^{xx}(\omega, \mathbf{q}=0)$ and $\pm|\omega|/\lambda$ for $\lambda=N_f/k=4$.

Finite momentum: $\mathbf{q}_1 = 2\pi(1,0)/L$ ($k=0$)




$$\mathbf{q}_1 = \frac{2\pi}{L}(1, 0)$$

Comparison with fermion energies, $\mathbf{q}=0$



Photon modes as a function of k



✧ For $k/N_f \rightarrow 0$, the modes approach the QED₃ values.

✧ As $k/N_f \rightarrow \infty$, $\omega_0 \rightarrow 0$ and all higher energy modes approach the free theory energy $E_f = 2|\mathbf{p}|$.

➡ The (infinite) ground state degeneracy is 'restored'.

Plot of energy modes for $\mathbf{q}=0$ as a function of $1/\lambda = k/N_f$

Thank you

AT, S. Sachdev, Phys. Rev. B 95, 205128 (2017)
arXiv:1607.05279

23