

Title: Isotropising an anisotropic cyclic cosmology

Date: Oct 10, 2017 11:00 AM

URL: <http://pirsa.org/17100077>

Abstract: <p>Standard models of cosmology use inflation as a mechanism to resolve the isotropy and homogeneity problem of the universe as well as the flatness problem. However, due to various well known problems with the inflationary paradigm, there has been an ongoing search for alternatives. Perhaps the most famous among these is the cyclic universe scenario or scenarios which incorporate bounces. As these scenarios have a contracting phase in the evolution of the universe, it is reasonable to ask whether the problems of homogeneity and isotropy can still be resolved in these scenarios. In my talk, I will focus on the problem of the resolution of isotropy. In the contracting phase of the evolution, the mechanism of ekpyrosis is used in most cosmological scenarios which incorporate a contracting phase to mitigate the problem of anisotropies blowing up on approaching the bounce. I will start by studying anisotropic universes and I shall examine the effect of the addition of ultra-stiff anisotropic pressures on the ekpyrotic phase. I will then consider evolving such anisotropic universes through several cycles with increasing expansion maxima at each successive bounce. This eventually leads to flatness in the isotropic case. My aim will be to see if the resolution of the flatness problem also leads to a simultaneous resolution of the isotropy problem. In the last section of my talk, I will briefly consider the effect of non comoving velocities on the shape of this anisotropic bouncing universe.</p>



Isotropising anisotropic cyclic cosmologies

Chandrima Ganguly

DAMTP, University of Cambridge

Seminar, Perimeter Institute
10th October, 2017

- 1 Introduction
- 2 In the contracting phase
- 3 Ekpyrosis meets anisotropic pressures!
- 4 The Bianchi IX universe
- 5 Solving the flatness problem within the framework of bouncing cosmologies
- 6 Adding a cosmological constant to the cocktail
- 7 Conclusions

How do we get a bounce?



- Coming out of the contracting phase the Hubble rate H is negative.
- $H > 0$ in the expanding phase
- So in the transition or 'bounce' phase, $H = 0$ and

$$\dot{H} = \frac{k}{a^2} - \frac{1}{2}(\rho + P)$$

- If the spatial curvature k is 0, then for $\dot{H} > 0$ and $H = 0$, we must have $\rho + P < 0$ (NEC violation)
- If we have positive spatial curvature, we can have a bounce, In the closed radiation FRW universe, exact solutions show this but need a NEC violating field to have the bounce occur at non-zero volume.

J.D.Barrow and Christos G.Tsagas, CQG Vol. 26, No. 19 (2009)



3 / 39

Do the most general cyclic universes isotropise?



- Closed FRW universe with ordinary matter or dust shows oscillatory behaviour
- Simple solutions in these scenarios have been found



We focus on the isotropy problem and split it up into 2 regimes of interest.

- In the contracting branch, on approach to the singularity, or in the case of non-singular cosmologies, on approach to the bounce



We focus on the isotropy problem and split it up into 2 regimes of interest.

- In the contracting branch, on approach to the singularity, or in the case of non-singular cosmologies, on approach to the bounce
- Over a large period of oscillations with increasing expansion maxima

- 1 Introduction
- 2 In the contracting phase**
- 3 Ekpyrosis meets anisotropic pressures!
- 4 The Bianchi IX universe
- 5 Solving the flatness problem within the framework of bouncing cosmologies
- 6 Adding a cosmological constant to the cocktail
- 7 Conclusions

A simple example of ekpyrosis



- The metric

$$ds^2 = dt^2 - a^2(t)dx^2 - b^2(t)dy^2 - c^2(t)dz^2$$

- Friedmann equation: $3H^2 = \sigma^2 + \rho_{matter}$,
- The shear evolves as,

$$\dot{\sigma}_{\alpha\beta} + 3H\sigma_{\alpha\beta} = 0$$

- ρ_{matter} should evolve as V^{-n} , $n \gg 2$

J. Khoury, B.A. Ovrut, P. J. Steinhardt and N. Turok, 2001, J. High Energy Phys. 11(2001)041

Bianchi Class A cosmologies



- The generalised metric

$$ds^2 = dt^2 - h_{ab}d\omega^a d\omega^b$$

- Having an isotropic ultra stiff field of density ρ with equation of state $p = (\gamma - 1)\rho$, such that $\gamma > 2$

The phase plane system



- We introduce

$$\begin{aligned}\sigma_+ &\equiv \frac{1}{2}(\sigma_{22} + \sigma_{33}), \\ \sigma_- &\equiv \frac{1}{2\sqrt{3}}(\sigma_{22} - \sigma_{33}).\end{aligned}$$

- Write EFE in terms of expansion normalised variables

$$\Omega \equiv \frac{\rho}{3H^2}, \quad \Sigma^2 \equiv \frac{\sigma^2}{3H^2}, \quad K \equiv -\frac{{}^{(3)}R}{6H^2}.$$

The phase plane system looks like...



- Einstein equations of the form $\mathbf{x}' = \mathbf{f}(\mathbf{x})$
- subject to the Friedmann constraint $\mathbf{g}(\mathbf{x}) = 0$
- where the state vector $\mathbf{x} \in \mathbb{R}^6$ is given by
$$\{H, \underbrace{\Sigma_+, \Sigma_-}, \underbrace{N_1, N_2, N_3}, \Omega\}$$

shear components spatial curvature variables

- The fact that the matter is ultra stiff $\gamma > 2$ is used and
- A no-hair theorem can be proved for all Bianchi types, I-VIII as well as IX(separately)

Cosmic no-hair theorem

All initially contracting, spatially homogeneous, orthogonal Bianchi Type I-VIII cosmologies and all Bianchi type IX universes sourced by an ultra-stiff fluid with an equation of state such that $(\gamma - 2)$ is positive definite, collapse into an isotropic singularity, where the sink is a spatially flat and isotropic FRW universe.

[J.E.Lidsey, CQG, 23, 3517,\(2005\)](#)

- 1 Introduction
- 2 In the contracting phase
- 3 Ekpyrosis meets anisotropic pressures!**
- 4 The Bianchi IX universe
- 5 Solving the flatness problem within the framework of bouncing cosmologies
- 6 Adding a cosmological constant to the cocktail
- 7 Conclusions

Why include anisotropic stress?



- Bouncing models of the universe, such as ekpyrotic scenarios or LQC models claim isotropisation occurs at early times. But this isn't true on addition of anisotropic stress.
- Interaction rates of particles

$$\Gamma = \sigma n v \sim g \alpha^2 T$$

- To remain in equilibrium, $\Gamma > H$
- Before isotropisation, anisotropic universe expands faster
- Harder to maintain equilibrium

Decoupled collisionless particles free stream and exert anisotropic stresses.

Anisotropic stresses in a Bianchi I universe



We go back to our simple flat anisotropic universe and add anisotropic pressures in.

- Friedmann equation

$$3H^2 = \sigma^2 + \rho_{matter},$$

- The shear evolves as,

$$\dot{\sigma}_{\alpha\beta} + 3H\sigma_{\alpha\beta} = \boxed{\mu\mathcal{P}_{\alpha\beta}}$$

anisotropic stress

The equation for the shear isn't homogeneous and we can't say straight away that an ultra stiff field will be able to dominate over it.

Anisotropic stresses in Bianchi Class A



- Resort to the expansion normalised variables and introduce $Z \equiv \frac{\mu}{3H^2}$ where μ is the anisotropic pressure field energy density with EOS, $p_i = (\gamma_i - 1)\mu$ and $\gamma_\star = (\gamma_1 + \gamma_2 + \gamma_3)/3 > \gamma$
- try to perform stability analysis on the state vector $\mathbf{x} = \{H, \Sigma_+, \Sigma_-, N_1, N_2, N_3, \Omega, Z\}$
- Linearise expansion normalised EFE around the FL point

$$\Sigma_+ = 0, \Sigma_- = 0, N_1 = 0, N_2 = 0, N_3 = 0, \Omega = 1, Z = 0$$

Stability analysis with anisotropic pressures: the results



We find the following eigenvalues

- $\frac{3}{2}(2 - \gamma)$ of multiplicity 2
- $\frac{3\gamma-2}{2}$ of multiplicity 3
- $3(\gamma - \gamma_*)$ of multiplicity 1
- Using the condition $\gamma_* > \gamma > 2$, FL equilibrium point stability cannot be determined

Stability analysis with anisotropic pressures: the results



We find the following eigenvalues

- $\frac{3}{2}(2 - \gamma)$ of multiplicity 2
- $\frac{3\gamma - 2}{2}$ of multiplicity 3
- $3(\gamma - \gamma_*)$ of multiplicity 1
- Using the condition $\gamma_* > \gamma > 2$, FL equilibrium point stability cannot be determined

We can no longer determine the stability of the FL point and can't prove a no hair theorem like before.

- 1 Introduction
- 2 In the contracting phase
- 3 Ekpyrosis meets anisotropic pressures!
- 4 The Bianchi IX universe**
- 5 Solving the flatness problem within the framework of bouncing cosmologies
- 6 Adding a cosmological constant to the cocktail
- 7 Conclusions

Bianchi Type IX: What it is and why we use it



- It is the most general closed homogeneous universe, describable by ODEs
- It has the closed FRW universe as its isotropic sub-case
- It has expansion anisotropy and anisotropic 3-curvature (which has no Newtonian analogue)
- The 3-curvature can change sign through the course of its evolution and is positive when the model is closest to isotropy.
- On approach to $t \rightarrow 0$, in an open interval $0 < t < T$, exhibits chaotic Mixmaster oscillations, however oscillations become finite in number even if $t \rightarrow t_{PI}$ on the finite interval $t_{PI} < t < T$ excluding $t \rightarrow 0$.

We have a Bianchi Type IX universe with




- an isotropic pressure field with energy density ρ which follows the equations of state $p = (\gamma - 1)\rho$ and is effectively NEC violating, to bring about a non-singular bounce
- Anisotropic pressure field with energy density μ and $p_i = (\gamma_i - 1)\mu$ with $i = 1, 2, 3$, such that $\gamma_\star = (\gamma_1 + \gamma_2 + \gamma_3)/3$ and $\gamma_\star > \gamma$
- Choose initial conditions satisfying the Friedmann constraint

Scale factor evolution



Figure: Scale factors with isotropic ghost field and with fields with anisotropic pressures respectively




- The scale factors with just an isotropic pressure ghost field **bounce** and start to expand.
- The scale factors with the anisotropic pressure field included seem to contract towards a singularity.



Evolution of the shear

- ☞ If we look at the evolution of the shear, we find,

Figure: Evolution of σ^2 with time



- 1 Introduction
- 2 In the contracting phase
- 3 Ekpyrosis meets anisotropic pressures!
- 4 The Bianchi IX universe
- 5 Solving the flatness problem within the framework of bouncing cosmologies**
- 6 Adding a cosmological constant to the cocktail
- 7 Conclusions

Bouncing cosmologies and the flatness problem



- Simple models of bouncing universes such as matter+ radiation closed FRW incorporated increasing radiation entropy to increase expansion maxima from cycle to cycle
- Universe seemed to **approach flatness**
- Suitable candidate for the current day universe?


Question

Would an anisotropic, bouncing cosmological model under similar increasing radiation entropy from cycle to cycle undergo isotropisation simultaneously with approach to flatness?

Present day flatness can perhaps be achieved by diluting the curvature with increasing volume



Figure: Scale factor with increasing entropy of radiation in closed FRW




J.D.Barrow, M.P.Dabrowski, MNRAS, 275, 850 – 862, 1995

The scale factors with increasing radiation entropy



Increasing entropy of radiation in Bianchi IX


Figure: Evolution of volume scale factor and individual scale factors respectively



Let's see how the shear and the 3-curvature behave



Figure: Evolution of σ^2 and ${}^{(3)}R$ respectively




- 1 Introduction
- 2 In the contracting phase
- 3 Ekpyrosis meets anisotropic pressures!
- 4 The Bianchi IX universe
- 5 Solving the flatness problem within the framework of bouncing cosmologies
- 6 Adding a cosmological constant to the cocktail**
- 7 Conclusions

The scale factors with increasing radiation entropy

- The volume scale factor and hence the individual scale factors evolve through a series of oscillations with increasing maxima until the cosmological constant starts to dominate and they expand exponentially


Figure: Evolution of volume scale factor and individual Hubble rates from left to right



Let's see how the shear and the 3-curvature behave



Figure: Evolution of σ^2 and ${}^{(3)}R$ respectively



- 1 Introduction
- 2 In the contracting phase
- 3 Ekpyrosis meets anisotropic pressures!
- 4 The Bianchi IX universe
- 5 Solving the flatness problem within the framework of bouncing cosmologies
- 6 Adding a cosmological constant to the cocktail
- 7 Conclusions**

Summary I



- In the initially contracting Bianchi Class A models, in the presence of ultra-stiff anisotropic stresses, FL is no longer an attractor in the asymptotic past
- In the Bianchi IX equations, including an ultra stiff anisotropic pressure field causes the scale factors to **contract towards a collapse near the singularity**.
They bounce with only an isotropic ghost field present.
- The shear remains small and nearly constant in the isotropic case but increases without bound when the anisotropic pressure field is included.

Summary II



- By future evolving the model, we find that with radiation entropy increase, the height of the **scale factor maxima increases**, but the shear and the curvature oscillate and **do not decrease** to indicate isotropisation at any time.
- On adding the cosmological constant to the analysis, at the point of cosmological constant domination, the scale factors stop oscillating and undergo exponential expansion.
- The shear and the curvature tensors oscillate as before and then **under cosmological constant domination, they fall to smaller and smaller values**

So the takeaway message...



Near the singularity...

Including anisotropic stress, does not always result in isotropisation near the singularity, even if the anisotropic stress field is ultra-stiff on average


On future-evolving the system..

On evolving the system into the future, isotropisation does not occur as the shear keeps oscillating with the oscillations of the volume scale factors. On adding a cosmological constant, the shear and curvature fall to very small values

The effect of non comoving velocities with entropy increase




Figure: Evolution of the square of one of the spatial velocity components



The effect of non comoving velocities after cosmological constant domination

Figure: Evolution of the square of one of the spatial velocity components



The effect of non comoving velocities, in brief



- On imposing momentum and angular momentum conservation, the spatial components of the velocities fall to smaller values with an increase in entropy density and vice versa
- On addition of cosmological constant, bounces cease, expansion tends to the quasi dS asymptote and velocities tend to oscillate with a constant amplitude, while one of them tends to a constant value.

References



- CG and John D. Barrow, 2017, arXiv:1710.00747
- John D. Barrow and CG, 2017, arXiv:1705.06647 (Accepted in Int. Journal Mod. Phys. D)
- John D. Barrow and CG, 2017, Physical Review D **95**, 083515
- John D. Barrow and CG, 2015, Class. Quantum Grav., Vol. 33, No. 12
- Dynamical Systems in Cosmology, edited by J. Wainwright and G.F.R. Ellis, Cambridge University Press
- V.A. Belinskii, I.M. Khalatnikov and E.M. Lifshitz, 1970, Adv. Phys. Vol. 19, No. 525
- V.A. Belinskii and I.M. Khalatnikov, 1972, Zh. Eksp. Teor. Fiz Vol. 63, No. 1121
- A.G. Doroshkevich, V.N. Lukash and I.D. Novikov, 1973, Zh. Eksper. Teor. Fiz, 64, No. 1457 – 1474