Title: Topological line operators and renormalization group flows in two dimensions

Date: Mar 06, 2018 02:30 PM

URL: http://pirsa.org/18030071

Abstract: Topological defect operators are extended operators in a quantum field theory (QFT) whose correlation functions are independent of continuous changes of the ambient space. They satisfy nontrivial fusion relations and put nontrivial constraints on the QFT itself and its deformations (such as renormalization group (RG) flows). Canonical examples of topological defect operators include generators of (higher-form) global symmetries whose constraints on the QFT have been well-studied. However they constitute just a tip of a much richer framework which we explore in detail in two spacetime dimensions.

More specifically, we study the crossing relations of general topological defect lines (TDL) in two dimensions, discuss their relation to the usual 't Hooft anomalies associated to global symmetries, and use them to constrain RG flows to either conformal fixed points or topological quantum field theories (TQFTs). We show that if certain non-invertible TDLs are preserved along a RG flow, then the vacuum cannot be a non-degenerate gapped state. For various massive flows, we determine the infrared TQFTs completely from the consideration of TDLs together with modular invariance.

Topological Line Operators & RG flows in 2d

arXiv: 1802.04445

C.M. Chang, Y-H. Lin, S-H. Shao, Y. Wu & X. Yin


Topological Line Operators & RG flows in 2d
arXiv: 1802.04445
C.M. Chang, Y-H. Lin, S-H. Shao, Y. Wu & X. Yin
extended QFT
RG flow
IR phase?
constrain
defect objects/ops
defects inherited

2d TDL commute with L_m, $\bar{L}_m$

- symmetry generator : $e^{S^{*j}}$

↓ generalize?

linear map on space of local ops

↓ consistency? $\lambda \Rightarrow \exists\, \mathcal{H}_\lambda$, modular inv.



$$\Rightarrow \mathrm{tr}\left(\lambda\, q^{L_0-\frac{c}{24}}\bar{q}^{\bar{L}_0-\frac{c}{24}}\right) = \mathrm{tr}_{\mathcal{H}_\lambda}\left(\tilde{q}^{L_0-\frac{c}{24}}\tilde{\bar{q}}^{\bar{L}_0-\frac{c}{24}}\right)$$


$\mathcal{L}$ = $\mathcal{L}$ $\Rightarrow$ $\mathrm{tr}\left(\mathcal{L}\, q^{L_0-\frac{c}{24}}\, \bar{q}^{\bar{L}_0-\frac{c}{24}}\right) = \mathrm{tr}_{\mathcal{H}_L}\left(\bar{q}^{L_0-\frac{c}{24}}\, \bar{q}^{\bar{L}_0-\frac{c}{24}}\right)$
· 2d RCFTs w/ diagonal modular inv. $\{V_i\}$ ← chiral primaries
$\hat{\mathcal{L}}_k |V_i\rangle = \frac{S_{ki}}{S_{0i}} |V_i\rangle$
$\mathcal{L}_i \longleftrightarrow V_i$ (verlinde ops)
V.F. $\Rightarrow$ $\hat{\mathcal{L}}_i \hat{\mathcal{L}}_j = \sum_k N_{ij}^k \hat{\mathcal{L}}_k$





$$\Rightarrow \quad \mathrm{tr}\left(\mathcal{L}\, q^{L_0-\frac{c}{24}}\, \bar{q}^{\bar{L}_0-\frac{c}{24}}\right) = \mathrm{tr}_{\mathcal{H}_L}\left(\tilde{q}^{L_0-\frac{c}{24}}\, \tilde{\bar{q}}^{\bar{L}_0-\frac{c}{24}}\right)$$

- 2d RCFTs w/ diagonal modular inv. $\{V_i\}$ ← chiral primaries

$$\hat{\mathcal{L}}_k |V_i\rangle = \frac{S_{ki}}{S_{0i}} |V_i\rangle \qquad \mathcal{L}_i \longleftrightarrow V_i \quad \text{(verlinde ops)}$$

V.F. $\Rightarrow$ $\hat{\mathcal{L}}_i \hat{\mathcal{L}}_j = \sum_k N_{ij}^k \hat{\mathcal{L}}_k$

- 2d non-Verlinde line ops in 2d CFTs (e.g. WZW G. $\partial_z j_{\bar{z}} = 0$)


junctions
L_1
L_2
L_3
$V_{L_1L_2L_3}$
$\mathcal{H}_{L_1L_2L_3}$
(0,0)
Fusion
L_1
L_2
L_k
local


locality
Modular Inv
mess
=
=


Modular Inv
mess
$\tau^h \bar{\tau}^{\bar{h}}$

Consequences.



$\in \underline{V_{1234}} \subset \mathcal{H}_{1234}$ $\Rightarrow$ $\bar{z}$



crossing kernel

$$\underline{K^{L_1 L_4}_{L_2 L_3}(L_5, L_6)}$$


Pentagon
$k\,k = k\,k\,k$


$\{1, X\}$
$X^2 = 1 + nX \quad n \in \mathbb{Z}^+$
$\exists K$
$n = 0, 1$
$\{1, X, Y\}$
$\leq$ possibilities

Ising $c = \frac{1}{2}$ $\mathcal{M}(3,4)$

	$1_{0,0}$	$\xi_{\frac{1}{2},\frac{1}{2}}$	$\sigma_{\frac{1}{16},\frac{1}{16}}$	
η	1	1	-1	← $\mathbb{Z}_2$ sym
$\hat{N}$	$\sqrt{2}$	$-\sqrt{2}$	0	← verlinde line

$\eta N = N\eta = \eta$, $N^2 = 1 + \eta$ ← fusion



$\mathcal{H}_\eta = \{1_{\frac{1}{2},0}, \hat{\psi}_{0,\frac{1}{2}}, \mathcal{M}_{\frac{1}{16},\frac{1}{16}}\}$, $\mathcal{H}_N = \{\sigma_{\frac{1}{16},0}, \xi_{0,\frac{1}{16}}, \Lambda_{\frac{1}{16},\frac{1}{2}}, \tilde{\Lambda}_{\frac{1}{2},\frac{1}{16}}\}$ ← defect Hilbert space


Constraints on RG flows
ϕ relevant def.
$\hat{\mathcal{L}}|\phi\rangle = \langle\hat{\mathcal{L}}\rangle|\phi\rangle$
$\Rightarrow$ $\mathcal{L}$ preserved by ϕ

Constraints on RG flows

ϕ relevant def.

$\hat{\mathcal{L}}|\phi\rangle = \langle \hat{\mathcal{L}} \rangle |\phi\rangle$ $\Rightarrow$ $\mathcal{L}$ preserved by ϕ



x


crossing
constrain
spin selection rule
Thm 1: if ∃ spin selection rule that forbids s=0 in
& preserved by RG
empty
empty
IR phase is a CFT satisfying the spin selection rule

Obs:

In the case when IR phase is gapped

$\Rightarrow$ all structure const determined by TDL


spin selection rule that forbids s=0 in $\mathcal{H}_2$
& 2 preserved by RG
$\mathcal{H}_2^{IR}$ empty
$\mathcal{H}_2^{IR}$ empty → IR phase is a CFT satisfying the spin selection rule
M(4,5) tricritical Ising $c = \frac{7}{10}$
RG: 2, σ'
$W^2 = 1 + W$ (LY)
$N^2 = 1 + \eta$ (Ising)
$\mathbb{Z}_2$ preserving
σ'
simple lines
$\frac{1}{10}$ $\frac{3}{5}$ $\frac{5}{2}$ $\frac{3}{80}$ $\frac{7}{16}$
1, η, W, N, ...
$\mathbb{Z}_2 \to \eta$
$\hat{W}$
$\hat{N}$
$\sqrt{2}$
$\xi = \frac{1+\sqrt{5}}{2}$

- $\mathcal{H}^{IR}$ ⊥ empty
- $\mathcal{H}^{IR}$ nonempty ⟶ IR phase is a CFT satisfying the spin selection rule

$M(4,5)$ tricritical Ising $c = \frac{7}{10}$

RG: ε, σ'

$\mathcal{H}_W$: $s \in \pm\frac{3}{5}, 0$

$\mathcal{H}_N$: $s \in \frac{2}{3} \pm \frac{1}{16}$

	1	ε	ε'	ε''	σ	σ'
	$\frac{1}{10}$	$\frac{3}{5}$	$\frac{3}{2}$		$\frac{3}{80}$	$\frac{7}{16}$
simple lines:	1	η	W	N	$\cdots$	$\zeta = \frac{1+\sqrt{5}}{2}$
$\mathbb{Z}_2 \to \eta$	1	1	1	1	−1	−1
W	ζ	$-\zeta^{-1}$	$-\zeta^{-1}$	ζ	$-\zeta^{-1}$	ζ
$\hat{N}$	$\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	0	0

$W^2 = 1 + W$ (LY)

σ' preserves

1 phase

$N^2 = 1 + \eta$ (Ising)

ε

2 phases

$\langle W \rangle = \zeta$

$\langle N \rangle = \sqrt{2}$


$\mathcal{H}^{IR}_{\mathbb{Z}_2}$ empty
$\mathcal{H}^{IR}_{\mathbb{Z}_2}$ nonempty → IR phase is a CFT satisfying the spin selection rule
$M(7,8)$ tricritical ising $c = \frac{7}{10}$
RG: ε, σ'
$\mathcal{H}_W$: $\gamma \in \{\frac{3}{5}, \frac{7}{5}, \frac{1}{10}\}$
$\mathcal{H}_N$: $\gamma \in \frac{3}{2} \pm \frac{1}{16}$
$W^2 = 1 + W$ (LY)
$N^2 = 1 + \eta$ (Ising)
$\langle W \rangle = \{$
$\langle N \rangle = \sqrt{2}$
σ' preserves
ε
2 phases
1 phase
$\langle W \rangle = \frac{\sqrt{5}+1}{2}$
$\mathrm{tr}(W_s) \in \mathbb{Z}^{\geq 0} \Rightarrow \{\frac{\sqrt{5}+1}{2}, \frac{1-\sqrt{5}}{2}$
simple lines: $1, \eta, W, N, \ldots$
$\{ = \frac{1+\sqrt{5}}{2}$
$\mathbb{Z}_2 \to \eta$
W
N
5