Title: The Master Chiral Algebra

Date: Mar 23, 2018 02:00 PM

URL: http://pirsa.org/18030089

Abstract:


KL(G)
FLE
Loc
Poinc


KL(G)
FLE
Whit(Gr)
Loc
Poinc
Whitk(GrG)
FLE


KL(G)
FLE
Whit(Gr)
Loc
Poinc
0 → C → ĝ_k → g((A)) → 0
g[[t]]
KL_k(G) = ĝ_{k-vol}(G[[t]])
FLE


Poinc
level_k
G((t))
G[[t]]
G[[t]]


Whit
Poinc
Dmod(Bun)
Γ
0 → C → ĝ_K → g((t)) → 0
g[[t]]
KL_K(G) = ĝ_K-mod^G[[t]]
level_K
G((t))
Bun_G
G[[t]]
Bun_G
Γ : Dmod_K(Bun_G^level)^G[[t]] → ĝ_K-mod^G[[t]]
D-mod(Bun_G)


Bun_G
level_x
$G(\mathbb{A})$
$G[\mathbb{A}]$
Bun_G
$\Gamma : \mathrm{D\text{-}M}(\mathrm{Bun}_G^{\mathrm{level}})$
$G[\mathbb{A}]$
$\hat{a}_{x\text{-loc}}$
$G(\mathbb{A})$
$\mathrm{D\text{-}M}(\mathrm{Bun}_G)$
$M \in \hat{a}_{x\text{-loc}}$
$\mathrm{Loc}(M)_1 = M_{a_{out}}$
$a_{out} = \Gamma(X - x, 0) \otimes y$


$\mathbb{C}[G]$
$V \in \hat{a}_{y_k\text{-loc}}^{\mathbb{C}[G]}$
$\| \hat{a}_{y_k}$
$\mathrm{Ind}_{y\mathbb{C}[G]\otimes\mathbb{C}}^{(G)}$
$\mathrm{loc}(V) = \mathcal{D}_K$
$a_{y_{out}}$
y
$\mathrm{Dmod}_K(G\backslash G)$
$\left(\mathrm{Dmod}_K(G\backslash G)\right)_{N(\theta,\chi)}$


C


Dmod_κ(GrG)
(Dmod_κ(GrG))^{W(G,S)}
C
Whit_κ(GrG)
GrG
BunG


$\mathrm{Dmod}_\kappa(\mathrm{Gr}_G)$
$\mathrm{Whit}_\kappa(\mathrm{Gr}_G)$
Gr_G
Bun_G
$\pi_* \cdot \mathrm{Dmod}_\kappa(\mathrm{Gr}_G) \to \mathrm{Dmod}(\mathrm{Bun}_G)$


C
$G_F G$
Bun_G
$\pi_* \cdot D_{mod_K}(G_F G) \to D_{mod}(\mathrm{Bun}_G)$
$\mathcal{F} \in D_{mod_K}(G_F G)$
$N_\alpha \subseteq N((t))$
$\pi_* \left(Av_!^{N_\alpha, \psi}(\mathcal{F}) \right) \in D_{mod_\alpha}(\mathrm{Bun}_G)$

$\overline{\mathcal{F}} \in \mathrm{Dmod}(G_F \backslash G)$

$N_\alpha \subseteq N((t))$

$\pi_* \left(\mathrm{Av}_*^{N_\alpha, \psi}(\overline{\mathcal{F}}) \right) \in \mathrm{Dmod}_\alpha(\mathrm{Bun}_G)$

$\mathrm{Dmod}(\mathrm{Bun}_G)$

Lemma For any $\overline{\mathcal{F}}$ (with finite support)

the sequence $\alpha \longrightarrow \pi_* \left(\mathrm{Av}_*^{N_\alpha, \psi}(\overline{\mathcal{F}}) \right)$ will stabilize


$\overline{\mathcal{F}} \in \mathrm{Dmod}_\kappa(G\mathrm{Gr}_G)$
$N_\alpha \subseteq N((t))$
$\pi_* \left(A\mathrm{V}_\alpha^{N_\alpha \cdot \infty}(\overline{\mathcal{F}}) \right) \in \mathrm{Dmod}_\kappa(\mathrm{Bun}_G)$
Lemma For any $\overline{\mathcal{F}}$ (with finite support)
the sequence
$\alpha \longrightarrow \pi_* \left(A\mathrm{V}_\alpha^{N_\alpha \cdot \infty}(\overline{\mathcal{F}}) \right)$ will stabilize
The resulting functor
$\mathrm{Dmod}_\kappa(G\mathrm{Gr}_G) \longrightarrow \mathrm{Dmod}_\kappa(\mathrm{Bun}_G)$
is $(N((t)), \delta)$ invariant $\Rightarrow$ gives rise Poinc.


KL(G)
FLE
Whit(Gr_G^∨)
Loc
Poinc
Whit(GrG)
FLE
λ-dominant weight for G
V^λ
G[[t]] ⟶ G ↷ V^λ


Whit
Gr
Poinc
Dmod
Bun
Γ
λ-dominant weight for G ⟺ dominant coweight for
Orbits of
on
are in bijection with
Whit


dominant coweight for $\check{G}$
Orbits of $\check{N}((t))$ on $Gr_{\check{G}}$
are in bijection with cowelnt
$\delta_\lambda \in Dmod_{+\kappa}(Gr_{\check{G}})$
$Whit_{-\kappa}(Gr_{\check{G}})$
$V^\lambda \longmapsto \delta_\lambda$
$KL(G)_\kappa \otimes KL(\check{G})_{\check{\kappa}} \to$
W-mod
Vect


δ_λ
$KL(G)_K \otimes KL(\check{G})_{\check{K}}$
$i\mathcal{N}\otimes FLE$
$KL(G)_K \otimes Whit(G_{FG})_{-K}$
W_{-w_0}
Verl
$KL(G)_K \otimes KL(\check{G})_{\check{K}}$
W_{-w_0}
Verl
$D_{mod_K}(G_{FG})$
$\left(D_{mod_K}(G_{FG})\right)_{(N(\mathbb{Q}),\chi)}$
$Whit_K(G_{FG})$
$\mathbb{C}$


KL(G)_κ ⊗ KL(Ǧ)_κ
id ⊗ FLE
KL(G)_κ ⊗ Whit(Gr_G)_κ
W-mod
Vect
KL(G)_κ ⊗ D-mod_{-κ}(Gr_G)
ω̃_{κ-mod}
Gr_G
Bun_G
Vect


KL(Ǧ)
id⊗FLE
Whit(GrG)
Vect
KL(G) ⊗ Dmod
(GrG)
DS
(h(t), n)
Vect
GrG
Bun
Vect


$KL_k(G)$
$KL_{-k}(G) \to V_{ect}$
$M_1, M_2 \to H^2(\mathrm{gr}(A), M_1 \otimes M_2)$
$KL_n(G) \otimes KL_k(\bar{G}) \to V_{ect}$
P_{glob}
Lemma
$P_{glob} = H_{dR}(W, P_{loc})$
time let Λ be a chiral
algebra with N-symmetry
$H_{dR}(X, B\Sigma\pi(n, A))$
$\cong$
$H_{dR}(B\mathrm{Sym}_N, \mathrm{Loc}(A))$


FLE
$Whit_{-\check{\kappa}}(G_{\mathrm{gr}}\check{G})$
Poinc
$D_{\mathrm{mod}}(Bun_{\check{G}})$
$-\check{\kappa}_{\mathrm{ny}}$
Γ
$KL_{-\check{\kappa}}(\check{G})$
FLE
$CD^0(G)_{\kappa,-\kappa} \in KL_\kappa(G) \otimes KL_{-\kappa}(G)$
$\mathrm{loc}_{G\times G}(CD^0(G)_{\kappa,-\kappa}) \in D_{\mathrm{mod}}{}_{\kappa,-\kappa}(Bun_G \times Bun_G)$
$CD^0(G)_{\kappa,-\kappa} =$


FLE
Whit
Poinc
Dmod
Γ
FL
FLE
CDG(G)
hoc
CDG(G)
Fun


Lemma
KL(G)_κ ⊗


Lemma


$KL_u(G) \otimes KL_{-u}(G) \otimes$
$KL(\check{G})_u \otimes KL(\check{G})_{\check{u}}$
$Id \otimes$
$p^{loc} \circ Id$
$KL_u(G)$
$KL(\check{G})_{\check{u}}$
$KL(G)_u \otimes KL(\check{G})_{\check{u}}$
Explicitly:
$M = \bigoplus_{\lambda, \check{\lambda}}$
$T^{\lambda, \check{\lambda}} = DS_{\chi^{\check{\lambda}}}(V^{\lambda})$
time let Λ' be a chiral
algebra with N-symmetry
$= H_{dR}(Bun_N, Loc(A))$


Explicitly:


Explicitly:




Lemma
M-mod
G((t))
Ǧ((t))
the functor


M-mod
M-mod
Dmod( )


M-mod
G((t))
Ǧ((t))
M-mod
Dmod(Fl)
C ⟷ Č
C^I ≃ Č^I