Title: Jocelyn Bell Burnell: University of Oxford

Date: Oct 25, 2018 07:00 PM

URL: http://pirsa.org/18100095

Abstract: ocelyn Bell Burnell, winner of the 2018 Special Breakthrough Prize in Fundamental Physics, is an accomplished scientist and champion for women in physics. As a graduate student in 1967, she co-discovered pulsars, a breakthrough widely considered one of the most important scientific advances of the 20th century. When the discovery of pulsars was recognized with the 1974 Nobel Prize in Physics, the award went to her graduate advisor. Undaunted, she persevered and became one of the most prominent researchers in her field and an advocate for women and other under-represented groups in physics.

She plans to use the \$3 million Breakthrough Prize to fund women and other under-represented groups pursuing physics to bring greater diversity to the field.


Pirsa: 18100095 Page 1/45

What is that? The discovery of pulsars - a graduate student's story

Jocelyn Bell Burnell

University of Oxford
Astrophysics
and
Mansfield College






Pirsa: 18100095 Page 2/45

Topics

- Introduction to radio astronomy
- Quasars
- Cambridge (UK) and the discovery of pulsars
- What are pulsars?



The electromagnetic spectrum



Radio astronomy




Page 5/45

Radio astronomy – quasars (QSR)

- Radio astronomy developed post WWII
- Developed by ex-radar 'boffins'; using the radar receivers (not transmitting).
- Found an undreamt-of richness of radio emitting objects in the sky.
- Some starlike-ish quasi-stellar
- Strong sources but VERY far (?)
- Called 'quasars'.




My path - some British geography...



Some British geography 2





Imposter Syndrome

- Everyone here is very clever!
- I'm not.....
- They've made a mistake admitting me.....
- They're going to discover their mistake....
- And throw me out......
- (I'd better leave before they throw me out)



Yakov Zeldovich (1914 – 87)

(Said of cosmologists.....!)

'Frequently in error, but never in doubt.'



Page 10/45

On arrival in Cambridge 1965



All new radio
 astronomy graduate
 students were given a
 set of tools......

Pirsa: 18100095 Page 11/45

First build your radio telescope (through hail, rain and sunshine)



Early 1967
Don Rolph + JBB
Checking impedance
of air-spaced cables

Area = 57 tennis courts. Grant £12k. 6 people worked for 2 years to build it.

Pirsa: 18100095 Page 12/45

The 4.5 acre* radio telescope (*1.8 hectares), looking West




2048 81.5MHz λ/2 antennae (16 E-W rows of 64 + 64), 1000+ wooden posts, 192 km wire and cable. Phased interferometric array. Valve receiver!

Pirsa: 18100095 Page 13/45

Interplanetary scintillation

- Compact radio source scintillates, extended does not.
- Quasars (quasi-stellar radio sources) are compact; other radio sources are not



Pirsa: 18100095 Page 14/45

Scintillation – a rough analogy



Swimming pool

- Swimming pool patterns of light
- Caustic network produced by ripples
- Pattern moves as wind blows ripples

Quasar Hunt!

- Only about 20 quasars known. Project was to find more, and measure their size, by repeated mapping of the sky.
- Pick quasars out by their rapid 'twinkling' interplanetary scintillation; use short time
 constant/short integration time. τ = 0.1 s.
- 6 months' observing
- Found c 180 more quasars



Data analysis

No computer!


3-pen chart paper

- 30m / day
- 120m/sky scan
- 5.3km total




Discovery of pulsars

- Occasionally 0.5 cm in the 120 m showed an unusual signal.
- 'Occasionally' = 20%
 of the occasions that
 part of sky observed
- This signal occupied approx 10 parts in a million of the chart paper!



High-speed recording



Top line - pulsed emission detected Bottom line - one second time pulses

What is it?

- Local radio interference? No keeps sidereal time.
- Faulty equipment? No seen by a separate telescope and receiver.
- Small (short pulses) and big (keeps accurate pulse period). ???????



What is it (contd)?

- Dispersion measurement showed it was 200 light years away (i.e. beyond the Solar System, but nearby in the Milky Way)
- Little Green Men? They'd be on a planet orbiting their Sun; no Doppler effect
- Finding second, third and fourth (1133, 0833, 0950)


The naming of pulsars

- Interviewed by Science Correspondent of The Daily Telegraph – Anthony Michaelis – shortly after the discovery
- What were we going to call them?
- He suggested pulsar cf quasar

The Daily Telegraph

What are pulsars? Overview

Pulsars (pulsating radio stars) model



- Star spins like a lighthouse, sweeping radio beam around the sky.
- We see a pulse each time beam sweeps across us.

Page 24/45

Overview

- Pulsars are small, very dense neutron-rich stars (neutron stars).
- Strong gravitational fields; very strong E and M fields); rapidly spinning.
- Probably formed in the explosion (supernova) that ends the life of a massive (>10M_{Sun}) star
- Primarily observed as pulsed, radio-emitting objects

Pulsar population

- Seen in RADIO, (visible), X-rays and gamma rays
- About 2700 known; ~20 visible; ~100 X-ray;
 ~200 gamma ray
- Of the 2700; ~10% in binaries (one PSR-PSR); 1 triple system; a few with planets
- About 10⁵ in the Galaxy (maybe)




Large Magellanic Cloud – SN1987a

When a large star (e.g. 10x bigger than the Sun) runs out of fuel the core of star collapses and becomes a neutron star. The outer 90 – 95% of star ejected giving a spectacular 'supernova' explosion





Crab Nebula ~ 1000 yrs after SN



- Collapsed core is a neutron star or pulsar
- This nebula emits synchrotron radiation – kept energised by pulsar in its centre

PROPERTIES OF PULSARS

In more detail

Pirsa: 18100095 Page 29/45

- Mass ≈ few x 10²⁷ tonnes, i.e. few thousand million million million tonnes (1.3 – 2 times mass of Sun)
- Radius ~10 km
- : average density comparable with density of nucleus of an atom
- Rich in neutrons: 'neutron star'



Properties of Pulsars - density



- A thimble-full of pulsar material has the same mass as the population of the Earth
- Strong surface gravity
- Tidal effects
- Very condensed matter physics

Pirsa: 18100095 Page 31/45

Gravity bends light – one can see 20° –
 30° over the horizon on a neutron star

Gravity red shifts light



 Gravity affects clocks – clocks tick two times slower than here on Earth




Pirsa: 18100095 Page 32/45


 The gradient of the gravitational force is strong – tidal disruption of bodies approaching pulsar



- The gradient of the gravitational force is strong – tidal disruption of bodies approaching pulsar
- Warning! Visiting a pulsar can seriously damage your health!



- Magnetic field of pulsar ~ 100 million Tesla
- (Frig magnet ~ 0.01T, strong lab magnet ~ 10T.)
- Voltage drops of up to 10 billion Volt/cm
- So electromagnetic forces are 100 billion times stronger than gravitational!



- Accurate timekeepers good to 1 part in 10000 million million (10¹⁶); the period of a typical pulsar has increased by about 1 second since the age of the dinosaurs!
- This means they can be used as clocks to test Einstein's theory of Relativity.

Things to watch over the next few years






FAST radio telescope – 500 m diameter, in S China. Now being commissioned – the world's largest. As of mid-August 2018, about 70 new PSRs. And some interesting/weird sources. (!?)

Pirsa: 18100095 Page 37/45

CHIME – Canada's new radio telescope



- Just starting operation
- Will be good for studying pulsars


SOME PULSAR RECORDS

 The pressure at the centre of a pulsar/neutron star is

times the atmospheric pressure here on Earth

Pirsa: 18100095 Page 41/45

 The first planets discovered beyond the Solar System are three planets around the pulsar PSR1257+12.
 There is also a tiny fourth planet or asteroid in the system.



• The roundest known thing in the universe is the orbit of the pulsar PSRJ1909- 3744 around another star. It is round to 5 μ m (one tenth the width of a human hair) in 567000km. (Eccentricity e = 1.30 x 10⁻⁷)

Pirsa: 18100095 Page 43/45

 If something falls on to the surface of a pulsar, it will hit the surface with a speed of 0.5c

Pirsa: 18100095 Page 44/45

The End

