Title: PSI 2018/2019 - Machine Learning - Lecture 7

Speakers: Lauren Hayward Sierens

Collection: PSI 2018/2019 - Machine Learning (Hayward Sierens)

Date: April 02, 2019 - 9:00 AM

URL: http://pirsa.org/19040002

TOMORROW'S LECTURE: 10:15am

Outline for today: Unsupervised Learning (UL)

- examples of UL tasks
- relationship to SL
- dimensional reduction
  - ↳ principal component analysis (PCA)

Unsupervised Learning (UL):

Goal: learn structural properties (features) of unlabelled data

Dataset $\mathcal{D} = \{\vec{x}\}$

↳ each datapoint has N components $\vec{x} = (x_1, x_2, \cdots, x_N)$

↳ dataset has M points: $\mathcal{D} = \{\vec{x}_1, \cdots \vec{x}_M\}$

Within this class, we will study two classes of UL algorithms:

1. Dimensional reduction
   - ↳ today & tomorrow

Idea: reduce the dimensionality of the dataset by finding the most important features

$N \rightarrow N' < N$

If $N' \leq 3$ then we can visualize the data.

In many cases the dataset might naturally form "clusters".

eg) Ising spin configs at diff temperatures

↳ we know magnetization is an important feature



② Generative modelling

↳ next week

Idea: use the datapoints $\vec{x}_1, \vec{x}_2, \ldots \vec{x}_M$ to learn about the underlying prob. dist $p(\vec{x})$ and generate new samples from it.


Relationship between SL and UL
UL: $\mathcal{D} = \{\vec{x}\} \longrightarrow$ learn $p(\vec{x})$
SL: $\mathcal{D} = \{(\vec{x}, \vec{y})\} \longrightarrow$ learn conditional prob dist $p(\vec{y}|\vec{x})$

$p(\vec{x})$

conditional dist $p(\vec{y}|\vec{x})$

UL and SL are not formally different due to the chain rule of prob.:

For $\vec{x} = (x_1, x_2, \ldots, x_N)$

$$p(\vec{x}) = p(x_N | x_1, x_2 \ldots x_{N-1})\, p(x_1, x_2 \ldots x_{N-1})$$

$$= \ldots$$

$$= \left[\prod_{i=2}^{N} p(x_i | x_1, \ldots, x_{i-1}) \right] p(x_1)$$

$\Rightarrow$ We can express a UL problem as
N-1 SL problems along with
the easier problem of learning $p(x_1)$

eg) for indep. spin configs. with $\mathbb{Z}_2$ symm.:

$$p(x) = p(\uparrow) = p(\downarrow) = \frac{1}{2}$$

We can also express a SL problem in terms of UL problems using:

$$p(\vec{y}|\vec{x}) = \frac{p(\vec{x},\vec{y})}{\sum_{\vec{y}'} p(\vec{x},\vec{y}')}$$

Dimensional Reduction using Principal Component Analysis (PCA)

Idea: Generate a lower-dim. representation of the N-dim. dataset $\mathcal{D}$ through a linear transformation.


An example for N=2 where PCA would be useful
X_2
X_1
X_1 and X_2 are highly (linearly) correlated. Knowing X_1 is enough to infer X_2 with high precision.


However, PCA is not always useful.

infer x_2 with high precis

Let us store our M datapoints in an MxN matrix:

$$X = \begin{bmatrix} \vec{x}_1 \\ \vec{x}_2 \\ \vdots \\ \vec{x}_M \end{bmatrix} = \begin{bmatrix} x_{11} & x_{12} & \cdots & x_{1N} \\ x_{21} & x_{22} & \cdots & x_{2N} \\ & & \ddots & \vdots \\ x_{M1} & x_{M2} & \cdots & x_{MN} \end{bmatrix}$$

infer X_2 with high precision.

To perform PCA, the mean of the datapoints must be zero, so first shift the "centre" of each column:

$$\vec{X}_i \longrightarrow \vec{x}_i - \frac{1}{M}\sum_{k=1}^{M} \vec{X}_k$$

$$\left(X_{ij} \longrightarrow X_{ij} - \frac{1}{M}\sum_{k=1}^{M} X_{kj} \quad \forall_j \right)$$

$1 \leq i \leq M, \; 1 \leq j \leq N$

$\cdots \; X_{1N}$

$\cdots \; X_{2N}$

$\cdots \; X_{MN}$

Now consider the $N \times N$ matrix

$$V_X \equiv \frac{1}{M-1} X^T X$$

- diag. elements store the variances of each component of $\vec{x}$
- off-diag. elements store the covariances between pairs of components

Goal of PCA: Find a new representation $X' = XP$ such that $V_{X'}$ is diagonal.

P is an $N \times N$ matrix encoding the linear transformation.

Now note that:

$$V_{X'} = \frac{1}{M-1} X'^{T} X' = \frac{1}{M-1} (XP)^{T} XP = \frac{1}{M-1} P^{T} X^{T} XP = P^{T} V_{X} P$$

When this matrix is diagonal:

$$P^{T} V_{X} P = D = \begin{pmatrix} \lambda_1 & & 0 \\ & \lambda_2 & \\ 0 & & \lambda_N \end{pmatrix}$$

Since $V_X = \frac{1}{M-1} X^T X$: $P^{-1} = P^T$

So PCA amounts to solving an eigenvalue problem

$$V_X = PDP^{-1}$$

where each eigenvalue λ_j is $\frac{1}{M-1} \sum_{i=1}^{M} X_{ij}'^2 \geq 0$

Let us sort the eigenvalues so $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_N$

$V_X^T \Sigma_X V_X = D = \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_N \end{pmatrix}$

Now the first principal component X_1' corresponds to the direction in the space of the original data with highest variance λ_1



Sort the eigenvalues so $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_N$

Let us plot λ_i vs i:



If these eigenvalues quickly become small, the lower-dim. representation $\vec{x}' = (x_1', x_2', \ldots, x_p')$ $(p \ll N)$ corresponding to the first few principal components will be a valid and efficient representation of the data's features.


Tomorrow: non-linear method for dim.
reduction called tSNE