

Title: General Relativity for Cosmology - Lecture 15

Speakers: Achim Kempf

Collection: General Relativity for Cosmology (Kempf)

Date: October 25, 2019 - 4:00 PM

URL: <http://pirsa.org/19100007>

## II Energy and momentum of matter in curved spacetime.

Recall: In flat spacetime, i.e., in Minkowski space, the invariance of the action,  $S$ , under

$$t \rightarrow t_0 + t \text{ and } \vec{x} \rightarrow \vec{x}_0 + \vec{x}$$

implies the conservation of energy and momentum,  $E, P$ , via Noether's theorem.

For fields  $\Psi_{i_1 i_2 \dots i_n}$ : Physical fields imply that at every point in space and time there are flows of energy and flows of momentum, that are conserved if spacetime is flat

## II Energy and momentum of matter in curved spacetime.

Recall: In flat spacetime, i.e., in Minkowski space, the invariance of the action,  $S$ , under

$$t \rightarrow t_0 + t \text{ and } \vec{x} \rightarrow \vec{x}_0 + \vec{x}$$

implies the conservation of energy and momentum,  $E, P$ , via Noether's theorem.

For fields  $\psi_{ijkl...b}$ : Physical fields imply that at every point in space and time there are flows of energy and flows of momentum, that are conserved if spacetime is flat.

### In curved spacetime:

A generic curved spacetime has no translation invariance.

⇒ The flows of energy and momentum will generally not be conserved! How then to identify these flows?

Idea: Whatever plays the role of energy and momentum flows in curved spacetimes must be very sensitive to any changes in the spacetime geometry. Therefore, to study energy and momentum flows in curved spacetime, study:

$$J^{\mu\nu}(x) := 2 \frac{\delta S^{\text{matter}}}{\delta g_{\mu\nu}(x)}$$

↑  
(unrelated to torsion)

convention

In curved spacetime:

A generic curved spacetime has no translation invariance.

⇒ The flows of energy and momentum will generally not be conserved! How then to identify these flows?

Idea: Whatever plays the role of energy and momentum flows in curved spacetimes must be very sensitive to any changes in the spacetime geometry. Therefore, to study energy and momentum flows in curved spacetime, study:

$$J^{\mu\nu}(x) := 2 \frac{\delta S^{(\text{matter})}}{\delta g_{\mu\nu}(x)}$$

↑  
(unrelated to torsion)

convention

In curved spacetime:

A generic curved spacetime has no translation invariance.

⇒ The flows of energy and momentum will generally not be conserved! How then to identify these flows?

Idea: Whatever plays the role of energy and momentum flows in curved spacetimes must be very sensitive to any changes in the spacetime geometry. Therefore, to study energy and momentum flows in curved spacetime, study:

$$J^{\mu\nu}(x) := \underset{\substack{\text{convention} \\ \text{(unrelated to torsion)}}}{2 \frac{\delta S^{(\text{matter})}}{\delta g_{\mu\nu}(x)}}$$

In curved spacetime:

A generic curved spacetime has no translation invariance.

⇒ The flows of energy and momentum will generally not be conserved! How then to identify these flows?

Idea: Whatever plays the role of energy and momentum flows in curved spacetimes must be very sensitive to any changes in the spacetime geometry. Therefore, to study energy and momentum flows in curved spacetime, study:

$$J^{\mu\nu}(x) := \underset{\substack{\text{convention} \\ \text{(unrelated to torsion)}}}{2 \frac{\delta S^{(\text{matter})}}{\delta g_{\mu\nu}(x)}}$$

Def:  $g_{\mu\nu}(\lambda, x)$  is called a deformation of  $g_{\mu\nu}(x)$  for  $x \in B$  if:

a.)  $g_{\mu\nu}(\lambda=0, x) = g_{\mu\nu}(x)$

b.)  $g_{\mu\nu}(\lambda, x) = g_{\mu\nu}(x) \text{ if } x \in M - B$

We then write:  $\delta g_{\mu\nu}(x) := \frac{dg_{\mu\nu}(\lambda, x)}{d\lambda} \Big|_{\lambda=0}$ .

or variation

} Notice: not all these

variations of  $g$  vary the

Riemannian structure!

Def:  $S$  is called functionally differentiable w. resp. to  $g_{\mu\nu}$  in  $B$  if

$$\delta S := \frac{dS}{d\lambda} \Big|_{\lambda=0}$$

exists for all smooth deformations and is of the form:

$$\frac{dS}{d\lambda} \Big|_{\lambda=0} = \frac{1}{2} \int_B \underbrace{\mathcal{T}^{\mu\nu}(x) \delta g_{\mu\nu}(x)}_{\text{any anti-symmetric part drops out.}} d^4x$$

convention

is symmetric:  $g_{\mu\nu} = g_{\nu\mu}$

We then write  $\frac{\delta S}{\delta g_{\mu\nu}(x)} = \frac{1}{2} \mathcal{T}^{\mu\nu}(x)$   $\rightarrow$  By definition, we choose  $\mathcal{T}^{\mu\nu}$  to be symmetric.

Def:  $g_{\mu\nu}(\lambda, x)$  is called a deformation of  $g_{\mu\nu}(x)$  for  $x \in B$  if:

a.)  $g_{\mu\nu}(\lambda=0, x) = g_{\mu\nu}(x)$

b.)  $g_{\mu\nu}(\lambda, x) = g_{\mu\nu}(x) \text{ if } x \in M - B$

We then write:  $\delta g_{\mu\nu}(x) := \frac{dg_{\mu\nu}(\lambda, x)}{d\lambda} \Big|_{\lambda=0}$ .

or variation

} Notice: not all these

variations of  $g$  vary the

Riemannian structure!

Def:  $S$  is called functionally differentiable w. resp. to  $g_{\mu\nu}$  in  $B$  if

$$\delta S := \frac{dS}{d\lambda} \Big|_{\lambda=0}$$

exists for all smooth deformations and is of the form:

$$\frac{dS}{d\lambda} \Big|_{\lambda=0} = \frac{1}{2} \int_B \underbrace{\mathcal{T}^{\mu\nu}(x) \delta g_{\mu\nu}(x)}_{\text{any anti-symmetric part drops out.}} d^4x$$

convention

is symmetric:  $g_{\mu\nu} = g_{\nu\mu}$

We then write  $\frac{\delta S}{\delta g_{\mu\nu}(x)} = \frac{1}{2} \mathcal{T}^{\mu\nu}(x)$   $\rightarrow$  By definition, we choose  $\mathcal{T}^{\mu\nu}$  to be symmetric.

Notice:  $T^{\mu\nu}$  is not a tensor! Why? Rewrite the integral:

$$\underbrace{\frac{dS}{d\lambda}|_{\lambda=0}}_{\text{scalar}} = \frac{1}{2} \int_B T^{\mu\nu}(x) \delta g_{\mu\nu}(x) d^4x$$

$$= \frac{1}{2} \int_B \underbrace{\frac{1}{\sqrt{g}}}_{\text{volume form}} \underbrace{T^{\mu\nu}(x)}_{\text{tensor}} \underbrace{\delta g_{\mu\nu}(x)}_{\text{tensor}} \underbrace{\sqrt{g} dx}_{\text{volume form}}$$

Def: If  $M_{\nu_1 \dots \nu_s}^{v_1 \dots v_r}$  is a tensor, then  $M_{\nu_1 \dots \nu_s}^{v_1 \dots v_r} := M_{\nu_1 \dots \nu_s}^{v_1 \dots v_r} \sqrt{g}$  is called a "Tensor Density".

Thus: □  $T^{\mu\nu}$  is a tensor density and

$$\square T^{\mu\nu} := \frac{1}{\sqrt{g}} T^{\mu\nu} = \frac{2}{\sqrt{g}} \frac{\delta}{\delta g_{\mu\nu}} S' \text{ is a tensor.}$$

Proposition:

$T^{\mu\nu}(x)$  always:


$$T^{\mu\nu}_{;\nu}(x) = 0$$

Proof: later

Strategy:

We need to identify how  $T^{\mu\nu}(x)$  relates to the flows of energy and momentum associated with the matter fields:

→ Consider the cases where spacetime has a symmetry:



$\exists$  diffeomorphism  $\phi: M \rightarrow M$   
so that  $\phi^*g = g$

→ We expect the flows of energy or momentum to be conserved then.  
→ Try to identify these flows by their conservation.

## How to find Killing vector fields?

**Proposition:** For metric connections, the Lie derivative is also:

$$\begin{aligned} L_{\xi} Q^{\alpha \dots b}_{\quad c \dots d} &= Q^{\alpha \dots b}_{\quad \dots d j k} \xi^k \\ &\quad - Q^{\alpha \dots b}_{\quad c \dots d} \xi^a_{j k} - \dots - Q^{\alpha \dots k}_{\quad c \dots d} \xi^b_{j k} \\ &\quad + Q^{\alpha \dots b}_{\quad k \dots d} \xi^k_{j c} + \dots + Q^{\alpha \dots b}_{\quad c \dots k} \xi^b_{j d} \end{aligned}$$

**Proof:** We know it is true with commas, instead of semicolons;

At origin of geodesic cds, can write it with ; too  
because there,  $\Gamma = 0$ . But with ; it is manifestly covariant.

⇒ The above eqn with the ; is true in all coordinate systems.

Apply to  $L_g g$ :

$$L_g g_{\mu\nu} = g_{\nu i k} \xi^k + g_{k\nu} \xi^k_{;i\nu} + g_{\mu k} \xi^k_{;\mu\nu}$$

Using  $\nabla g = 0$  i.e.:  $g_{\nu i k} = 0$  we find:

$$L_g g = 0 \text{ means } g_{k\nu} \xi^k_{;i\nu} + g_{\mu k} \xi^k_{;\mu\nu} = 0$$

$\Rightarrow$  To search for Killing vector fields, i.e., to find out if  $(M, g)$  has symmetries, is to search for vector fields  $\xi$  that obey:

$$\xi_{\mu;i\nu} = -\xi_{\nu;i\mu}$$

Apply to  $L_g g$ :

$$L_g g_{\mu\nu} = g_{\nu i k} \xi^k + g_{k\nu} \xi^k_{;i\mu} + g_{\mu k} \xi^k_{;j\nu}$$

Using  $\nabla g = 0$  i.e.:  $g_{\nu i k} = 0$  we find:

$$L_g g = 0 \text{ means } g_{k\nu} \xi^k_{;i\mu} + g_{\mu k} \xi^k_{;j\nu} = 0$$

$\Rightarrow$  To search for Killing vector fields, i.e., to find out if  $(M, g)$  has symmetries, is to search for vector fields  $\xi$  that obey:

$$\xi_{\mu ; \nu} = - \xi_{\nu ; \mu}$$

Assume the spacetime  $(M, g)$  has a symmetry, described by some Killing vector field  $\xi$ .

Can we then identify flows that are conserved?

Prop.: For every symmetry, i.e., for every Killing vector field  $\xi$  that a spacetime  $(M, g)$  possesses, its matter fields possess a conserved quantity which flows according to the vector field  $P^\mu(x)$ :

$$P^\mu := T^{\mu\nu} \xi_\nu$$

Proof:

$$\begin{aligned} P^\mu_{;\mu} &= (T^{\mu\nu} \xi_\nu)_{;\mu} = \overbrace{T^{\mu\nu}}^0{}_{;\mu} \xi_\nu + T^{\mu\nu} \xi_{\nu;\mu} \\ &= 0 \end{aligned}$$

Symmetric      anti-symmetric  
                        ⇒ = 0

Can we then identify flows that are conserved?

Prop.: For every symmetry, i.e., for every Killing vector field  $\xi$  that a spacetime  $(M, g)$  possesses, its matter fields possess a conserved quantity which flows according to the vector field  $P^\mu(x)$ :

$$P^\mu := T^{\mu\nu} \xi_\nu$$

Proof:

$$\begin{aligned} P^\mu_{;\nu} &= (T^{\mu\nu} \xi_\nu)_{;\nu} = \underbrace{T^{\mu\nu}}_0 + \underbrace{T^{\mu\nu}_{;\nu} \xi_\nu}_{\text{symmetric}} + \underbrace{T^{\mu\nu} \xi_{\nu;\mu}}_{\text{anti-symmetric}} \\ &= 0 \end{aligned}$$

In integral form:

$$0 = \int_B P^r_{;p} \sqrt{g} dx^p = \int_B \text{div}_p P \stackrel{\text{Gauß}}{=} \int_{\partial B} i_p \Omega$$

Note:  $T^r_{;v}=0$  is not a conservation law because  $T^r_{;v}$  is not a divergence!

Thus: As much of  $P^r$  flows into a volume  $B$ ,  
that much also flows out of it.

Def: If the Killing vector field is time-like, i.e., if it generates a time-like translation, then the flow described by the conserved vector field  $K^r(x) = T^r_{;v}(x) \xi_v(x)$  is called the flow of energy. If spacelike, momentum.

→ Def:  $T^{r\nu}(x)$  is called the energy-momentum tensor of fields.  
(with and without Killing fields)

## Energy and momentum of point particles?

Assume the spacetime  $(M, g)$  possesses a Killing field  $\xi$ .

Assume a point-like particle travels on a geodesic  $\gamma$ .

Then:

The quantity  $Q := \xi^\mu j_\mu$  is conserved on the trip  $\gamma$ :

It is called an energy or momentum etc. depending on  $\xi$ .

(In Minkowski space  $\xi^t = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$  and  $\hat{\xi}^t = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$  are Killing fields and  $j_0, j_i$  are energy & momentum)

Proof: Denote the geodesic's tangent vector field by  $u$ . Then:

$$\nabla_u (\xi^\mu u_\mu) = u^\kappa (\xi^\mu u_\mu)_{;\kappa} = \underbrace{u^\kappa \xi^\mu}_{\text{rate of change of } \xi^\mu \text{ along the geodesic } \gamma} \underbrace{j_\kappa}_{\substack{\text{anti-symmetric} \\ \text{symmetric}}} + \underbrace{u^\kappa \xi^\mu}_{=0} \underbrace{u_{;\kappa \mu}}_{\text{because } u^\lambda u_{\lambda ;\kappa} = \nabla_u u = 0 \text{ because geodesic.}} = 0 \quad \checkmark$$

Assume the spacetime  $(M, g)$  possesses a Killing field  $\xi$ .

Assume a point-like particle travels on a geodesic  $\gamma$ .

Then:

The quantity  $Q := \xi^\mu j_\mu$  is conserved on the trip  $\gamma$ :

It is called an energy or momentum etc. depending on  $\xi$ .

(In Minkowski space  $\xi^\mu = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $\hat{\xi}^\mu = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  are Killing fields and  $j_0, j_i$  are energy & momentum)

Proof: Denote the geodesic's tangent vector field by  $u$ . Then:

$$\nabla_u (\xi^\mu u_\mu) = u^\kappa (\xi^\mu u_\mu)_{;\kappa} = \underbrace{u^\kappa \xi^\mu}_{\text{rate of change of } \xi^\mu \text{ along the geodesic } \gamma}{}_{;\kappa} + \underbrace{u^\kappa \xi^\mu u_{\mu;\kappa}}_{\substack{\text{anti-symmetric} \\ \text{symmetric}}} = 0 \quad \checkmark$$

because  
 $u^\kappa u_{\mu;\kappa} = \nabla_\mu u^\kappa = 0$  because geodesic.

Why is  $T^{\mu\nu}_{;\nu} = 0$ ?

Intuition: Many variations

$$g_{\mu\nu}(x) \rightarrow \tilde{g}_{\mu\nu}(x) = g_{\mu\nu}(x) + \delta g_{\mu\nu}(x)$$

do not change the shape of the manifold because  $(M, g)$  and  $(\tilde{M}, \tilde{g})$  describe the same Riemannian structure, i.e., because there is an isometric diffeomorphism, i.e., a coordinate change that relates them.

But  $T^{\mu\nu}(x) = \frac{1}{2} \frac{1}{\sqrt{g}} \frac{\partial S}{\partial g_{\mu\nu}(x)}$

depends on all  $\delta g_{\mu\nu}$ , even the trivial ones!

$\Rightarrow$  Expect  $T^{\mu\nu}$  to contain redundant information.

## Why is $T'^{\nu}_{\mu\nu} = 0$ ?

Intuition: Many variations

$$g_{\mu\nu}(x) \rightarrow \tilde{g}_{\mu\nu}(x) = g_{\mu\nu}(x) + \delta g_{\mu\nu}(x)$$

do not change the shape of the manifold because  $(M, g)$  and  $(\tilde{M}, \tilde{g})$  describe the same Riemannian structure, i.e., because there is an isometric diffeomorphism, i.e., a coordinate change that relates them.

But  $T'^{\nu}_{\mu\nu}(x) = \frac{1}{2} \sqrt{g} \frac{\delta S}{\delta g_{\mu\nu}(x)}$

depends on all  $\delta g_{\mu\nu}$ , even the trivial ones!

$\Rightarrow$  Expect  $T'^{\nu}$  to contain redundant information.

## Proof of $T^{\mu}_{\nu; \nu=0}$ :

□ Assume  $\phi_t: M \rightarrow M$  is a diffeomorphism  $\xleftarrow{t \text{ is the flow parameter, i.e.: } \phi_0 = \text{id}}$   
that is generated by the flow of a vector field,  $\xi$ ,  
that vanishes outside the region  $B \subset M$ , i.e.

$$\phi_t(p) = p \text{ if } p \in M - B$$

(i.e. only the points in  $B$  get re-labeled)

□ Every integral, including the action integral, is invariant under the change of variable, i.e., here under the diffeomorphism  $\phi_s$ ,

Every integral, including the action integral, is invariant under the change of variable, i.e., here under the diffeomorphism  $\phi_t$ , including when the diffeomorphism is infinitesimal. Thus:

start for all matter fields

$$\int_B \mathcal{L}(\psi, \partial\psi, g) d^4x = \int_B \mathcal{L}(\psi, \partial\psi, g) d^4\tilde{x}$$

$\nwarrow$   
Lagrangian density

$$\Rightarrow 0 = \frac{1}{t} \int_B [\mathcal{L} - \phi_t^{-1}(\mathcal{L})] d^4x$$

(total dependence on  $\psi$  and  $\partial\psi$   
vanishes because of eqn  
of motion for the matter  
fields  $\psi$ :

for small  $t$

$$\approx \frac{1}{t} \int_B \left[ \sum_i \frac{\delta \mathcal{L}}{\delta \psi_{(i)}^{a..b}} \text{end} (\psi_{(i)}^{a..b} \text{end} - \phi_t^{-1}(\psi)^{a..b} \text{end}) \right] d^4x = 0$$

$$\text{recognize: } \frac{1}{2} T^{ab} \frac{\delta \mathcal{L}}{\delta g_{ab}} = + \frac{\delta \mathcal{L}}{\delta g_{ab}} (g_{ab} - \phi_t^{-1}(g)_{ab}) \Big] d^4x$$

becomes  $\lim_{t \rightarrow 0} \frac{1}{t} (g - \phi_t^{-1}(g)) = L_P(g)$

Every integral, including the action integral, is invariant under the change of variable, i.e., here under the diffeomorphism  $\phi_t$ , including when the diffeomorphism is infinitesimal. Thus:

$$\int_B \mathcal{L}(\psi, \partial\psi, g) d^4x = \int_B \mathcal{L}(\psi, \partial\psi, g) d^4\tilde{x}$$

start for all matter fields  
↓  
 $B \nearrow$  Lagrangian density

$$\Rightarrow 0 = \frac{1}{t} \int_B [\mathcal{L} - \phi_t^{-1}(\mathcal{L})] d^4x$$

(total dependence on  $\psi$  and  $\partial\psi$   
vanishes because of eqn  
of motion for the matter  
fields  $\psi$ :

$$\approx \frac{1}{t} \int_B \left[ \sum_i \frac{\delta \mathcal{L}}{\delta \psi_{(i)}^{ab}} \text{end} (\psi_{(i)}^{ab} - \phi_t^{-1}(\psi)^{ab}) \right] d^4x$$

for small  $t$

$$+ \frac{\delta \mathcal{L}}{\delta g_{\mu\nu}} (g_{\mu\nu} - \phi_t^{-1}(g)_{\mu\nu}) \right] d^4x$$

recognize:  
 $\frac{1}{2} T^{ab} \sqrt{g} =$

becomes  $\lim_{t \rightarrow 0} \frac{1}{t} (g - \phi_t^{-1}(g)) = L_P(g)$

Every integral, including the action integral, is invariant under the change of variable, i.e., here under the diffeomorphism  $\phi_t$ , including when the diffeomorphism is infinitesimal. Thus:

$$\int_B \mathcal{L}(\psi, \partial\psi, g) d^4x = \int_B \mathcal{L}(\psi, \partial\psi, g) d^4\tilde{x}$$

start for all matter fields  
↓  
 $B \nearrow$  Lagrangian density

$$\Rightarrow 0 = \frac{1}{t} \int_B [\mathcal{L} - \phi_t^{-1}(\mathcal{L})] d^4x$$

(total dependence on  $\psi$  and  $\partial\psi$   
vanishes because of eqn  
of motion for the matter  
fields  $\psi$ :

$$\approx \frac{1}{t} \int_B \left[ \sum_i \frac{\delta \mathcal{L}}{\delta \psi_{(i)}^{ab}} \text{end} (\psi_{(i)}^{ab} - \phi_t^{-1}(\psi)^{ab}) \right] d^4x$$

for small  $t$

$$+ \frac{\delta \mathcal{L}}{\delta g_{\mu\nu}} (g_{\mu\nu} - \phi_t^{-1}(g)_{\mu\nu}) \right] d^4x$$

recognize:  
 $\frac{1}{2} T^{ab} \sqrt{g} =$

becomes  $\lim_{t \rightarrow 0} \frac{1}{t} (g - \phi_t^{-1}(g)) = L_P(g)$

□ Take  $\lim_{\epsilon \rightarrow 0}$   $\Rightarrow$  obtain Lie derivative:

$$O = \int_B \frac{1}{2} T^{ab} \sqrt{g} L_g(g_{ab}) d^4x$$

□ Notice:  $L_g(g_{ab}) = \underbrace{g_{ab;K}\xi^K}_{=0} + g_{kb}\xi^k_{;a} + g_{ak}\xi^k_{;b}$

□ Thus:  $O = \int_B T^{ab} \left( g_{kb}\xi^k_{;a} + g_{ak}\xi^k_{;b} \right) \sqrt{g} d^4x$

$$\begin{aligned} &= \int_B \underbrace{T^{ab} (\xi_{b;a} + \xi_{a;b})}_{\text{symmetric}} \sqrt{g} d^4x \\ &= 2\xi_{b;a} + \underbrace{(\xi_{a;b} - \xi_{b;a})}_{\text{anti-symmetric}} \end{aligned}$$

$$= \int_B 2 T^{ab} \xi_{b;a} \sqrt{g} d^4x$$

$$= 2 \int_B \underbrace{\left( T^a \xi_{b;a} + T^a{}_{jk} \xi_a - T^a{}_{jk} \xi_a \right) \sqrt{g} d^4x}_{=0}$$

$$= 2 \int_B \left[ \underbrace{\left( T^{ab} \xi_b \right)_{;a} \sqrt{g}}_{=0} - T^{ak}{}_{jk} \xi_a \sqrt{g} \right] d^4x$$

Why? define  $r^a := T^{ab} \xi_b$ , then:

$$\int_B r^a{}_{;a} \sqrt{g} d^4x = \int_{\partial B} i_r \Omega = 0$$

because  $\xi = 0$   
on  $\partial B$  by assumption,  
i.e. also  $r^a = T^{ab} \xi_b = 0$   
there.

□ Thus:

$$\int_B T^{ak}{}_{jk} \xi_a \sqrt{g} d^4x = 0 \text{ for all } \xi$$

$$\Rightarrow \boxed{T^{ak}{}_{jk} = 0} \quad \begin{array}{l} \text{consequence of } \checkmark \\ \text{diffeomorphism invariance.} \end{array}$$

But: How to calculate  $T^{\alpha\beta}(x) = \frac{2}{\sqrt{g(x)}} \frac{\delta S^{(\text{matter})}}{\delta g_{\alpha\beta}(x)}$ ?

$$\text{Recall: } S = \int L(\psi, \partial\psi) \sqrt{g} d^4x$$

$$\begin{aligned} \frac{\delta S}{\delta \lambda} \Big|_{\lambda=0} &= \int_B \left( \frac{\partial L}{\partial g_{ab}} \delta g_{ab} + \sum_i \frac{\partial L}{\partial \psi_{(i)\text{end}}^{a-b}} \underbrace{\delta \psi_{(i)\text{end}}^{a-b}}_{=0 \text{ because } \delta \psi = 0} + \int_B L \frac{\partial \Gamma^k}{\partial g_{ab}} \delta g_{ab} \right. \\ &\quad \left. + \sum_i \frac{\partial L}{\partial \psi_{(i)\text{end}}^{a-b}} \underbrace{\delta (\psi_{(i)\text{end}}^{a-b})}_{(*)} \right) \sqrt{g} d^4x \end{aligned}$$

Exercise: prove this.

(\*) Notice:  $\delta(\psi_{(i)\text{end}}^{a-b}) \neq 0$  even though  $\delta \psi = 0$ , because it contains  $\Gamma$  and if  $\delta g \neq 0$  then  $\delta \Gamma \neq 0$ :

$$\delta(\psi_{(i)\text{end}}^{a-b}) = \sum_{k,m,n} \frac{\partial \psi_{(i)\text{end}}^{a-b}}{\partial \Gamma^k_{mn}} \delta \Gamma^k_{mn}$$

But: How to calculate  $T^{\alpha\beta}(x) = \frac{2}{\sqrt{g(x)}} \frac{\delta S^{(\text{matter})}}{\delta g_{\alpha\beta}(x)}$ ?

$$\text{Recall: } S = \int L(\psi, \partial\psi) \sqrt{g} d^4x$$

$$\begin{aligned} \frac{\delta S}{\delta \lambda} \Big|_{\lambda=0} &= \int_B \left( \frac{\partial L}{\partial g_{ab}} \delta g_{ab} + \sum_i \frac{\partial L}{\partial \psi_{(i)\text{c.d}}} \underbrace{\delta \psi_{(i)\text{c.d}}}_{=0 \text{ because } \delta \psi = 0} + \int_B L \frac{\partial \Gamma^a}{\partial g_{ab}} \delta g_{ab} \right. \\ &\quad \left. + \sum_i \frac{\partial L}{\partial \psi_{(i)\text{c.d.e}}} \underbrace{\delta (\psi_{(i)\text{c.d.e}})}_{(*)} \right) \sqrt{g} d^4x \end{aligned}$$

Exercise: prove this.

(\*) Notice:  $\delta(\psi_{(i)\text{c.d.e}}) \neq 0$  even though  $\delta\psi = 0$ , because it contains  $\Gamma$  and if  $\delta g \neq 0$  then  $\delta\Gamma \neq 0$ :

$$\delta(\psi_{(i)\text{c.d.e}}) = \sum_{k,m,n} \frac{\partial \psi_{(i)\text{c.d.e}}}{\partial \Gamma^k_{mn}} \delta \Gamma^k_{mn}$$

But: How to calculate  $T^{\mu\nu}(x) = \frac{2}{\sqrt{g(x)}} \frac{\delta S^{(\text{matter})}}{\delta g_{\mu\nu}(x)}$ ?

$$\text{Recall: } S = \int L(\psi, \partial\psi) \sqrt{g} d^4x$$

$$\begin{aligned} \left. \frac{\partial S}{\partial \lambda} \right|_{\lambda=0} &= \int_B \left( \frac{\partial L}{\partial g_{ab}} \delta g_{ab} + \sum_i \underbrace{\frac{\partial L}{\partial \psi_{(i)}^{a-b} \text{c.d.}}}_{=0 \text{ because } \delta \psi = 0} \delta (\psi_{(i)}^{a-b} \text{c.d.}) + \int_B L \frac{\partial \Gamma^a}{\partial g_{ab}} \delta g_{ab} \right. \\ &\quad \left. + \sum_i \frac{\partial L}{\partial \psi_{(i)}^{a-b} \text{c.d.e}} \delta (\psi_{(i)}^{a-b} \text{c.d.e}) \right) \sqrt{g} d^4x \end{aligned}$$

Exercise: prove this.

(\*) Notice:  $\delta(\psi_{(i)}^{a-b} \text{c.d.e}) \neq 0$  even though  $\delta \psi = 0$ , because it contains  $\Gamma$  and if  $\delta g \neq 0$  then  $\delta \Gamma \neq 0$ :

$$\delta(\psi_{(i)}^{a-b} \text{c.d.e}) = \sum_{k,m,n} \frac{\partial \psi_{(i)}^{a-b} \text{c.d.e}}{\partial \Gamma^k_{mn}} \delta \Gamma^k_{mn}$$

Recall:  $\delta \Gamma^a_{bc}$  is a tensor. It is:

$$\delta \Gamma^a_{bc} = \frac{1}{2} g^{ad} (\delta g_{db;c} + \delta g_{dc;b} - \delta g_{bc;d})$$

(easiest to prove in geodesic i.e. normal coords)

$$\Rightarrow \delta(\Psi_{(i)}^{a..b}{}_{c..d;e}) = \sum_{k=1}^n \frac{\partial \Psi}{\partial r^k} \frac{1}{2} g^{ad} (\delta g_{db;c} + \delta g_{dc;b} - \delta g_{bc;d})$$

Integrate again "by parts"

$$\Rightarrow \delta(\Psi_{(i)}^{a..b}{}_{c..d;e}) \sim \cancel{\delta g_{\mu\nu}}$$

Exercise: work this out

$$\Rightarrow \left. \frac{dS}{d\lambda} \right|_{\lambda=0} = \int_B T^{\mu\nu}(x) \delta g_{\mu\nu}(x) \sqrt{g} d^4x$$

$\Rightarrow$  One can read off  $T^{\mu\nu}(x)$  for any  $L$ .

Example:

4 is scalar, i.e.  $\psi_{,\mu} = \psi_{,\nu}$  e.g.  $V(4) = \frac{m^2}{2\lambda} 4^2 + \frac{\lambda}{4!} 4^4$

$$S' := -\frac{1}{2} \int \left( \psi_{,a} \psi_{,b} g^{ab} + 2V(4) \right) \sqrt{g} d^4x$$

Then:

(Klein Gordon field, e.g. inflaton field)

$$\begin{aligned} \frac{\partial S'}{\partial \lambda} \Big|_{\lambda=0} &= -\frac{1}{2} \int \left( \psi_{,a} \psi_{,b} (\delta g^{ab}) \sqrt{g} + \psi_{,a} \psi_{,b} g^{ab} \frac{\partial \sqrt{g}}{\partial g_{ij}} \delta g_{ij} \right. \\ &\quad \left. + 2V'(4) \frac{\partial \sqrt{g}}{\partial g_{ij}} \delta g_{ij} \right) d^4x \end{aligned}$$

Recall:  $\frac{\partial \sqrt{g}}{\partial g_{ij}} = \frac{1}{2} g^{ij} \sqrt{g}$  i.e.  $\delta g^{ab} = -g^{ai} g^{bj} \delta g_{ij}$

We also notice:  $S(g_{ab} g^{bc}) = 0 = g_{ab} \delta g^{bc} + (\delta g_{ab}) g^{bc}$

Thus:

$$\frac{\partial S'}{\partial \lambda} \Big|_{\lambda=0} = -\frac{1}{2} \int \left( \psi_{,a} \psi_{,b} \sqrt{g} (-g^{ai} g^{bj} \delta g_{ij}) + \psi_{,a} \psi_{,b} g^{ab} \frac{1}{2} g^{ij} \sqrt{g} \delta g_{ij} \right)$$

Example:

$\psi$  is scalar, i.e.  $\psi_{,\mu} = \psi_{,\nu}$  e.g.  $V(\psi) = \frac{m^2}{2\lambda} \psi^2 + \frac{\lambda}{4!} \psi^4$

$$S' := -\frac{1}{2} \int \left( \psi_{,\alpha} \psi_{,\beta} g^{\alpha\beta} + 2V(\psi) \right) \sqrt{g} d^4x$$

Then:

(Klein Gordon field, e.g. inflaton field)

$$\begin{aligned} \frac{\partial S'}{\partial \lambda} \Big|_{\lambda=0} &= -\frac{1}{2} \int \left( \psi_{,\alpha} \psi_{,\beta} (\delta g^{\alpha\beta}) \sqrt{g} + \psi_{,\alpha} \psi_{,\beta} g^{\alpha\beta} \frac{\partial \sqrt{g}}{\partial g_{,\beta}} \delta g_{,\alpha} \right. \\ &\quad \left. + 2V'(\psi) \frac{\partial \sqrt{g}}{\partial g_{,\beta}} \delta g_{,\beta} \right) d^4x \end{aligned}$$

Recall:  $\frac{\partial \sqrt{g}}{\partial g_{,\beta}} = \frac{1}{2} g^{ij} \nabla_j g_{,\beta}$  i.e.  $\delta g^{\alpha\beta} = -g^{\alpha i} g^{\beta j} \delta g_{ij}$

We also notice:  $S(g_{\alpha\beta} g^{\beta\gamma}) = 0 = g_{\alpha\beta} \delta g^{\beta\gamma} + (\delta g_{\alpha\beta}) g^{\beta\gamma}$

Thus:

$$\frac{\partial S'}{\partial \lambda} \Big|_{\lambda=0} = -\frac{1}{2} \int \left( \psi_{,\alpha} \psi_{,\beta} \sqrt{g} (-g^{\alpha i} g^{\beta j} \delta g_{ij}) + \psi_{,\alpha} \psi_{,\beta} g^{\alpha\beta} \frac{1}{2} g^{ij} \nabla_j \sqrt{g} \delta g_{ij} \right)$$

We also notice:  $\partial(g_{\mu\nu}g^{\mu\nu}) = 0 = g_{\mu\nu}\partial g^{\mu\nu} + (\partial g_{\mu\nu})g^{\mu\nu}$

Thus:

$$\frac{\partial S}{\partial \lambda} \Big|_{\lambda=0} = -\frac{1}{2} \int \left( \Psi_{,\alpha} \Psi_{,\beta} \sqrt{g} \left( -g^{ai} g^{bj} \delta g_{ij} \right) + \Psi_{,\alpha} \Psi_{,\beta} g^{ab} \frac{1}{2} g^{ij} \sqrt{g} \delta g_{ij} \right.$$

$$\left. + 2 V(4) \frac{1}{2} g^{ij} \sqrt{g} \delta g_{ij} \right) d^4x$$

$$\Rightarrow \frac{\delta S'}{\delta g_{\mu\nu}} = \frac{1}{2} T^{\mu\nu} \text{ with:}$$

$$T^{\mu\nu} = \left( \underbrace{\Psi_{;\mu} \Psi_{;\nu}}_{= \Psi_{,\alpha} g^{\alpha\mu} g^{\alpha\nu}} - \frac{1}{2} \Psi_{,\alpha} \Psi_{,\beta} g^{\mu\nu} - V(4) g^{\mu\nu} \right) \sqrt{g}$$

i.e. the energy-momentum tensor reads:

$$T_{\mu\nu}^{\text{K.G.}} = \Psi_{;\mu} \Psi_{;\nu} - \frac{1}{2} g_{\mu\nu} \left( \Psi_{,\alpha} \Psi_{,\alpha} + 2 V(4) \right)$$

## Perfect fluid case:

(traditional sense: thermodynamically reversible dynamics)

$$V^M V_\mu = -1$$

□ A perfect (classical) fluid has at every point a unique time-like flux direction vector  $V^\mu$ , the flux is conserved, and the fluid is completely characterized by its local energy density  $\rho$  and pressure  $p$  (i.e., e.g. no shear, no viscosity).

as measured by a co-moving observer:  
 $T^{\mu\nu}, v_\nu = \left(\begin{matrix} \rho & \\ & 0 \end{matrix}\right)$ , so  $T_{\mu\nu} = \left(\begin{matrix} \rho & p \\ 0 & p \end{matrix}\right)$ .

□ Then:  $T_{\mu\nu}^{\text{P.F.}} = (\rho + p) V_\mu V_\nu + p g_{\mu\nu}$  if  $p = 0$ , call it perfect "dust".

□ Note: Eqn. of motion is  $T^{\mu\nu}_{;\nu} = 0$  and dust ( $\rho = 0$ ) travels on geodesics

□ Terminology: (Hawking & Ellis) Any fluid with this  $T_{\mu\nu}$  is called perfect.

### Definition:

The "equation of state" of a perfect fluid is the relation between its energy density,  $\rho$  and its pressure,  $p$ . It depends on the fluid and so one can characterize the fluids by this parameter:

$$w := \frac{p}{\rho}$$

### Important later for cosmology:

The two tensors

$$T_{\mu\nu}^{KL} = g_{\mu\nu} g_{ij} - \frac{1}{2} g_{\mu\nu} (g_{ia} g^{ia} + 2 V(g))$$

$$\text{and } T_{\mu\nu}^{P,T} = (\rho + p) v_\mu v_\nu + g_{\mu\nu} p$$

are of similar form (unlike e.g.  $T_{\mu\nu}^{EM}$ )

applies to the inflaton field.

### Definition:

The "equation of state" of a perfect fluid is the relation between its energy density,  $\rho$  and its pressure,  $p$ . It depends on the fluid and so one can characterize the fluids by this parameter:

$$w := \frac{p}{\rho}$$

### Important later for cosmology:

The two tensors

$$T_{\mu\nu}^{KL} = g_{\mu\nu} g_{ij} - \frac{1}{2} g_{\mu\nu} (g_{ia} g^{ia} + 2 V(g))$$

$$\text{and } T_{\mu\nu}^{P,T} = (\rho + p) v_\mu v_\nu + g_{\mu\nu} p$$

are of similar form (unlike e.g.  $T_{\mu\nu}^{EM}$ )

applies to the inflaton field.

if  $\Psi$  is almost homogeneous, i.e.  $\Psi_{j,i} \xrightarrow{i=1,2,3} 0$ :

Then, define:  $v_\mu := \frac{\Psi_{i\mu}}{\sqrt{|g^{ab}\Psi_{ja}\Psi_{ib}|}}$  (so that  $v_\mu v^\mu = -1$ )

$$\text{i.e.: } T_{\mu\nu}^{\text{K.G.}} = |g^{ab}\Psi_{ja}\Psi_{ib}| v_\mu v_\nu + g_{\mu\nu} \left( -\frac{1}{2} \Psi_{ja} \Psi^{ja} - V(\Psi) \right)$$

$$\approx +\dot{\Psi}^2 v_\mu v_\nu + g_{\mu\nu} \left( \frac{1}{2} \dot{\Psi}^2 - V(\Psi) \right)$$

□ Compare with  $T^{\text{PF}}$ :

$$\frac{\mu+P}{P} = \frac{1}{w} + 1 = \frac{\dot{\Psi}^2}{\frac{1}{2} \dot{\Psi}^2 - V(\Psi)}$$

$$\Rightarrow \quad \frac{1}{w} = \frac{\dot{\Psi}^2}{\frac{1}{2} \dot{\Psi}^2 - V(\Psi)} - \frac{\dot{\Psi}^2/2 - V(\Psi)}{\frac{1}{2} \dot{\Psi}^2 - V(\Psi)} = \frac{\dot{\Psi}^2/2 + V(\Psi)}{\dot{\Psi}^2/2 - V(\Psi)}$$

$$\text{□ Thus: } w = \frac{\dot{\Psi}^2/2 - V(\Psi)}{\dot{\Psi}^2/2 + V(\Psi)} \in (-1, 1)$$

potential dominated, i.e.  
 $V(\Psi) \gg \dot{\Psi}^2$  (see inflation later)

no potential:  $V(\Psi) = 0$

if  $\Psi$  is almost homogeneous, i.e.  $\Psi_{j;i} \approx 0$ :

Then, define:  $v_\mu := \frac{\Psi_{i;\mu}}{\sqrt{|g^{ab}\Psi_{j;a}\Psi_{j;b}|}}$  (so that  $v_\mu v^\mu = -1$ )

$$\begin{aligned} \text{i.e.: } T_{\mu\nu}^{\text{vac.}} &= |g^{ab}\Psi_{j;a}\Psi_{j;b}| v_\mu v_\nu + g_{\mu\nu} \left( -\frac{1}{2} \Psi_{j;a}\Psi^{ja} - V(\Psi) \right) \\ &\approx +\dot{\Psi}^2 v_\mu v_\nu + g_{\mu\nu} \left( \frac{1}{2} \dot{\Psi}^2 - V(\Psi) \right) \end{aligned}$$

□ Compare with  $T^{\text{PF}}$ :

$$\frac{\mu+P}{P} = \frac{1}{w} + 1 = \frac{\dot{\Psi}^2}{\frac{1}{2} \dot{\Psi}^2 - V(\Psi)}$$

$$\Rightarrow \quad \frac{1}{w} = \frac{\dot{\Psi}^2}{\frac{1}{2} \dot{\Psi}^2 - V(\Psi)} - \frac{\dot{\Psi}^2/2 - V(\Psi)}{\frac{1}{2} \dot{\Psi}^2 - V(\Psi)} = \frac{\dot{\Psi}^2/2 + V(\Psi)}{\dot{\Psi}^2/2 - V(\Psi)}$$

□ Thus:  $w = \frac{\dot{\Psi}^2/2 - V(\Psi)}{\dot{\Psi}^2/2 + V(\Psi)} \in (-1, 1)$

potential dominated, i.e.  
 $V(\Psi) \gg \dot{\Psi}^2$  (see inflation later)  
no potential:  $V(\Psi) = 0$