Title: General Relativity for Cosmology - Lecture 18

Speakers: Achim Kempf

Collection: General Relativity for Cosmology (Kempf)

Date: November 12, 2019 - 4:00 PM

URL: http://pirsa.org/19110003

2019-gr18.pdf - HP Sure Click Secure View

Horizons & Singularities

Local causal structure

The metric, g, not only defines the "shape" of a pseudo-Riemannian manifold, it also defines what is causal and what is acausal: (by defining what is space-, null- or time-like)

Preparation: Consider an arbitrary point $p \in M$ and an arbitrary "convex normal neighborhood" of p, i.e., a set $U \subset M$ with $p \in U$ for which holds:

$q, r \in U \Rightarrow$ there exist a unique geodesic connecting q and r.

Type here to search
ENG 4:06 PM 2019-11-12

Now consider in $\mathcal{U}$:



past light cone is determined by the metric.

some arbitrary coordinate system in which the hyperplanes $t = \text{const.}$ are space-like.

Definition: In order for the laws of matter fields Ψ to be called "locally causal" (and therefore reasonable), their equations of motion must allow one to calculate $\Psi(p)$ from only the values $\Psi(q)$ and finite order derivatives $\Psi(q), \ldots$ for all $q \in F$.

Remark:
For massless fields of spin >1, there is no natural linear equation of motion with such well-defined causality.
Note: Gravitons are spin $s=2$ but their dynamics is ultimately nonlinear.
(See Wald p. 375)

2019-gr18.pdf - HP Sure Click Secure View

Remark: In Newton's theory these data don't suffice, because there: $c = \infty$

Equivalently: The laws of matter fields are locally causal if signals can be sent between events $q, p \in \mathcal{U}$ only iff there is a curve $\gamma \subset \mathcal{U}$ with $\gamma(t_1) = q$, $\gamma(t_2) = p$ whose tangents are non-space-like:

$$g(\dot\gamma(t), \dot\gamma(t)) \leq 0 \quad \text{for all } t \in [t_a, t_b]$$

Question: Assume that on a differentiable manifold $\mathcal{M}$ only a causal structure is given. To what extent fixes this g?

Answer: Nearly completely!

We will obtain yet another way to describe the "shape" of a curved spacetime!

Type here to search
ENG 4:12 PM 2019-11-12

2019-gr18.pdf - HP Sure Click Secure View

Theorem:

Assume that on a differentiable manifold M we don't know the metric, i.e., we can't evaluate

$$g(\xi, \eta)$$

but assume that for all $p \in M$ and all $\xi \in T_p(M)$ we know for each ξ whether it is space-, light- or time-like, i.e. assume we know: $\in \{-1, 0, 1\}$

$$\operatorname{sign}(g(\xi, \xi)) \text{ for all } p \in M,\ \xi \in T_p(M)$$

Then, this information already determines the metric tensor up to conformal transformations, i.e., we obtain:

$$c(x)\, g_{\mu\nu}(x)$$

$c(x)$: unspecified scalar function: "conformal factor"

$g_{\mu\nu}(x)$: metric in canonical frame (also called "holonomic frame")

Type here to search

ENG 4:16 PM 2019-11-12

2019-gr18.pdf - HP Sure Click Secure View

Remark:

Conformal transformations affect only the length of vectors but leave their mutual "angles" invariant:

$$\cos(\sphericalangle(\xi,\eta)) = \frac{g(\xi,\eta)}{\sqrt{g(\xi,\xi)\,g(\eta,\eta)}} \qquad \left(\frac{c}{\sqrt{c}\sqrt{c}}\right)$$

Proof: ▯ Consider a timelike ξ and a spacelike η.

Are there linear combinations

$$\xi + \lambda\eta$$

that are light-like? If yes, we can assume that we know these λ from knowing the causal structure!

Type here to search

ENG 4:18 PM 2019-11-12


$\cos(\measuredangle(a,b)) = \frac{a \cdot b}{|a|\,|b|}$

2019-gr18.pdf - HP Sure Click Secure View

Proof: Consider a timelike ξ and a spacelike η.

Are there linear combinations

$$\xi + \lambda \eta$$

that are light-like? If yes, we can assume that we know these λ from knowing the causal structure!

Need to solve this quadratic equation in λ:

$$f(\lambda) = g(\xi + \lambda\eta, \xi + \lambda\eta) = 0 \qquad (*)$$

i.e.: $g^{\mu\nu}(\xi_\mu + \lambda\eta_\mu)(\xi_\nu + \lambda\eta_\nu) = 0$

Type here to search
ENG 4:22 PM 2019-11-12

- Need to solve this quadratic equation in λ:

$$f(\lambda) = g(\xi + \lambda\eta, \xi + \lambda\eta) = 0 \qquad (*)$$

i.e.: $g^{\mu\nu}(\xi_\mu + \lambda\eta_\mu)(\xi_\nu + \lambda\eta_\nu) = 0$

- Eq. (*) has two roots λ_1, λ_2. Are they real?

Yes, because:

ξ timelike $\Rightarrow$ $f(0) < 0$

η spacelike $\Rightarrow$ $f(\lambda) > 0$ for large enough λ

$\Rightarrow$ $f(\lambda) = 0$ has one real root

$\Rightarrow$ Both roots, λ_1, λ_2 of $f(\lambda) = 0$ are real.

- Since by assumption we can identify all null vectors we can assume λ_1, λ_2 known.

2019-gr18.pdf - HP Sure Click Secure View

Lemma:

$$\frac{g(\xi,\xi)}{g(\eta,\eta)} = \lambda_1 \lambda_2$$

Thus, the ratio $\frac{g(\xi,\xi)}{g(\eta,\eta)}$ can be assumed known for all timelike ξ and all spacelike η.

Proof: From $g(\xi+\lambda_{1,2}\eta, \xi+\lambda_{1,2}\eta)=0$

we have: $g(\xi,\xi) + 2\lambda_1 g(\xi,\eta) + \lambda_1^2 g(\eta,\eta) = 0$

and: $g(\xi,\xi) + 2\lambda_2 g(\xi,\eta) + \lambda_2^2 g(\eta,\eta) = 0$

Eliminate $g(\xi,\eta)$ $\Rightarrow$ $\frac{g(\xi,\xi)}{g(\eta,\eta)} = \lambda_1 \lambda_2$ ✓

Exercise: show this

Type here to search
ENG 4:29 PM 2019-11-12

2019-gr18.pdf - HP Sure Click Secure View

Corollary:

Also the ratios $\frac{g(\xi,\xi)}{g(\xi',\xi')}$ for ξ, ξ' both timelike (or both spacelike) can be assumed known:

$$\frac{g(\xi,\xi)}{g(\eta,\eta)} = \lambda_1\lambda_2 \;;\; \frac{g(\xi',\xi')}{g(\eta,\eta)} = \lambda_1'\lambda_2' \Rightarrow \frac{g(\xi',\xi')}{g(\xi,\xi)} = \frac{\lambda_1'\lambda_2'}{\lambda_1\lambda_2}$$

Corollary:

Consider arbitrary non-null vectors α, β.

Then

$$g(\alpha,\beta) = \frac{-1}{2}\left[g(\alpha,\alpha) + g(\beta,\beta) - g(\alpha+\beta, \alpha+\beta)\right]$$

and thus: (By lemma, all these ratios can be assumed known)

$$\frac{g(\alpha,\beta)}{g(\xi,\xi)} = \frac{-1}{2}\left[\frac{g(\alpha,\alpha)}{g(\xi,\xi)} + \frac{g(\beta,\beta)}{g(\xi,\xi)} - \frac{g(\alpha+\beta,\alpha+\beta)}{g(\xi,\xi)}\right]$$

We can consider $g(\xi,\xi)$ to be a fixed, unknown scalar function.

Type here to search
ENG 4:31 PM 2019-11-12

2019-gr18.pdf - HP Sure Click Secure View

Implications: Spacetimes (M, g) and $(M, \tilde{g})$ for which

$$\tilde{g} = \phi\, g$$

(if $\phi = 0$ then not invertible; if $\phi < 0$ then change signature)

↑ some positive scalar function

possess the same causal structure.

⇒ Spacetimes fall into "conformal equivalence classes" within which the local causal structure is invariant.

⇝ This is very useful to help intuition:

Choose a conformally equivalent spacetime, for which space and time are conformally so much squeezed that infinities turn into a finite distance, all while 45° remain 45° degrees b/c conformality.

Type here to search
ENG 4:40 PM 2019-11-12

$\tilde{g} = \phi g$

(if $\phi = 0$ then g not invertible
if $\phi < 0$ then change signature)

↑ some positive scala function

possess the same causal structure.

⇒ Space times fall into "conformal equivalence classes" within which the local causal structure is invariant.

⇝ This is very useful to help intuition:

Choose a conformally equivalent spacetime, for which space and time are conformally so much squeezed that infinities turn into a finite distance, all while 45° remain 45° degrees b/c conformality.

2019-gr18.pdf - HP Sure Click Secure View

Application: Penrose diagrams

Example: Consider Minkowski space, (M, g) in spherical coordinates:

$$g = -dx^0 \otimes dx^0 + dx^1 \otimes dx^1 + dx^2 \otimes dx^2 + dx^3 \otimes dx^3$$
$$= -dt \otimes dt + dr \otimes dr + r^2 (d\theta \otimes d\theta + \sin^2\theta \, d\phi \otimes d\phi)$$

with $-\infty < t < \infty$, $0 \le r < \infty$, $0 \le \phi < 2\pi$, $0 \le \theta < \pi$

Now consider the spacetime $(\bar{M}, \bar{g})$ given by:

$$\bar{g} = d\bar{t} \otimes d\bar{t} + d\bar{r} \otimes d\bar{r} + \sin^2(\bar{r}^2)(d\theta \otimes d\theta + \sin^2\theta \, d\phi \otimes d\phi)$$

with $\underbrace{-\pi < \bar{t} + \bar{r} < \pi, -\pi < \bar{t} - \bar{r} < \pi}_{\text{finite!}}, \bar{r} \ge 0, 0 \le \phi < 2\pi, 0 \le \theta < \pi$

The spacetimes $(M, g), (\bar{M}, \bar{g})$ are related by a diffeomorphism $\bar{M} \to M$:

$$t := \frac{1}{2}\tan\left(\frac{1}{2}(\bar{t} + \bar{r})\right) + \frac{1}{2}\tan\left(\frac{1}{2}(\bar{t} - \bar{r})\right)$$
$$r := \frac{1}{2}\tan\left(\frac{1}{2}(\bar{t} + \bar{r})\right) - \frac{1}{2}\tan\left(\frac{1}{2}(\bar{t} - \bar{r})\right)$$

Type here to search
ENG 4:48 PM 2019-11-12

The diffeomorphism is not isometric, but it is conformal:

$$g_{\mu\nu} = \phi\, \bar{g}_{\mu\nu} \quad \text{with} \quad \phi = \frac{1}{4}\sec^2\left(\tfrac{1}{2}(\bar{t}+\bar{r})\right)\sec^2\left(\tfrac{1}{2}(\bar{t}-\bar{r})\right)$$

Thus, $(\mathcal{M}, g)$ and $(\bar{\mathcal{M}}, \bar{g})$ have the same causal structure, although $-\pi < \bar{t}+\bar{r} < \pi$ and $-\pi < \bar{t}-\bar{r} < \pi$ and $\bar{r} > 0$.

⇝ Use this to study the causal structure using $(\bar{\mathcal{M}}, \bar{g})$ which is of finite size:

Legend:

- Continuous (green) lines: Infinities
- Dotted (green) line: Radius = 0
- Singularities (hats): double line.



2019-gr18.pdf - HP Sure Click Secure View

Examples:

A) geodesic, massive observers, sitting at r_i.

B) same but then uniformly accelerating.

C) light ray



Type here to search
ENG
5:00 PM
2019-11-12

Definition:

An observer's Event horizon (if any) is the boundary of the past of this observer's future causal infinity.

I.e., the event horizon is the boundary of the set of those events that can possibly ever influence the observer, i.e., it's the boundary of the set of events the observer can ever learn about.



Recall FL metric for $K=0$ (i.e. spatially flat $\mathbb{R}^3$)

$$g^{(FL)} = -dt\otimes dt + a^2(t)\,dr\otimes dr + a^2(t)r^2\left(d\theta\otimes d\theta + \sin^2\theta\, d\phi\otimes d\phi\right)$$

Change to a new time variable τ: "conformal time"

$$\tau(t) := \int_{t_0}^{t} \frac{1}{a(t')}\,dt'$$

Why useful? Notice: $\frac{d\tau}{dt} = \frac{1}{a(t)} \Rightarrow dt = a\,d\tau$

$$\Rightarrow g^{(FL)} = a^2(\tau)\left(-d\tau\otimes d\tau + dr\otimes dr + r^2\left(d\theta\otimes d\theta + \sin^2\theta\, d\phi\otimes d\phi\right)\right)$$

$$\Rightarrow g^{(FL)} = a^2(\tau)\,\eta$$

$\Rightarrow g^{(FL)}$ is conformally equivalent to Minkowski space!

$\Rightarrow$ Can re-use some Penrose diagram, except range of time τ may be smaller!

Why useful: Notice: $\frac{d}{dt} = \frac{1}{a(t)} \frac{d}{d\tau} \Rightarrow dt = a\, d\tau$

$$\Rightarrow g^{(FL)} = a^2(\tau)\left(-d\tau \otimes d\tau + dr \otimes dr + r^2(d\theta \otimes d\theta + \sin^2\theta\, d\phi \otimes d\phi)\right)$$

$$\Rightarrow g^{(FL)} = a^2(\tau)\,\eta$$

$\Rightarrow g^{(FL)}$ is conformally equivalent to Minkowski space !

$\Rightarrow$ Can re-use some Penrose diagram, except range of time τ may be smaller !

Penrose diagrams of F.L. cosmologies: (with $K=0$)

Example: Radiation dominated universe: $a(t) = \sqrt{t}$, $t > 0$

$$\Rightarrow \tau(t) := \int_0^t \frac{1}{\sqrt{t}}\, dt = 2\sqrt{t}\Big|_0^t = 2\sqrt{t} \quad \Rightarrow \tau > 0$$

Penrose diagrams of F.L. cosmologies: (with $K=0$)

Example: Radiation dominated universe: $a(t) = \sqrt{t}$, $t > 0$

$$\Rightarrow \tau(t) := \int_0^t \frac{1}{\sqrt{t}}\, dt = 2\sqrt{t}\Big|_0^t = 2\sqrt{t} \quad \boxed{\Rightarrow \tau > 0}$$

$\Rightarrow$ Obtain (τ, r) diagram with $\tau > 0$:

A.) geodesic, massive observer

B) same but then uniformly accelerating

C) light rays



Notice: Singularity at $t=0$ assumed. (Some FL spacetimes are without, e.g. de Sitter: $a(t) = e^{Ht}$)

At finite t, an observer can see only a finite distance.

Def: This distance is called the observer's "Particle Horizon" at time t.

2019-gr18.pdf - HP Sure Click Secure View

Particle horizon:

How far away, d_p, is the particle horizon at time t?



Recall: (r is the comoving radius, i.e., galaxies sit at fixed (r,θ,ϕ) at all times. Recall that by definition, $a(t_{today})=1$, i.e., comov. distance = proper distance today.)

$$g = -dt\otimes dt + a^2(t)\,dr\otimes dr + r^2(d\theta\otimes d\theta + \sin^2\theta\, d\phi\otimes d\phi)$$

$t(\lambda)$ $r(\lambda)$ $\theta(\lambda)$ $\phi(\lambda)$

Consider a light ray $\gamma(\lambda) = (\gamma^0(\lambda), \gamma^1(\lambda), 0, 0)$, i.e., emitted radially.

Its tangent is null $g_{\mu\nu}\frac{\partial\gamma^\mu}{\partial\lambda}\frac{\partial\gamma^\nu}{\partial\lambda}=0$, i.e.:

$$\left(\frac{\partial\gamma^0(\lambda)}{\partial\lambda}\right)^2 - a^2(t)\left(\frac{\partial\gamma^1(\lambda)}{\partial\lambda}\right)^2 = 0$$

Note: $\gamma^0(\lambda) = t(\lambda)$

Thus: $\frac{dt}{d\lambda} = \pm a(t)\frac{dr}{d\lambda}$ i.e. $\frac{dr}{dt} = \pm\frac{1}{a(t)}$

Note: this speed is not $=1=c$ because r is the comoving distance. At late times, $a(t)\gg 1$, i.e., $\frac{dr}{dt}$ small, i.e., light crosses comoving distances slowly, because the same comoving distance becomes larger and larger.

Type here to search
ENG 5:20 PM 2019-11-12

2019-gr18.pdf - HP Sure Click Secure View

Thus: $d_p = \int_{t=0}^{t} \frac{1}{a(t')}\, dt'$ (It's the comoving distance travelled, and with $a(\text{today}) = 1$, it's also the current proper distance to what's the furthest we can see.)

For example for us today: $d_p \approx 4 \cdot 10^{10}$ light years. (Say since CMB emission)

Recall event horizon:

An observer's event horizon is the boundary of the past of this observer's future infinity.

⇒ If we have a cosmological event horizon, it is the particle horizon that we will have at future infinity.

Q: Do we have a cosmological event horizon?

A: Depends on behavior of $a(t)$ for $t \to \infty$:

I.e., does $d_p^{\infty} = \int_0^{\infty} \frac{1}{a(t)}\, dt$ converge to a finite comoving distance?

Type here to search
ENG 5:27 PM 2019-11-12

2019-gr18.pdf - HP Sure Click Secure View

Thus: $d_p = \int_{t=0}^{t} \frac{1}{a(t')}\, dt'$ (It's the comoving distance travelled, and with $a(\text{today}) = 1$, it's also the current proper distance to what's the furthest we can see.)

For example for us today: $d_p \approx 4 \cdot 10^{10}$ light years. (Say since CMB emission)

Recall event horizon:

An observer's event horizon is the boundary of the past of this observer's future infinity.

⇒ If we have a cosmological event horizon, it is the particle horizon that we will have at future infinity.

Q: Do we have a cosmological event horizon?

A: Depends on behavior of $a(t)$ for $t \to \infty$:

I.e., does $d_p^{\infty} = \int_{t_0}^{\infty} \frac{1}{a(t)}\, dt$ converge to a finite comoving distance?

Type here to search
ENG 5:28 PM 2019-11-12

2019-gr18.pdf - HP Sure Click Secure View

Recall: $a(t) \sim t^{\frac{2}{3(1+w)}}$

Notice: convergence iff $\tau < -1$

$$\Rightarrow d_p^{\infty} = \int_0^{\infty} \frac{1}{a(t)}\, dt \sim \int_0^{\infty} t^{\frac{-2}{3(1+w)}}\, dt \qquad (t^{\tau})$$

$\Rightarrow \exists$ Event horizon iff $w < -\frac{1}{3}$, i.e., if "inflation", i.e., iff $\ddot{a} > 0$!

Notice:

$$d_p(t) = \int_0^{t} \frac{1}{a(t)}\, dt = \tau(t) = \underline{\text{Conformal time !}}$$

$\Rightarrow$ $d_p^{\infty} = d_p(t=\infty)$ is finite $\Leftrightarrow$ $\tau(t=\infty)$ is finite

$\Rightarrow$ Iff inflation then Penrose diagram truncated above at a $\tau = \tau_{max}$ line:

Type here to search

ENG 5:34 PM 2019-11-12

2019-gr18.pdf - HP Sure Click Secure View

Recall:

FLRW spacetime with $K = 0$, big bang and no late inflation:



Type here to search
ENG 5:39 PM 2019-11-12

Now with inflation:

(as we have today and presumably in the future)

FL spacetime with $K=0$, big bang and late inflation:



2019-gr18.pdf - HP Sure Click Secure View

Black holes:

The metric of an eternal, nonrotating, uncharged classical black hole was first found by Schwarzschild in Dec. 1915. It can be written as:

$$ds^2 = \left(1 - \frac{r_s}{r}\right) dt^2 - \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 - r^2\left(d\theta^2 + \sin^2\theta\, d\varphi^2\right)$$

where $r_s = 2GM$ is the Schwarzschild radius.

Notice: At $r = r_s$ this representation of the metric, g, becomes singular. E.g. Kruskal coordinates show that g is regular there.

$\rightsquigarrow$ $r = r_s$ is merely the event horizon (which is light-like!)

Only $r = 0$ is a singularity (it is spacelike).

Type here to search

ENG 5:41 PM 2019-11-12

2019-gr18.pdf - HP Sure Click Secure View

Q: Can we analyse the causal structure using a Penrose diagram, i.e., a conformally equivalent diagram whose light rays are at $\pm 45°$?

Q: I.e., is the metric conformally equivalent to Minkowski space?

Q: Also, can we include the full dynamics of the black hole?

A: Yes, if we consider only the r,t plane. Why?

Theorem: The metric $g_{\mu\nu}$ of any 2-dimensional Lorentzian manifold or sub-manifold reads in suitable coordinates:

$$g_{\mu\nu}(x) = \Omega(x)\,\eta_{\mu\nu}$$

(Ω: scalar function)

Why? In 2D, $g_{\mu\nu}(x)$ has 3 independent entries and 2 of them can be fixed by choosing the 2 coordinate change functions $\bar{x}_1(x), \bar{x}_2(x)$.

$\Rightarrow$ Every 2D Lorentzian submanifolds of any 3+1 metric has a Penrose diagram.

Type here to search
ENG 5:43 PM 2019-11-12

A: Yes, if we consider only the r,t plane. Why?

Theorem: The metric $g_{\mu\nu}$ of any 2-dimensional Lorentzian manifold or sub-manifold reads in suitable coordinates:

$$g_{\mu\nu}(x) = \alpha(x)\,\eta_{\mu\nu}$$

↑ scalar function

Why? In 2D, $g_{\mu\nu}(x)$ has 3 independent entries and 2 of them can be fixed by choosing the 2 coordinate change functions $\bar{x}_1(x), \bar{x}_2(x)$.

⇒ Every 2D Lorentzian submanifolds of any 3+1 metric has a Penrose diagram.

Example: Collapsing star, forming black hole (non-rotating)

2019-gr18.pdf - HP Sure Click Secure View

Example: Collapsing star, forming black hole (non-rotating)



For the transformation, see, e.g., the text by Susskind and Lindsay.

Type here to search
ENG 5:46 PM 2019-11-12