Title: The Cohomology of Groups (Johnson-Freyd/Guo) - Lecture 9

Speakers: Meng Guo, Theo Johnson-Freyd

Collection: The Cohomology of Groups (Johnson-Freyd/Guo)

Date: November 27, 2019 - 10:00 AM

URL: http://pirsa.org/19110045

1707.08388 + 1810.00463
Ej + w/ D. Treumann.

Goal: Thm. $H^3(\mathbb{M}, U(1)) \cong \mathbb{Z}_{24} \oplus (\leq 4)$
(probably trivial)

$\mathbb{M}$ = Fischer-Griess monster is the largest sporadic simple gp.

$$\#\mathbb{M} = 2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 13^3 \cdot 11^2 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71 \sim 10^{54}$$

$\#\mathbb{M} = 2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 13^3 \cdot 11^2 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71 \sim 10^{54}$

remark: $\in \#SL_{15}(\mathbb{F}_2)$.

$H^1(\mathbb{M}, U(1)) = \hom(\mathbb{M}, U(1)) = \{*\}$ because $\mathbb{M}$ is simple

$H^2(\mathbb{M}, U(1)) = \{\text{central extensions}\} = \{*\}$ ~1980s.

$H^3(\mathbb{M}, U(1))$ $\exists$ 't Hooft anomalies of actions of $\mathbb{M}$ on (1+1)d QFTs

Method. work prime by prime. $H^\bullet(\mathbb{M}, U(1))$ is a finite ab. gp.

$\bigoplus_{p \mid \#\mathbb{M}} H^\bullet(\mathbb{M}; U(1))_{(p)}$ "p-part"; order is a power of p

s.t. $\chi(\#\mathbb{M}) = 0$ on this finite gp.

$\mathbb{Z}_p \oplus \mathbb{Z}_p$ or $\mathbb{Z}_{p^2}$ or ...

Sylow theory: $\forall p$. $\exists S \subseteq M$ s.t. $\frac{\#M}{\#S}$ is coprime to p.

$H^\cdot(M; \mathcal{U}(1))_{(p)} \underset{\text{"transfer"}}{\overset{\text{rest}}{\rightleftarrows}} H^\cdot(S, \mathcal{U}(1))_{(p)}$

a sort of integral / push forward

$\circlearrowleft$ = mult by $\frac{\#M}{\#S}$.

is an IO on $H^\cdot_{(p)}$.

rest is an injection onto a direct summand.
on (p)-parts.

rest is an injection onto a direct summand on (p)-parts.

Eg: $p \geq 17$ p-Sylow in M is a $\mathbb{Z}_p$. in fact $M \geq S = \mathbb{Z}_p \rtimes \mathbb{Z}_{\frac{p-1}{2}}$.

$$\begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \in PSL_2(\mathbb{F}_p).$$

$$H^3(M; \mathcal{U}(1))_{(p)} \subseteq H^3(S; \mathcal{U}(1))_{(p)}$$

$\uparrow$ compat w/ S.S.

$$\#M = 2^{46} \cdot 3^{20} \cdot 5^{9} \cdot 7^{6} \cdot 13^{3} \cdot \underline{11^{2} \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71} \sim 10^{54}$$

remark: $\in \#SL_{13}(\mathbb{F}_2)$.

$$E_2^{i,j} = H^i\left(\overbrace{\mathbb{Z}_{p^{\frac{1}{2}}}}^{\text{acts on } (p)\text{-torsion}}; \overbrace{H^j(\mathbb{Z}_p; U(1))}^{\text{purely } (p)\text{-torsion}}\right)_{(p)} \Rightarrow H^{i+j}(S; U(1))_{(p)}$$

if $i > 0$, then H^i = "purely 2-torsion" $= 0$.

$E_2^{i,j} = \{i = 0 \text{ columns}\}$.

$\mathbb{Z}_p$ if j odd

0 if j even.

$$H^{j+1}(\mathbb{Z}_p, \mathbb{Z}) = \mathbb{Z}[t]/(p^t = 0)$$

$\mathbb{Z}_{p^{\frac{1}{2}}}$ acts faithfully on $\mathbb{Z}_p \cdot t$

acts by k^{th} power of the faithful action on $\mathbb{Z}_p \cdot t^i$

$H^3(M; \mathcal{U}(1))_{(p)} \subseteq H^3(S; \mathcal{U}(1))_{(p)}$

↑ compat w/ S.S.

$\begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \in PSL_2(\mathbb{F}_p)$

$\mathbb{Z}_p$

$\deg \leq p$ $PSL_2(\mathbb{F}_p)$

$\mathbb{F}_p^\times \simeq \mathbb{Z}_{p-1}$

E_2 $\quad \mathbb{Z}_p$ $\quad \mathbb{Z}_p$ $\quad \mathbb{Z}_p$

$\mathcal{U}(1)$ $\quad 0$ $\quad 0$ $\quad 0$

$j=0$

$= E_\infty = H^*(S)$.

So $H^3(S; \mathcal{U}(1))_{(p)} = 0$.

So $H^3(M; \mathcal{U}(1))_{(p)} = 0$.

$p=11$ use $S = (\mathbb{Z}_{11} \rtimes \mathbb{Z}_5)^3 \subseteq M$ ...

$3 \subseteq M$

Push Quad

$$\#\mathbb{M} = 2^{46}\cdot 3^{20}\cdot 5^{9}\cdot 7^{6}\cdot 13^{3}\cdot 11^{2}\cdot 17\cdot 19\cdot 23\cdot 29\cdot 31\cdot 41\cdot 47\cdot 59\cdot 71 \sim 10^{54}$$

✓

remark: $\in \#SL_{13}(\mathbb{F}_2)$.

The primes $p = 2, 3, 5, 7, 13$ are special. (cf. Conway–Norton)
Monstrous moonshine conjectures

- there are non-conjugate cyclic subgps of $\mathbb{M}$ of order p
- p-Sylow in $\mathbb{M}$ is nonabelian.
- $p-1$ divides 24.

Why? I don't know. But I know the main ingredient.


Remind: ∃ special lattice "Leech" Λ
full rank discrete subgp of $\mathbb{R}^{\text{rank}=24}$
• rank 24
so ∃ an isom. of gps $\Lambda \simeq \mathbb{Z}^{24}$.
e.g. E_8 →
• even: if $\lambda \in \Lambda$, $\|\lambda\|^2 \in 2\mathbb{Z}$. ($\Rightarrow \langle \lambda, \lambda' \rangle \in \mathbb{Z}$)
• self-dual. $\Lambda^* := \{\lambda' \in \mathbb{R}^{24} \text{ s.t. } \langle \lambda, \lambda' \rangle \in \mathbb{Z} \ \forall \lambda \in \Lambda\} = \Lambda$
• Λ has no "holes" i.e. no vectors λ w/ $\|\lambda\|^2 = 2$.

• self-dual. $\Lambda^* := \{\lambda' \in \mathbb{R}^{24} \text{ s.t. } \langle \lambda, \lambda' \rangle \in \mathbb{Z} \ \forall \lambda \in \Lambda\} = \Lambda$

• Λ has no roots i.e. no vectors λ w/ $\|\lambda\|^2 = 2$.

• $O(\Lambda) \subseteq O(\mathbb{R}^{24}) = C_0 = 2C_1$ C_1 is sporadic simple.

Given even + self dual lattice, Λ of rank 24

$\Rightarrow$ holomorphic bosonic (1+1)d CFT V_Λ with $c_L = \text{rank} = 24$, $c_R = 0$.

$\sim$ EOM: holomorphic map from (1+1)d spacetime to $\mathbb{R}^{24}/\Lambda = T$ abelian WZW model.

↑ no right movers

Some manifest automorphisms of V_Λ

• $T = \mathbb{R}^{24}/\Lambda$ acts by translation;

• $O(\Lambda)$ acts by rotation;

∃ anomaly to lifting these classical symmetries to quantum ones.

for Leech, these are all the automorphisms

but for a general lattice, there are more

in particular: every root $\alpha \in \Lambda$ generates S^1s.

$\Rightarrow$ actually $\text{Aut}(V_\Lambda) \supseteq T \cdot O(\Lambda)$. non-split.

$$\#\mathbb{M} = 2^{46}\cdot 3^{20}\cdot 5^{9}\cdot 7^{6}\cdot 13^{3}\cdot \underbrace{11^{2}\cdot 17\cdot 19\cdot 23\cdot 29\cdot 31\cdot 41\cdot 47\cdot 59\cdot 71}_{\checkmark} \sim 10^{54}$$

remark: $\in \#SL_{13}(\mathbb{F}_2)$.

The primes $p \in \{2, 3, 5, 7, 13\}$ are special. (cf. Conway–Norton Monstrous moonshine conjectures)

- $\exists\, g \in O(\Lambda)$ of order p s.t. $g \subset \mathbb{R}^{24}$ has no (nonzero) fixed pts
  Unique up to conjugacy.
- $\mathbb{Z}_p \subseteq O(\Lambda)$ means this $\mathbb{Z}_p$ lifts in a unique-up-to-conjugacy way to $\mathbb{Z}_p \subseteq \mathrm{Aut}(V_\Lambda)$.

In fact, to a monstrous auto.


Piece rigidity
$\mathbb{Z}_{pB}$
$=$
$P \cdot P^{\alpha} \cdot J \subseteq \Pi$
new sym.
abelian calc
$\sim$
inspect known facts
about Π
$P \cdot P^{\alpha} \supset$
extra special
$P^{1+\alpha}$
$= H^2(P^{\alpha}, \mathbb{Z}_p)$


Choice of $\mathbb{Z}_p \subset V_\Lambda$ breaks $Aut(V_\Lambda) = T \cdot O(\Lambda)$
to the normalizer of $\mathbb{Z}_p$
$J = \frac{Norm(g \in O(\Lambda))}{\langle g \rangle}$
$(\mathbb{Z}_p$-fixed pt in $T) \cdot (\text{normalizer of } \mathbb{Z}_g \in O(\Lambda))$
P
P, J
Piece that fixes
Z_{FB}


new sym.
abelian calc
inspect known facts
extraspecial
classified by
$\mathrm{AH}^2(p^d) \subseteq H^2(p^d, \mathbb{Z}_p)$


The primes p=2, 3, 5, 7, 13 are special.
p-1 divides 24
d = 24/(p-1)
M2
1+d
J
2
p-Sylow.
's.
p
J
2
3
5
7
13
C0.
2S3 ≅ 2.
2 # J. 4
3 × 2S7
3 × 2S4


primes p ≤ 2, 3, 5, 7, 13 are special.
p-1 divides 24
d = 24/(p-1)
M2
(1+d)
J
≅ S.
p-Sylow.
p | J
2 | C0.
3 | 2S3 ≅ Z2.
5 | 2H.J.4
7 | 3×2S7
13 | 3×2S4
p = 13.
S = 13^(1+2) ⋊ (3×2S4)
J
Heisenberg gp
(1 * *; 1 *; 1) ⊆ SL3(F13).
≅ 13's.
H°(J; H(13^(1+2), U(1))) ⇒ H(S; U(1))
(J)


central
$13 \cdot 13^2$
E_2 page is
$H^0(J; H^3(13^{1+2}; \mathcal{U}(1))) \cong H^3(M; \mathcal{U}(1))_{(13)}$
$H^3(13^{1+2}, \mathcal{U}(1))$
$\cong Sym^2(13^2)$
$H^i(13^2; H^j(13; \mathcal{U}(1))) \Rightarrow H^{i+j}(13^{1+2})$
$\mathcal{U}(1)$
$\mathbb{Z}_{13}$
0
$\mathbb{Z}_{13}$


primes p ≤ 2, 3, 5, 7, 13 are special.
M2
p^{1+2} J
2
p-Sylow.
For p = 7, 5
E_2 page.
H^0(J; H^3(p^{1+2}))
H^1(J; H^2(p^{1+2}))
H^2(J; H^1)
H^3(J; Q(1))
p | J
2 | Co1
3 | 2Suz.2
5 | 2HJ.4
7 | 3×2S7
13 | 3×2S4


p=2, 3, 5, 7, 13
are special.
P^{1+2} ⋊ J
2 p-Sylow.
$E := P^2 = \Phi_P^2$
$H^1(P^{1+2}) = E^*$.
For p=7, 5
E_2 page. Calc.
$H^0(J; Sym^2(E^*) \otimes (\det)^{-1})$ by cultural clue.
$H^1(J; \Lambda^2(E^*))$ extension $= 0$ for p=5, 7.
$H^2(J; E^*) = 0$ by central char considerations.
$H^3(J; \mathbb{Q}(1)) = 0$
p=7 known. p=5 we had to work.
p | J
2 | C_3
3 | $2S_4(2)$.
5 | $2 \times J_4$
7 | $3 \times 2S_7$
13 | $3 \times 2S_4$

$\#SL_{13}(\mathbb{F}_2)$.

× × × × done, no H^3.

$p = 2, 3, 5, 7, 13$ are special.

$p^{1+2}:J$

p	J
2	C_3
3	$2S_{4,2,2}$
5	$2HJ.4$
7	$3 \times 2S_7$
13	$J \times 2S_4$

$E := p^2 = \mathbb{F}_p^2$

$H^1(p^{1+2}) = E^*$.

For $p = 7, 5$

E_2 page. $\to$ Calc.

$H^0(J; Sym^2(E^*) \otimes (\det)^{-1}) = 0$ by central char.

$H^1(J; \Lambda^2(E^*)) = 0$ for $p = 5, 7$.

$H^2(J; E^*) = 0$ by central char condition.

$H^3(J; \mathbb{Q}(1)) = 0$

$\mathbb{Z}$


At p=3:
$E^* = \pi_3^{12} \oplus 2S_{UZ.2}$
we need to know:
$H^0(2S_{UZ.2};\ Sym^2(E^*))$
$H^1(2S_{UZ.2};\ \Lambda^2(E^*))$
Sym. pr.
$H^3(2S_{UZ.2})_{(3)} = 0.$
$2S_{UZ.2}$
U1
$3^5 \times (M_{11} \times 2)$
U1
3-Sylow of $2S_{UZ.2}$


At p=3:
$E^* = TF_3^{12} \circlearrowleft 2S_{UZ.2.}$
we need to know:
0 easy $= H^0(2S_{UZ.2.}; Sym^2(E^*))$
$H^1(2S_{UZ.2.}; \overline{\Lambda^2(E^*)})$
Symp. Pr.
$H^3(2S_{UZ.2.})_{(3)} = 0.$
$2S_{UZ.2.}$
U1
$3^5 \times (M_{11} \times 2)$
U1
3-Sylow of $2S_{UZ.2.}$
$= H^1(3^5 \times (M_{11} \times 2); 1) = \mathbb{Z}_3.$


$H^3(M;U(1))_{(3)} \leq \mathbb{Z}_3$
enough to show $\neq 0$.
$\omega_3^5 \in H^3(M;U(1))$
'tHooft anomaly of $M \subset V$
$H^*(G;H(3)) \Rightarrow H^*(3G)$
V
G
12
$3 \times 6Sur$
both central extensions
$3 \cdot G$
36
$6Sur$


Learned: $H^3(M; U(1))_{(3)} \leq \mathbb{Z}_3$.
Claim: $= \mathbb{Z}_3$. enough to show $\neq 0$.
$V^{\uparrow}$ $\mathbb{Z}_3$ gauging V
$\uparrow G$
$3^{1+12} \cdot 2S_{U_2}$ $\leftrightarrow$ $3^{12} \rtimes 6S_{U_2}$ ← both central extensions
$3G$
$G = 3^{12} \rtimes 6S_{U_2}$ $3G$ $3 \cdot G$
$\omega_3^{\uparrow} \in H^3(M$
't Hooft
$H(G,$


E_2: $H^i(G; H^j(\mathbb{Z}_3, U(1))) \Rightarrow H^{i+j}(3G)$
j
3
2
1
0
$H^3(\mathbb{Z}_3)$
0
$H^1(\mathbb{Z}_3, \mathbb{Z}_3)$
$\mathbb{Z}_3$
$U(1)$
0
0
$H^2(G, \mathbb{Z}_3)$
$H^3(G, U(1))$
$H^4(G, U(1))$
0
1
2
3


$G = 3^{12} \rtimes SL_2$
$3G$
$K \in H^2(G; \mathbb{Z}_3)$ classifying extension.
$E_2: \quad H^i(G, H^j(\mathbb{Z}_3, U(1))) \Rightarrow H^{i+j}(3G)$
d_2 differential
$H^0(G; \hat{\mathbb{Z}}_3) \to H^{i+2}(G; U(1))$
is $\chi \mapsto \langle \alpha \cup K \rangle$
$H^3(\mathbb{Z}_3)$
$H^1(\mathbb{Z}_3, \mathbb{Z}_3)$
$\hat{\mathbb{Z}}_3$
$U(1)$
$H^2(G, \hat{\mathbb{Z}}_3)$
$H^3(G, U(1))$
$H^4(G, U(1))$
d_2
d_3
d_2


remark: $\in \#SL_{13}(\mathbb{F}_2)$.
done, no H^3.
$\omega^4 \in H^3(BG) = (\text{sub of } H^3(\mathbb{Z}_3)) \cdot (\ker d_2: H^2(G;\hat{\mathbb{Z}}_3) \to H^4(G)) \cdot (\text{coker } d_2: H^1(G;\hat{\mathbb{Z}}_3) \to H^3(G))$
"β"
A cocycle $\omega \in H^3(BG)$ s.t. $\omega|_{\mathbb{Z}_3}$ 2-cocycle
α 2-cocycle coeff in $\hat{\mathbb{Z}}_3$.
β solving $d\beta = \langle \alpha \cup k \rangle$ 3-cochain
$\omega^4|_{\mathbb{Z}_3} = 0$ from $\mathbb{Z}_3$ 3-manifold
So ω^4 lives here
2-cocycle
$\alpha \in H^2(G;\hat{\mathbb{Z}}_3)$ image of ω^4.
$(d\alpha = 0.)$ also $d_2\alpha = \langle \alpha \cup k \rangle = 0$ in $H^4(G)$
3 cochain β $d\beta = \langle \alpha \cup k \rangle$


Learned: $H^3(M;U(1))_{(3)} \leq \mathbb{Z}_3$
Claim: $= \mathbb{Z}_3$.
enough to show $\neq 0$.
$\omega = (\alpha, \beta)$
$\omega = (\ast, \beta)$
$\omega \in H^3(M;U(1))$
$\alpha \neq 0$.
$G = 3^{1+12} \cdot 2_{+}^{1+12}$
$3^{12} \times 6S_{12}$
$G = 3^{12} \times 6S_{12}$
$K \in H^2(G, \mathbb{Z}_3)$ classifying extension
$H^i(G, H(\mathbb{Z}_3)) \Rightarrow H^i(3G)$


$\#\mathcal{M} = 2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 13^3 \cdot 11^2 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41$
$H^3(\mathcal{M}; U(1))_{(3)} = \mathbb{Z}_3$.
done, no H^3.
$\omega^4 \in H^3(BG) = (\text{sub of } H^3(\mathbb{Z}_3)) \cdot (\ker d_2 : H^2(G; \hat{\mathbb{Z}}_3) \to H^4(G))$
A cocycle $\omega^4 \in H^3(BG)$ s.t. $\omega|_{\mathbb{Z}_3}$ 2-cocycle
$\omega^4|_{\mathbb{Z}_3} = 0$
α 2-cocycle exists in $\hat{\mathbb{Z}}_3$.
β solving $d\beta = \langle \alpha \cup K \rangle$
3-cochain
So ω^4 lives
2 cocycle
$\alpha \in H^2(G; \hat{\mathbb{Z}}_3)$
$(d\alpha = 0$, also
3 cochain
$-\beta$
$d\beta =$


$H^3(M; U(1))_{(3)} = \mathbb{Z}_3$.
done, no H^3.
$p=2$.
$H^i(2^{1+24}, \wedge^i$
$J=3$ $2^{2300} \cdot 2 \leq \mathbb{Z}_4$
exp=4
$J=2$ $\mathbb{Z}_2$
me. but free w/ matrices.
$J=1$ 2^{24} $H^1(G_1, 2^{24}) = 0$ $H^2(G_1, 2^{24}) = \mathbb{Z}_2$
Leech Hut
$J=0$ $U(1)$ $H^1(G) = *$ $H^2(G_1, U(1)) = \mathbb{Z}_2$ $H^3(G_1)_{(2)} = \mathbb{Z}_4$ Tw w/ DT.
$H^i(G_1, -)$


$\omega^4 \in H^3(M, U(1))$ use for lower bound.
ω^4
$M^3 = S^3/\Gamma$ $\Gamma = 2S_{U_2} \subset 6S_{U_2} = 2A_4 \subset 2D_8$
restricts to a class here which is not 0 in H^3
next
order 8: $\mathbb{Z}'' \in \mathbb{Z}_{16}$
$S^3 \to S^3/2D_8$
3-oriented manifold
M^3
Meng told you $H^*(D_8)$
$H(2D_8, \mathbb{Z})$
$H(2D_8, \mathbb{Z})$
$2D_8 \subset SU(2) = S^3$
$\Rightarrow H^3(2D_8, U(1))$
$D_8 \subset O(2) \subset SO(3)$
$H(2D_8, H(S^3)) \Rightarrow H^*(M^3)$
$H^1(\cdot, \mathbb{Z}) = \mathbb{Z}/16$.


M = Fischer-Griess monster is the largest sporadic simple gp.
#M = 2^46 · 3^20 · 5^9 · 7^6 · 13^3 · 11^2 · 17 · 19 · 23 · 29 · 31 · 41 · 47 · 59 · 71 ~ 10^54
H^3(M; U(1)) = Z_8 ⊕ Z_3 ⊕ Z_3
H^3(M; U(1))_(3) = Z_3
done, no H^3
exp=4
me: brute force w/ matrices!
Deloch Hat
is pulled back from Co_1
Co_1 not DT
⇒ 8ω^3 = 0
E_2^{s,t} = H^s(Co_1; H^t(Z^{1+24}))
t=2
t=1
t=0
U(1)
H^0(Co_1, Z^24) = 0
H^1(Co_1, Z^24) = Z_2
H^0(Co_1) = *
simple.
H^1(Co_1) = *
H^2(Co_1; U(1)) = Z_2
ext to Co_0
H^3(Co_1)_(2) = Z_4
H^4(Co_1, -)
Thm w/ DT.

1707.08388 + 1810.00463
Z_2 w/ D. Treumann.

Goal: Thm. $H^3(\mathbb{M}, U(1)) \cong \mathbb{Z}_{24} \oplus (\underbrace{\leq 4}_{\text{probably trivial}})$

$\mathbb{M}$ = Fischer-Griess monster is the largest sporadic simple gp.

$$\#\mathbb{M} = 2^{46}\cdot 3^{20}\cdot 5^9\cdot 7^6\cdot 13^3\cdot \underbrace{11^2\cdot 17\cdot 19\cdot 23\cdot 29\cdot 31\cdot 41\cdot 47\cdot 59\cdot 71}_{\text{done, no } H^3} \sim 10^{54}$$

$H^3(\mathbb{M}; U(1)) = \mathbb{Z}_8 \oplus$ [illegible] $H^3(\mathbb{M}; U(1))_{(3)} = \mathbb{Z}_3$.